

CAFPE-121/09  
UF-FT-251/09

## Standard Model Prediction of the Muon Anomalous Magnetic Moment <sup>1</sup>

**Joaquim Prades**

CAFPE and Departamento de Física Teórica y del Cosmos, Universidad  
de Granada, Campus de Fuente Nueva, E-18002 Granada, Spain.

I review the present Standard Model prediction of the muon anomalous magnetic moment. The discrepancy with its experimental determination is  $(25.5 \pm 8.0) \times 10^{-10}$ , i.e., 3.2 standard deviations.

September 2009

---

<sup>1</sup> Invited talk given by J.P. at Topical FLAVIANet Workshop on “Low Energy Constraints on Extensions of the Standard Model”, July 24-26 2009, Kazimierz, Poland.

## 1. Introduction

The general vertex  $\Gamma_\mu$  between a fermion  $f$  and an external electromagnetic (EM) field  $A^\mu(q = p - p')$  can be written as

$$\bar{u}(p')\Gamma_\mu u(p) = \bar{u}(p') \left[ \gamma_\mu F_1(q^2) + i \frac{\sigma_{\mu\nu} q^\nu}{2m_f} F_2(q^2) + \dots \right] \quad (1)$$

The magnetic dipole moment for a charged fermion ( $f = e, \mu, \tau, \dots$ ) is proportional to the spin  $\vec{s}$  through the gyromagnetic factor  $g_f \equiv 2(F_1(0) + F_2(0))$

$$\vec{\mu} = g_f \frac{e}{2m_f} \vec{s}. \quad (2)$$

The Dirac vertex predicts  $F_1(0) = 1$  and  $F_2(0) = 0$  at tree-level, quantum loops modify this prediction to  $F_2(0) \equiv a_f \neq 0$  while  $F_1(0)$  is a conserved charge. The quantity  $a_f = (g_f - 2)/2$  is commonly called the fermion anomaly.

The muon anomalous magnetic moment  $a_\mu$  has been measured by the E821 experiment (Muon g-2 Collaboration) at BNL with an impressive accuracy of 0.72 ppm [1] yielding the present world average

$$a_\mu^{\text{exp}} = (11, 659, 208.9 \pm 6.3) \times 10^{-10} \quad (3)$$

with an accuracy of 0.54 ppm. New experiments [2,3] are being designed to measure  $a_\mu$  with an accuracy better than 0.14 ppm. Here, I'm interested in what is the present status of the Standard Model (SM) prediction for this very precise measurement. Is there room for new physics in  $a_\mu$ ? Recent reviews can be found in [4–9].

First, I will shortly recall recent advances in the electron anomaly  $a_e$  which is a necessary ingredient to predict the muon anomaly  $a_\mu$ . I'll discuss then briefly the main contribution to  $a_e$  that is QED, and the much smaller hadronic and weak contributions. Secondly, I'll discuss briefly the main contribution to  $a_\mu$  that is QED and the smaller weak contribution. Then I discuss the also smaller but dominant in the uncertainty hadronic contribution to  $a_\mu$ . Finally, I'll give the conclusions and prospects for improving  $a_\mu$  both theoretic and experimentally.

## 2. The electron anomaly $a_e$

The Harvard group [10] has very recently made a very precise measurement of the electron anomaly

$$a_e^{\text{exp}} = (11, 596, 521, 807.3 \pm 2.8) \times 10^{-13} \quad (4)$$

improving the 1987 measurement of the University of Washington group [11]

$$a_e^{\text{exp}} = (11, 596, 521, 883 \pm 42) \times 10^{-13} \quad (5)$$

by more than a factor 15.

The new  $a_e$  measurement provides the most accurate value for the fine structure constant <sup>1</sup>

$$\alpha^{-1} = 137.035\,999\,084(33)(39) = 137.035\,999\,084(51) \quad (6)$$

which is one order of magnitude more precise than the best previous determination<sup>2</sup>. The best non- $a_e$  based determinations of  $\alpha$  come from atomic physics, and in particular from the precise measurement of Cs [16] and Rb [17] atomic masses,

$$\begin{aligned} \text{Cs atom (2002)} : \alpha^{-1} &= 137.036\,000\,000(1\,100) \\ \text{Rb atom (2008)} : \alpha^{-1} &= 137.035\,999\,450(620). \end{aligned} \quad (7)$$

It is expected that the Rb atomic mass measurement can reach the level of accuracy of  $a_e$  determinations soon. At present,  $a_e$  together from  $\alpha$  from Cs and Rb atomic mass determinations checks QED at 3-loops ! If both determinations were at the same level of accuracy, it would check QED at 4-loops and a possible substructure of the electron.

### 3. Standard Model Contributions to $a_e$

The dominant SM contribution comes from QED

$$a_e^{\text{QED}} = \sum_{n=1} C_{2n} \left( \frac{\alpha}{\pi} \right)^n \quad (8)$$

with  $C_2 = 1/2$  calculated by Schwinger [18] 61 years ago; one can find the values of the rest of the coefficients in different reviews [4–9]. The coefficients  $C_4$  [19] and  $C_6$  [20] are known analytically including lepton mass corrections.  $C_8$  is only numerically known and being continuously improved since early 1980s [21]. Actually, it was corrected recently in 2007 [13]. For  $C_{10}$ , one uses  $0.0 \pm 4.6$  as an estimate where the error is based on the  $C_{2n}$  values growing. Its full numerical calculation is in progress, it

<sup>1</sup> For a very recent discussion of the extraction of  $\alpha$  from  $a_e$  and other methods, see [12].

<sup>2</sup> Notice that due to the 2007 corrected value for the four-loop order QED coefficient [13], the value quoted in both PDG 2008 [14] and CODATA 2006 [15]  $\alpha^{-1} = 137.035\,999\,679(94)$  is not correct.

involves 12,672 diagrams and some partial results are already available [22]. Recently, specific classes of the QED eighth and tenth orders have been calculated analytically using Mellin-Barnes transform techniques [23] and agree with the numerical results in [13, 21, 22].

There are other much smaller contributions to  $a_e$  but that at the level of present precision start to be needed. The largest is the leading order hadronic vacuum polarization contribution (see Fig 1) [24]

$$a_e^{\text{LOHadronic}} = (18.75 \pm 0.18) \times 10^{-13}. \quad (9)$$

while higher order hadronic vacuum polarization contribution is [25]

$$a_e^{\text{HOHadronic}} = -(2.25 \pm 0.05) \times 10^{-13} \quad (10)$$

and the hadronic light-by-light contribution (see Fig 2) is [26]

$$a_e^{\text{HadronicLbL}} = (0.35 \pm 0.10) \times 10^{-13}. \quad (11)$$

The total hadronic contribution to  $a_e$  is

$$a_e^{\text{Hadronic}} = (16.85 \pm 0.21) \times 10^{-13}. \quad (12)$$

Notice that this is the first time that  $a_e^{\text{exp}}$  is sensitive to the hadronic contribution. The electroweak contribution is much smaller [5]

$$a_e^{\text{EW}} = (0.385 \pm 0.004) \times 10^{-13}. \quad (13)$$

## 4. Standard Model Contributions to $a_\mu$

### 4.1. QED

A precise value of the fine structure constant is very important to the determination of the muon anomaly since QED is again the dominant SM contribution. One can write the QED contribution as

$$a_\mu^{\text{QED}} = \sum_{n=1} C_{2n} \left( \frac{\alpha}{\pi} \right)^n \quad (14)$$

and the same comments and authors can be quoted for the  $C_{2n}$  coefficients that can be found in the reviews [4–9]. For  $C_{10}$ , one uses  $663 \pm 20$  which is also estimate and its calculation is progress in parallel to  $C_{10}$  for  $a_e$  [22]. Using those coefficients and  $\alpha$  from the latest  $a_e$  in (6), one gets

$$a_\mu^{\text{QED}} = (11, 658, 471.810 \pm 0.015) \times 10^{-10} \quad (15)$$

where the largest uncertainty comes from  $C_{10}$ . Given that precision, the difference between the experimental value (3) and its QED prediction

$$\Delta_{\text{Not QED}}^{\text{exp}} = (736.2 \pm 6.3) \times 10^{-10} \quad (16)$$

can be considered as a “measurement”. In fact,  $a_\mu$  has been sensitive to the hadronic contribution since 1975 and to the electroweak ones since 2001.




Fig. 1. Lowest order hadronic vacuum polarization contribution to muon  $g - 2$ .

#### 4.2. Electroweak

The one loop electroweak contribution is fully known since 1972 [28]

$$\begin{aligned} a_{\mu}^{\text{EW one-loop}} &= \frac{5G_{\mu}m^2}{24\sqrt{2}\pi^2} \left[ 1 + \frac{1}{5} \left( 1 - 4\sin^2(\theta_W) \right)^2 + \mathcal{O}(m/M_{Z,W,H}) \right] \\ &= (19.5 \pm 0.2) \times 10^{-10}. \end{aligned} \quad (17)$$

The full two-loops electroweak contribution has been finished much more recently (in 2004) though its potential importance was already pointed out in 1992 [29]. The full result can be found in [30].

The final result for the two-loop electroweak contribution [27, 30] is  $-(4.1 \pm 0.1) \times 10^{-10}$  and the total electroweak contribution

$$a_{\mu}^{\text{EW}} = [(19.5 \pm 0.2) - (4.1 \pm 0.1)] \times 10^{-10} = (15.4 \pm 0.2) \times 10^{-10}. \quad (18)$$

Again, given the experimental precision, the difference between the precise experimental value in (3) and its QED plus EW prediction

$$\Delta_{\text{Not QED+EW}}^{\text{exp}} = (720.8 \pm 6.3) \times 10^{-10} \quad (19)$$

can be considered as a “measurement”. This also tells us that we need to know the hadronic contribution to  $a_{\mu}$  with an accuracy below the 1 % level!

#### 4.3. Hadronic Contributions

##### 4.3.1. Vacuum Polarization Contribution

This contribution is depicted in Fig. 1. Its leading-order (LO) contribution is order  $\alpha^2$  and can be written in terms of the one-photon  $e^+e^- \rightarrow \gamma^* \rightarrow \text{hadrons}$  cross-section,  $\sigma^{(0)}(s)$  [31].

$$a_{\mu}^{\text{LO Hadronic}} = \frac{1}{4\pi^3} \int_{4m_{\pi}^2}^{\infty} ds K(s) \sigma^{(0)}(s) \quad (20)$$

with

$$K(s) = \int_0^1 dx \frac{x^2(1-x)}{x^2 + \frac{s}{m^2}(1-x)}. \quad (21)$$

The kernel  $K(s)$  behaves as  $m^2/s$  making the low-energy region ( $\rho$ -region) dominate in the integrand. In fact, the  $\pi^+\pi^-$  channel below 1 GeV gives 72 % of the total of  $a_\mu^{\text{LO Hadronic}}$ .

The one-photon cross-section  $\sigma^{(0)}(s)$  is (almost) experimentally obtained. One has to correct the experimental cross-section to include final state photons and exclude the running of  $\alpha(s)$  not to make double counting with higher orders. It is also welcome to eliminate systematics to have the cross-section  $\sigma^{(0)}$  normalized to the  $e^+e^- \rightarrow \mu^+\mu^-$  cross-section measured by the same experiment. In the case of the initial state radiation (ISR) method, this is done by BaBar which measures  $e^+e^- \rightarrow \pi^+\pi^-\gamma(\gamma)$  normalized to  $e^+e^- \rightarrow \mu^+\mu^-\gamma(\gamma)$  cross-section measured by the same experiment and is planned to be done in the forthcoming KLOE-2 measurement.

For the low energy contribution to  $\sigma^{(0)}$ , we have very precise  $e^+e^-$  data from several low-energy experiments, namely, from CMD-2 and SND detectors at VEPP-2M (Novosibirsk), KLOE at DAΦNE (Frascati) and BaBar at PEP-II (SLAC). The last results from CMD-2 [32, 33] and SND [34] are in nice agreement. Final KLOE data [35] are at the same level of accuracy as CMD-2 and SND data and there is an overall agreement at energies between 0.630 and 0.958 GeV with CMD-2 and SND though KLOE data lies somewhat lower. Very recently, BaBar using the ISR method has also released  $e^+e^-$  results for the dominant  $\pi^+\pi^-$  at energies between 0.3 GeV and 3 GeV [36]. Belle at KEK (Tsukuba) which has larger statistics, will do so soon.

At high energy, higher than 13 GeV, and in between  $\bar{c}c$  and  $\bar{b}b$  thresholds, perturbative QCD is used to calculate  $\sigma^{(0)}(s)$ .

Combining all precision  $e^+e^-$  data, including the latest BaBar data [36], the authors [37] quote

$$a_\mu^{\text{LO Hadronic}} = (695.5 \pm 4.0_{\text{exp}} \pm 0.7_{\text{QCD}}) \times 10^{-10} = (695.5 \pm 4.1) \times 10^{-10}. \quad (22)$$

A new analysis of isospin corrections to  $\tau^+ \rightarrow \pi^+\pi^0\nu$  data when compared to the the CVC  $e^+e^- \rightarrow \pi^+\pi^-$  data has been also recently released [38]. The discrepancy between  $\tau$  data and the combined  $e^+e^-$  data for the dominant  $\pi^+\pi^-$  contribution to  $\sigma^{(0)}$  of the measurements quoted above is reduced to  $1.5 \sigma$  from the previous  $4.5 \sigma$  discrepancy [37, 38].

There are also hadronic vacuum polarization contributions at higher order included in Fig 1. The most recent evaluation including  $\alpha^3$  corrections




Fig. 2. Hadronic light-by-light contribution to  $a_\mu$  is in Fig. 2.

is in [39] using also  $e^+e^-$  data,

$$a_\mu^{\text{HO Hadronic}} = -(9.79 \pm 0.08_{\text{exp}} \pm 0.03_{\text{rad}}) \times 10^{-10} = -(9.79 \pm 0.09) \times 10^{-10}.$$

Where the uncertainty also takes into account non-included radiative corrections [39].

#### 4.3.2. Light-by-Light Contribution

One of the six possible momenta routing to the hadronic light-by-light contribution to  $a_\mu$  is in Fig. 2. Recent work on  $a_\mu^{\text{HLbL}}$  can be found in [26, 40–47]. For reviews on the status, prospects of  $a_\mu^{\text{HLbL}}$  and references see [4, 5, 48]. In particular, the discussion in [26] lead the authors to give the following value

$$a_\mu^{\text{Hadronic LbL}} = (10.5 \pm 2.6) \times 10^{-10} \quad (23)$$

for their present best estimate.

### 5. Result for $a_\mu^{\text{SM}}$

Adding all pieces discussed above contributing to  $a_\mu$ , one gets

$$\begin{aligned} 10^{10} a_\mu^{\text{SM}} &= \overbrace{(11, 658, 471.810 \pm 0.015)}^{\text{QED}} + \overbrace{(15.4 \pm 0.2)}^{\text{EW}} \\ &+ \underbrace{(695.5 \pm 4.1)}_{\text{LO Had}} - \underbrace{(9.79 \pm 0.09)}_{\text{HO Had}} + \underbrace{(10.5 \pm 2.6)}_{\text{Had LbL}} \\ &= \underbrace{(11, 658, 487.2 \pm 0.2)}_{\text{QED+EW}} + \underbrace{(696.2 \pm 4.9)}_{\text{Hadronic}} \\ &= 11, 659, 183.4 \pm 4.9. \end{aligned} \quad (24)$$

Using this result and the experimental value of  $a_\mu$  in (3), one gets

$$a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (25.5 \pm 8.0) \times 10^{-10}. \quad (25)$$

## 6. Conclusions and Prospects

Combining all recent precise  $e^+e^-$  data obtained with different methods (energy scanning and ISR) in different experiments to calculate  $\sigma^{(0)}$  [37] and using the new evaluation of the hadronic light-by-light contribution in [26, 48] one gets more than  $3\sigma$  of discrepancy between the SM value for  $a_\mu$  (24) and its experimental determination (3). This discrepancy has been slowly growing due to impressive theory and experiment recent achievements. In fact, both theory and experiment uncertainties have been reduced by more than a factor two in the last eight years <sup>3</sup>.

There are planned new  $e^+e^-$  experiments at Novosibirsk (VEPP-2000) and Frascati (DAΦNE-2) which will cross-check present results and reduce the present uncertainty in  $a_\mu^{\text{LOHad}}$  which dominates the final SM uncertainty now. There are also new  $\tau$  data at B-factories and a new  $\tau$ -charm factory at Beijing which will cross-check the  $\tau$  result. It will also help to understand better the isospin violation corrections in order to use the forthcoming precise  $\tau$  data. In addition, a new full calculation of  $a_\mu^{\text{HadLbL}}$  is desirable and possible. The goal in this case is to reduce its present uncertainty to the level of  $(1.5 \sim 2.0) \times 10^{-10}$ . With all these eventual theory improvements the uncertainty of the SM prediction can be reduced soon enough for the new experiments being designed. As said before, these experiments are being designed to measure  $a_\mu$  with an accuracy better than 0.14 ppm in parallel to the expected theory advances. The proposed experiment at Fermilab is designed to reduce the  $a_\mu$  uncertainty to the level of  $1.6 \times 10^{-10}$  [2] and the one at J-PARC to somewhere between  $1.2 \times 10^{-10}$  and  $0.6 \times 10^{-10}$  [3].

With all the theory activity detailed above and the tantalizing more than  $3\sigma$  discrepancy in  $a_\mu$  (25) between the SM prediction and its experimental determination, I believe that those new  $g-2$  experiments are very timely, necessarily complementary to direct searches like LHC and ILC and should be done as soon as possible. Its very likely that they give the first new physics discovery !

## Acknowledgments

This work is supported in part by the European Commission (EC) RTN network, Contract No. MRTN-CT-2006-035482 (FLAVIANet), by the Spanish Consolider-Ingenio 2010 Programme CPAN Grant No. CSD2007-00042, MEC (Spain) and FEDER (EC) Grant No. FPA2006-05294, Junta de Andalucía Grant Nos. P05-FQM 437 and P07-FQM 03048.

---

<sup>3</sup> In 2001 , this discrepancy was  $a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (23.1 \pm 16.9) \times 10^{-10}$ . [49]

## REFERENCES

- [1] B.L. Roberts, arXiv:1001.2898; G.W. Bennett *et al.* [Muon  $g-2$  Collaboration], Phys. Rev. D **73** (2006) 072003; Phys. Rev. Lett. **92** (2004) 161802.
- [2] D.W. Hertzog, Nucl. Phys. B (Proc. Suppl.) **181-182** (2008) 5; D.W. Hertzog, *et al* arXiv:0705.4617; B.L. Roberts [E821 and E969 Collaborations], Nucl. Phys. B (Proc. Suppl.) **155** (2006) 372; hep-ex/0501012.
- [3] J. Imazato, Nucl. Phys. B (Proc. Suppl.) **129-130** (2004) 81; “Letter of Intent: An Improved Muon ( $g-2$ ) Experiment at J-PARC”, R.M. Carey *et al.* (2003).
- [4] E. de Rafael, Nucl. Phys. B (Proc. Suppl.) **186** (2009) 211; PoS (**EFT09**) 050.
- [5] F. Jegerlehner and A. Nyffeler, Phys. Rept. **477** (2009) 1.
- [6] F. Jegerlehner, Lect. Notes Phys. **745** (2008) 9; Acta Phys. Polon. B **38** (2007) 3021.
- [7] J.P. Miller, E. de Rafael and B.L. Roberts, Rept. Prog. Phys. **70** (2007) 795.
- [8] M. Passera, J. Phys. G **31** (2005) R75.
- [9] M. Knecht, Lect. Notes Phys. **629** (2004) 37.
- [10] D. Hanneke, S. Fogwell, and G. Gabrielse, Phys. Rev. Lett. **100** (2008) 120801; B. Odom, *et al*, *ibid.* **97** (2006) 030801.
- [11] R.S. Van Dyck, Jr., P.B. Schwinberg, and H.G. Dehmelt, Phys. Rev. Lett. **59** (1987) 26.
- [12] G. Gabrielse, in “Lepton Dipole Moments”, B.L. Roberts and W.J. Marciano (eds) (World Scientific, Singapore, 2009), Advanced Series on Directions in High Energy Physics, Vol. 20.
- [13] T. Aoyama *et al.* Phys. Rev. D **77** (2008) 053012.
- [14] C. Amsler, *et al*, Phys. Lett. B **667** (2008) 1.
- [15] P.J. Mohr, B.N. Taylor and D.B. Newell, Rev. Mod. Phys. **80** (2008) 633.
- [16] A. Wicht, *et al*, Phys. Scripta **T102** (2002) 82.
- [17] M. Cadoret, *et al*, Phys. Rev. Lett. **101** (2008) 230801.
- [18] J.S. Schwinger, Phys. Rev. **73** (1948) 416.
- [19] C.M. Sommerfield, Phys. Rev. **107** (1957); Ann. Phys. (N.Y.) **5** (1958) 677; A. Petermann, Hel. Phys. Acta **30** 407; Nucl. Phys. **5** (1958) 677; H. Suura and E. Wichmann, Phys. Rev. **105** (1957) 1930; A. Petermann, *ibid.* **105** (1957) 1931; H.H. Elend, Phys. Lett. **20** (1966) 682; [Errat-*ibid* **21** (1966) 720.]
- [20] J.A. Mignaco and E. Remiddi, Nuovo Cim. A **60** (1969) 519 R. Barbieri and E. Remiddi, Phys. Lett. B **49** (1974) 468; Nucl. Phys. B **90** (1975) 233; R. Barbieri, M. Caffo and E. Remiddi, Phys. Lett. B **57** (1975) 460; M. Levine, E. Remiddi and R. Roskies, Phys. Rev. D **20** (1979) 2068; S. Laporta and E. Remiddi, Phys. Lett. B **265** (1991) 182; *ibid.* **301** (1993) 440; *ibid.* **356** (1995) 390; *ibid.* **379** (1996) 283; S. Laporta, Phys. Rev. D **47** (1993) 4793; Phys. Lett. B **343** (1995) 421; Nuovo Cim. A **106** (1996) 675; T. Kinoshita, Nuovo Cim. B **51** (1967) 140; B.E. Lautrup and E. de Rafael, Phys. Rev. **174** (1968) 1835; B.E. Lautrup and M.A. Samuel, Phys. Lett. B **72** (1977) 114; M.A. Samuel and

- G. Li, Phys. Rev. D **44** (1991) 3935; [Errat-ibid **48** (1993) 1879]; A. Czarnecki and M. Skrzypek, Phys. Lett. B **449** (1999) 354.
- [21] T. Kinoshita and W.B. Lindquist. Phys. Rev. Lett. **47** (1981) 1573; Phys. Rev D **271** (1983) 867; *ibid.* **27** (1983) 877; *ibid.* **27** (1983) 886; *ibid.* **39** (1989) 2407; *ibid.* **42** (1990) 636; T. Kinoshita, B. Nizic and Y. Okamoto, *ibid.* **41** (1990) 593; *ibid.* **52** (1984) 717.
- [22] T. Aoyama, *et al.*, Phys. Rev. D **78** (2008) 113006; *ibid.* **78** (2008) 053005.
- [23] J.P. Aguilar, D. Greynat and E. de Rafael, Phys. Rev. D **77** (2008) 093010.
- [24] M. Davier and A. Höcker, Phys. Lett. B **435** (1998) 427.
- [25] B. Krause, Phys. Lett. B **390** (1997) 392.
- [26] J. Prades, E. de Rafael and A. Vainshtein, arXiv:0901.0306 in “Lepton Dipole Moments”, B.L. Roberts and W.J. Marciano (eds) (World Scientific, Singapore, 2009), Advanced Series on Directions in High Energy Physics, Vol. 20.
- [27] A. Czarnecki, B. Krause and W.J. Marciano, Phys. Rev. Lett. **76** (1996) 3267.
- [28] R. Jackiw and S. Weinberg, Phys. Rev. D **5** (1972) 2396; I. Bars and M. Yoshimura, *ibid.* **6** (1972) 374; K. Fujikawa, B.W. Lee and A.I. Sanda, *ibid.* **6** (1972) 2923; G. Altarelli, N. Cabibbo and L. Maiani, Phys. Lett. B **40** (1972) 415; W.A. Bardeen, R. Gastmans and B. Lautrup, Nucl. Phys. B **46** (1972) 319.
- [29] T.V. Kukhto *et al.*, Nucl. Phys. B **371** (1992) 567.
- [30] A. Czarnecki, B. Krause and W.J. Marciano, Phys. Rev. D **52** (1995) 2619; T. Kaneko and N. Nakazawa, arXiv:hep-ph/9505278; S. Peris, M. Perrottet and E. de Rafael, Phys. Lett. B **355** (1995) 523; G. Degrossi and G.F. Giudice, Phys. Rev. D **58** (1998) 053007; M. Knecht *et al.*, JHEP **11** (2002) 003; A. Czarnecki, W.J. Marciano and A. Vainshtein, Phys. Rev D **67** (2003) 073006; Acta. Phys. Polon. B **34** (2003) 5669; S. Heinemeyer, D. Stöckinger and G. Weiglein, Nucl. Phys. B **699** (2004) 103.
- [31] C. Bouchiat and L. Michel, J. Phys. Radium **22** (1961) 121; L. Durand, Phys. Rev. **128** (1962) 441; [Erratum-*ibid* **129** (1963) 2835]; S.J. Brodsky and E. de Rafael, Phys. Rev. **168** (1968) 1620; M. Gourdin and E. de Rafael, Nucl. Phys. B **10** (1969) 667.
- [32] R.R. Akhmetshin *et al.*, [CMD2 Collaboration], Phys. Lett. B **648** (2007) 28; Phys. Lett. B **466** (1999) 392; V.M. Aulchenko *et al.*, [CMD2 Collaboration], JETP Lett. **84** (2006) 413; *ibid.* **82** (2005) 743.
- [33] I.B. Logashenko, Nucl. Phys. B (Proc. Suppl.) **162** (2006) 13.
- [34] M.N. Achasov *et al.*, Nucl. Phys. B (Proc. Suppl.) **162** (2006) 11; JETP Lett. **103** (2006) 103.
- [35] F. Ambrosino *et al.* [KLOE Collaboration], Phys. Lett. B **670** (2009) 285.
- [36] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **103** (2009) 231801.
- [37] M. Davier *et al.*, arXiv:0908.4300; M. Davier *et al.*, [BABAR Collaboration], Nucl. Phys. B (Proc. Suppl.) **189** (2009) 222.
- [38] M. Davier *et al.*, arXiv:0906.5443.
- [39] K. Hagiwara, *et al.*, Phys. Lett. B **649** (2007) 173.

- [40] M. Hayakawa and T. Kinoshita, Phys. Rev. D **57** (1998) 465 [Erratum-ibid. D **66** (2002) 019902]; M. Hayakawa, T. Kinoshita and A.I. Sanda, Phys. Rev. Lett. **75** (1995) 790.
- [41] J. Bijnens, E. Pallante and J. Prades, Nucl. Phys. B **474** (1996) 379; Phys. Rev. Lett. **75** (1995) 1447 [Erratum-ibid. **75** (1995) 3781]; Nucl. Phys. B **626** (2002) 410.
- [42] J. Bijnens and F. Persson, Lund University Master Thesis, hep-ph/0106130.
- [43] M. Knecht and A. Nyffeler, Phys. Rev. D **65** (2002) 073034; M. Knecht *et al*, Phys. Rev. Lett. **88** (2002) 071802.
- [44] K. Melnikov and A. Vainshtein, Phys. Rev. D **70** (2004) 113006.
- [45] A.E. Dorokhov and W. Broniowski, Phys. Rev. D **78** (2008) 073011.
- [46] A. Nyffeler, Phys. Rev. D **79** (2009) 073012
- [47] T. Blum and S. Chowdhury, Nucl. Phys. B (Proc. Suppl.) **189** (2009) 251; M. Hayakawa, *et al*, PoS **LAT2005** (2006) 353.
- [48] J. Prades, PoS (**CD09**) 079; arXiv:0907.2938; arXiv:0905.3164 (to be published in Eur. Phys. J C); Nucl. Phys. B (Proc. Suppl.) **181-182** (2008) 15; J. Bijnens and J. Prades, Mod. Phys. Lett. A **22** (2007) 767; Acta Phys. Polon. B **38** (2007) 2819.
- [49] J. Prades, arXiv:hep-ph/0108192.