



(51) International Patent Classification:

C07D 237/16 (2006.01) A61P 35/00 (2006.01)
C07D 237/20 (2006.01) A61K 31/501 (2006.01)

(21) International Application Number:

PCT/EP2018/081545

(22) International Filing Date:

16 November 2018 (16.11.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

17202871.4 21 November 2017 (21.11.2017) EP

(71) Applicants: **BAYER AKTIENGESELLSCHAFT** [DE/DE]; Kaiser-Wilhelm-Allee 1, 51373 Leverkusen (DE). **BAYER PHARMA AKTIENGESELLSCHAFT** [DE/DE]; Müllerstr. 178, 13353 Berlin (DE). **DEUTSCHES KREBSFORSCHUNGSZENTRUM** [DE/DE]; Im Neuenheimer Feld 280, 69120 Heidelberg (DE).

(72) Inventors: **ZORN, Ludwig**; Osianderweg 45a, 13509 Berlin (DE). **RÖHN, Ulrike**; Boyenstr. 42, 10115 Berlin (DE). **GUTCHER, Ilona**; Paul-Robeson-Str. 28, 10439 Berlin (DE). **RÖSE, Lars**; Fuchsring 82, 13465 Berlin (DE). **BADER, Benjamin**; Hillmannstr. 14, 13467 Berlin (DE). **KOBER, Christina**; Heidelberger Str.18, 76646 Bruchsal (DE). **CARRETERO, Rafael**; Mühlingstraße 6, 69121 Heidelberg (DE). **STÖCKIGT, Detlef**; Clara-Zetkin-Strasse 27, 14471 Potsdam (DE). **PLATTEN, Michael**; Cambridgestr. 7, 69115 Heidelberg (DE).

(74) Agent: **BIP PATENTS**; Alfred-Nobel-Str. 10, 40789 Monheim am Rhein NRW (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,

SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))

(54) Title: SULPHUR SUBSTITUTED 3-OXO-2,3-DIHYDROPYRIDAZINE-4-CARBOXAMIDES



(57) Abstract: The present invention covers sulphur substituted 3-oxo-2,3-dihydropyridazine-4-carboxamide compounds of general formula (I): (I) in which R¹, R², A, X, Y and Z are as defined herein, methods of preparing said compounds, intermediate compounds useful for preparing said compounds, pharmaceutical compositions and combinations comprising said compounds and the use of said compounds for manufacturing pharmaceutical compositions for the treatment or prophylaxis of diseases, in particular of cancer or conditions with 10 dysregulated immune responses or other disorders associated with aberrant AHR signaling, as a sole agent or in combination with other active ingredients.



SULPHUR SUBSTITUTED 3-OXO-2,3-DIHYDROPYRIDAZINE-4-CARBOXAMIDES

The present invention covers sulphur substituted 3-oxo-2,3-dihydropyridazine-4-carboxamide compounds of general formula (I) as described and defined herein, methods of preparing said
5 compounds, intermediate compounds useful for preparing said compounds, pharmaceutical compositions and combinations comprising said compounds, and the use of said compounds for manufacturing pharmaceutical compositions for the treatment or prophylaxis of diseases, in particular cancer or conditions with dysregulated immune responses, as a sole agent or in combination with other active ingredients.

10

BACKGROUND

The AHR (Aryl Hydrocarbon Receptor) is a ligand-activated transcription factor, belonging to the basic helix-loop-helix/Per-Arnt-Sim (bHLH/PAS) family, and is located in the cytosol. Upon ligand binding, the AHR translocates to the nucleus where it heterodimerises with ARNT (AHR
15 Nuclear Translocator) upon which it interacts with DREs (Dioxin Response Elements) of AHR-responsive genes to regulate their transcription. The AHR is best known for binding to environmental toxins and inducing the metabolic machinery, such as cytochrome P 450 enzymes (e.g. CYP1A1, CYP1A2 and CYP1B1), required for their elimination (Reyes et al., Science, 1992, 256(5060):1193-5). Activation of AHR by xenobiotics has demonstrated its role
20 in numerous cellular processes such as embryogenesis, tumorigenesis and inflammation.

AHR is expressed in many cells of the immune system, including dendritic cells (DCs), macrophages, T cells and NK cells, and plays an important role in immunoregulation (Nguyen et al., Front Immunol, 2014, 5:551). The classic exogenous AHR ligands TCDD and 3-methylcholanthrene, for example, are known to induce profound immunosuppression, promote
25 carcinogenesis and induce tumour growth (Gramatzki et al., Oncogene, 2009, 28(28):2593-605; Bui et al., Oncogene, 2009, 28(41):3642-51; Esser et al., Trends Immunol, 2009, 30:447-454). In the context of immunosuppression, AHR activation promotes regulatory T cell generation, inhibits Th1 and Th17 differentiation, directly and indirectly, and decreases the activation and maturation of DCs (Wang et al., Clin Exp Immunol, 2014, 177(2):521-30;
30 Mezrich et al., J Immunol, 2010, 185(6): 3190-8; Wei et al., Lab Invest, 2014, 94(5):528-35; Nguyen et al., PNAS, 2010, 107(46):19961-6). AHR activation modulates the innate immune response and constitutive AHR expression has been shown to negatively regulate the type-I interferon response to viral infection (Yamada et al., Nat Immunol, 2016). Additionally, mice with a constitutively active AHR spontaneously develop tumours (Andersson et al., PNAS,
35 2002, 99(15):9990-5).

In addition to xenobiotics, the AHR can also bind metabolic products of tryptophan degradation. Tryptophan metabolites, such as kynurenine and kynurenic acid, are endogenous

AHR ligands that activate the AHR under physiological conditions (DiNatale et al., Toxicol Sci, 2010, 115(1):89-97; Mezrich et al., J Immunol, 2010, 185(6):3190-8; Opitz et al., Nature, 2011, 478(7368):197-203). Other endogenous ligands are known to bind the AHR although their physiological roles are currently unknown (Nguyen & Bradfield, Chem Res Toxicol, 2008, 21(1):102-116).

The immunosuppressive properties of kynurenine and tryptophan degradation are well described and are implicated in cancer-associated immunosuppression. The enzymes indoleamine-2,3-dioxygenases 1 and 2 (IDO1/IDO2) as well as tryptophan-2,3-dioxygenase 2 (TDO2) are responsible for catalysing the first and rate-limiting step of tryptophan metabolism.

IDO1/2-mediated degradation of tryptophan in tumours and tumour-draining lymph nodes reduces anti-tumour immune responses and inhibition of IDO can suppress tumour formation in animal models (Uyttenhove et al., Nat Med, 2003, 9(10):1269-74 ; Liu et al., Blood, 2005, 115(17): 3520-30; Muller et al., Nat Med, 11(3):312-9; Metz, Cancer Res, 2007, 67(15):7082-7).

TDO2 is also strongly expressed in cancer and can lead to the production of immunosuppressive kynurenine. In glioma, activation of the AHR by kynurenine, downstream of TDO-mediated tryptophan degradation, enhances tumour growth as a consequence of inhibiting anti-tumour immune responses as well as directly promoting tumour cell survival and motility (Opitz et al., Nature, 2011, 478(7368):197-203). AHR ligands generated by tumour cells therefore act in both an autocrine and paracrine fashion on tumour cells and lymphocytes, respectively, to promote tumour growth.

The present invention covers sulphur substituted 3-oxo-2,3-dihydropyridazine-4 carboxamide compounds of general formula (I) which inhibit the AHR.

State of the Art

WO 2010/059401 relates to compounds and compositions for expanding the number of CD34+ cells for transplantation. In particular, WO 2010/059401 relates *inter alia* to heterocyclic compounds capable of down-regulating the activity and/or expression of AHR.

WO 2012/015914 relates to compositions and methods for modulating AHR activity. In particular, WO 2012/015914 relates *inter alia* to heterocyclic compounds that modulate AHR activity for use in therapeutic compositions.

WO 2007/058392 relates to novel heterocyclic compounds and a pharmaceutical use thereof. In particular, WO 2007/058392 relates *inter alia* to heterocyclic compounds having an hepatitis C virus cell infection inhibitory activity.

WO 2002/022587 relates to novel compounds exhibiting inhibitory activities against AMPA receptor and/or kainate receptor. In particular, WO 2002/022587 relates *inter alia* to pyridazinone and triazinone compounds.

US 5,418,233 relates to heterobiaryl derivatives inhibiting cell-cell aggregation and cell-matrix interactions. In particular, US 5,418,233 relates to heterobiaryl derivatives which are histamine receptor antagonists.

WO 2015/143164 relates to antimicrobial agents and screening methods. In particular,
5 WO 2015/143164 relates *inter alia* to pyridazinone compounds as antibiotics.

WO 2009/142732 relates to substituted pyridazinone derivatives and their use as H₃ antagonists/inverse agonists.


However, the state of the art does not describe the sulphur substituted 3-oxo-2,3-dihydropyridazine-4 carboxamide compounds of general formula (I) of the present invention as
10 described and defined herein.

It has now been found, and this constitutes the basis of the present invention, that the compounds of the present invention have surprising and advantageous properties.

In particular, the compounds of the present invention have surprisingly been found to
15 effectively inhibit AHR for which data are given in biological experimental section and may therefore be used for the treatment or prophylaxis of cancer or other conditions where exogenous and endogenous AHR ligands induce dysregulated immune responses, uncontrolled cell growth, proliferation and/or survival of tumour cells, immunosuppression in the context of cancer, inappropriate cellular immune responses, or inappropriate cellular
20 inflammatory responses or diseases which are accompanied with uncontrolled cell growth, proliferation and/or survival of tumour cells, immunosuppression in the context of cancer inappropriate cellular immune responses, or inappropriate cellular inflammatory responses, particularly in which the uncontrolled cell growth, proliferation and/or survival of tumour cells, immunosuppression in the context of cancer, inappropriate cellular immune responses, or
25 inappropriate cellular inflammatory responses is mediated by AHR, such as, for example, liquid and solid tumours, and/or metastases thereof, e.g. head and neck tumours including brain tumours and brain metastases, tumours of the thorax including non-small cell and small cell lung tumours, gastrointestinal tumours including colon, colorectal and pancreatic tumours, liver tumours, endocrine tumours, mammary and other gynecological tumours, urological tumours
30 including renal, bladder and prostate tumours, skin tumours, and sarcomas, and/or metastases thereof.

DESCRIPTION of the INVENTION

In accordance with a first aspect, the present invention covers compounds of general formula (I):



5

(I)

in which

- A represents phenyl or monocyclic heteroaryl, wherein said phenyl is substituted with R⁴, R⁵ and R⁶ and said monocyclic heteroaryl is optionally substituted one to three times, independently from each other, with R^{6a}.
- 10 X represents CR^{4a} or N;
 Y represents CR³ or N;
 Z represents CH or N, wherein
 if X represents N, Y represents CR³ and Z represents CH, and
 if X represents CR^{4a}, Z represents CH and Y represents CR³ or N, and
 15 if Z represents N, Y represents N and X represents CH;
- R¹ represents C₂-C₆-alkyl substituted once with R⁷ and optionally once with hydroxy and optionally one to three times with halogen, or
 C₃-C₆-cycloalkyl substituted once with R⁷ and optionally once with hydroxy, or
 5- to 6-membered heterocycloalkyl optionally substituted once with hydroxy or oxo;
- 20 R² represents chloro, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy or -NR⁸R⁹;
 R³ represents hydrogen, halogen or methyl;
 R⁴ represents hydrogen, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, halogen or cyano;
- 25 R^{4a} represents hydrogen or halogen;
 R⁵ represents hydrogen or halogen;
 R⁶ represents hydrogen or halogen;
 R^{6a} represents C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₆-cycloalkyl, C₁-C₄-alkoxy, halogen or cyano;
- 30 R⁷ represents -SR¹⁰, -SO-R¹⁰, -SO₂-R¹⁰, -SO₂NR⁸R⁹ or -SO(NR⁸)-R¹⁰;
 R⁸ and R⁹ are the same or different and represent, independently from each other, hydrogen, C₁-C₃-alkyl or trifluoroacetyl, or
 together with the nitrogen atom to which they are attached form a 4- to 6-membered

nitrogen containing heterocyclic ring, said ring optionally containing one additional heteroatom selected from O, S, NH, NR^a in which R^a represents a C₁-C₄-alkyl group; R¹⁰ represents C₁-C₄-alkyl;

5 their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

Further, it covers their use in combination with other anti cancer medications such as immunotherapeutics, targeted anti cancer agents or chemotherapy.

10

DEFINITIONS

The term "substituted" means that one or more hydrogen atoms on the designated atom or group are replaced with a selection from the indicated group, provided that the designated atom's normal valency under the existing circumstances is not exceeded. Combinations of substituents and/or variables are permissible.

The term "optionally substituted" means that the number of substituents can be equal to or different from zero. Unless otherwise indicated, it is possible that optionally substituted groups are substituted with as many optional substituents as can be accommodated by replacing a hydrogen atom with a non-hydrogen substituent on any available carbon atom. Commonly, it is possible for the number of optional substituents, when present, to be 1, 2 or 3.

The term "comprising" when used in the specification includes "consisting of".

If within the present text any item is referred to as "as mentioned herein", it means that it may be mentioned anywhere in the present text.

25 The terms as mentioned in the present text have the following meanings:

The term "halogen" means a fluorine, chlorine, bromine or iodine, particularly a fluorine, chlorine or bromine atom.

The term "C₁-C₆-alkyl" means a linear or branched, saturated, monovalent hydrocarbon group having 1, 2, 3, 4, 5 or 6 carbon atoms, e.g. a methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, isobutyl, *tert*-butyl, pentyl, isopentyl, 2-methylbutyl, 1-methylbutyl, 1-ethylpropyl, 1,2-dimethylpropyl, *neo*-pentyl, 1,1-dimethylpropyl, hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1-ethylbutyl, 2-ethylbutyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl, 2,3-dimethylbutyl, 1,2-dimethylbutyl or 1,3-dimethylbutyl group, or an isomer thereof. Particularly, said group has 1, 2, 3 or 4 carbon atoms ("C₁-C₄-alkyl"), e.g. a methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl isobutyl, or *tert*-butyl group, more particularly 1, 2 or 3 carbon atoms ("C₁-C₃-alkyl"), e.g. a methyl, ethyl, *n*-propyl or isopropyl group.

35

The term "C₂-C₈-alkyl" means a linear or branched, saturated, monovalent hydrocarbon group having 2, 3, 4, 5, 6, 7 or 8 carbon atoms, e.g. a ethyl, propyl, isopropyl, butyl, *sec*-butyl, isobutyl, *tert*-butyl, pentyl, isopentyl, 2-methylbutyl, 1-methylbutyl, 1-ethylpropyl, 1,2-dimethylpropyl, *neo*-pentyl, 1,1-dimethylpropyl, hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1-ethylbutyl, 2-ethylbutyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl, 2,3-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 3-ethyl-pentyl or 3-ethyl-hexyl group, or an isomer thereof. Particularly, said group has 2, 3 or 4 carbon atoms ("C₂-C₄-alkyl"), e.g. a ethyl, propyl, isopropyl, butyl, *sec*-butyl isobutyl, or *tert*-butyl group, more particularly 2 or 3 carbon atoms ("C₂-C₃-alkyl"), e.g. a ethyl, *n*-propyl or isopropyl group.

10 The term "C₂-C₈-hydroxyalkyl" means a linear or branched, saturated, monovalent hydrocarbon group in which the term "C₂-C₈-alkyl" is defined *supra*, and in which 1 or 2 hydrogen atoms are replaced with a hydroxy group, e.g. a 1-hydroxyethyl, 2-hydroxyethyl, 1,2-dihydroxyethyl, 3-hydroxypropyl, 2-hydroxypropyl, 1-hydroxypropyl, 1-hydroxypropan-2-yl, 2-hydroxypropan-2-yl, 2,3-dihydroxypropyl, 1,3-dihydroxypropan-2-yl, 3-hydroxy-2-methyl-propyl, 2-hydroxy-2-methyl-propyl, 1-hydroxy-2-methyl-propyl, 3-ethyl-2-hydroxypentyl or 3-ethyl-2-hydroxyhexyl group.

The term "C₁-C₆-haloalkyl" means a linear or branched, saturated, monovalent hydrocarbon group in which the term "C₁-C₆-alkyl" is as defined *supra*, and in which one or more of the hydrogen atoms are replaced, identically or differently, with a halogen atom. Particularly, said halogen atom is a fluorine atom. Said C₁-C₆-haloalkyl group is, for example, fluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, 3,3,3-trifluoropropyl or 1,3-difluoropropan-2-yl. Particularly, said group has 1, 2, 3 or 4 carbon atoms ("C₁-C₄-haloalkyl"), more particularly 1, 2 or 3 carbon atoms ("C₁-C₃-haloalkyl"), e.g. a fluoromethyl, difluoromethyl or trifluoromethyl group.

25 The term "C₁-C₄-alkoxy" means a linear or branched, saturated, monovalent group of formula (C₁-C₄-alkyl)-O-, which means methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-butoxy, *sec*-butoxy, isobutoxy or *tert*-butoxy.

The term "C₃-C₆-cycloalkyl" means a saturated, monovalent, monocyclic hydrocarbon ring which contains 3, 4, 5 or 6 carbon atoms ("C₃-C₆-cycloalkyl"). Said C₃-C₆-cycloalkyl group is a monocyclic hydrocarbon ring, e.g. a cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

30 The term "5- to 6-membered heterocycloalkyl" means a monocyclic, saturated heterocycle with 5 or 6 ring atoms in total, which contains a heteroatom-containing group selected from the group consisting of -SO-, -SO₂-, -SO₂-NR⁸-, -SO(=NR⁸)- and optionally one nitrogen atom, wherein R⁸ means a hydrogen atom or a C₁-C₃-alkyl group. It being possible for said heterocycloalkyl group to be attached to the rest of the molecule via any one of the carbon atoms or, if present, a nitrogen atom.

Said heterocycloalkyl group, without being limited thereto, can be a 5-membered ring, such as tetrahydrothiophene 1-oxide, 1,2-thiazolidine 1-oxide, 1,3-thiazolidine 1-oxide, tetrahydrothiophene 1,1-dioxide, 1,2-thiazolidine 1,1-dioxide, 1,3-thiazolidine 1,1-dioxide, 1,2,5-thiadiazolidine 1,1-dioxide, 1,2,4-thiadiazolidine 1,1-dioxide, 1,2,3-thiadiazolidine 1,1-dioxide, tetrahydro-1H-1 λ^4 -thiophen-1-imine 1-oxide, 1 λ^4 ,2-thiazolidin-1-imine 1-oxide or 1 λ^4 ,3-thiazolidin-1-imine 1-oxide, for example; or a 6 membered ring, such as tetrahydro-2H-thiopyran 1-oxide, 1,2-thiazinane 1-oxide, 1,3-thiazinane 1-oxide, thiomorpholine 1-oxide, tetrahydro-2H-thiopyran 1,1-dioxide, 1,2-thiazinane 1,1-dioxide, 1,3-thiazinane 1,1-dioxide, thiomorpholine 1,1-dioxide, 1,2,6-thiadiazinane 1,1-dioxide, 1,2,5-thiadiazinane 1,1-dioxide, 1,2,4-thiadiazinane 1,1-dioxide, 1,2,3-thiadiazinane 1,1-dioxide, hexahydro-1 λ^4 -thiopyran-1-imine 1-oxide, 1 λ^4 ,2-thiazinan-1-imine 1-oxide, 1 λ^4 ,3-thiazinan-1-imine 1-oxide or 1 λ^4 -thiomorpholin-1-imine 1-oxide, for example.

The term "monocyclic heteroaryl" means a monovalent, aromatic ring having 5 or 6 ring atoms (a "5- or 6-membered heteroaryl" group), which contains at least one ring heteroatom and optionally one or two further ring heteroatoms from the series: N, O and/or S, and which is bound via a ring carbon atom or optionally via a ring nitrogen atom (if allowed by valency).

Said heteroaryl group can be a 5-membered heteroaryl group, such as, for example, thienyl, furanyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl or tetrazolyl; or a 6-membered heteroaryl group, such as, for example, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl or triazinyl.

In general, and unless otherwise mentioned, the heteroaryl or heteroarylene groups include all possible isomeric forms thereof, e.g.: tautomers and positional isomers with respect to the point of linkage to the rest of the molecule. Thus, for some illustrative non-restricting examples, the term pyridinyl includes pyridin-2-yl, pyridin-3-yl and pyridin-4-yl; or the term thienyl includes thien-2-yl and thien-3-yl.

Particularly, the heteroaryl group is a pyrazolyl, isothiazolyl or pyridinyl group.

An oxo substituent in the context of the invention means an oxygen atom, which is bound to a carbon atom via a double bond.

Where the plural form of the word compounds, salts, polymorphs, hydrates, solvates and the like, is used herein, this is taken to mean also a single compound, salt, polymorph, isomer, hydrate, solvate or the like.

By "stable compound" or "stable structure" is meant a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

The compounds of the present invention optionally contain one or more asymmetric centres, depending upon the location and nature of the various substituents desired. It is possible that one or more asymmetric carbon atoms are present in the (R) or (S) configuration, which can

result in racemic mixtures in the case of a single asymmetric centre, and in diastereomeric mixtures in the case of multiple asymmetric centres. In certain instances, it is possible that asymmetry also be present due to restricted rotation about a given bond, for example, the central bond adjoining two substituted aromatic rings of the specified compounds.

5 Preferred compounds are those which produce the more desirable biological activity. Separated, pure or partially purified isomers and stereoisomers or racemic or diastereomeric mixtures of the compounds of the present invention are also included within the scope of the present invention. The purification and the separation of such materials can be accomplished by standard techniques known in the art.

10 Preferred isomers are those which produce the more desirable biological activity. These separated, pure or partially purified isomers or racemic mixtures of the compounds of this invention are also included within the scope of the present invention. The purification and the separation of such materials can be accomplished by standard techniques known in the art.

The optical isomers can be obtained by resolution of the racemic mixtures according to
15 conventional processes, for example, by the formation of diastereoisomeric salts using an optically active acid or base or formation of covalent diastereomers. Examples of appropriate acids are tartaric, diacetyltartaric, ditoluoyltartaric and camphorsulfonic acid. Mixtures of diastereoisomers can be separated into their individual diastereomers on the basis of their physical and/or chemical differences by methods known in the art, for example, by
20 chromatography or fractional crystallisation. The optically active bases or acids are then liberated from the separated diastereomeric salts. A different process for separation of optical isomers involves the use of chiral chromatography (e.g., HPLC columns using a chiral phase), with or without conventional derivatisation, optimally chosen to maximise the separation of the enantiomers. Suitable HPLC columns using a chiral phase are commercially available, such as
25 those manufactured by Daicel, e.g., Chiracel OD and Chiracel OJ, for example, among many others, which are all routinely selectable. Enzymatic separations, with or without derivatisation, are also useful. The optically active compounds of the present invention can likewise be obtained by chiral syntheses utilizing optically active starting materials.

In order to distinguish different types of isomers from each other reference is made to IUPAC
30 Rules Section E (Pure Appl Chem 45, 11-30, 1976).

The present invention includes all possible stereoisomers of the compounds of the present invention as single stereoisomers, or as any mixture of said stereoisomers, e.g. (R)- or (S)- isomers, in any ratio. Isolation of a single stereoisomer, e.g. a single enantiomer or a single diastereomer, of a compound of the present invention is achieved by any suitable state of the
35 art method, such as chromatography, especially chiral chromatography, for example.

Further, the compounds of the present invention can exist as N-oxides, which are defined in that at least one nitrogen of the compounds of the present invention is oxidised. The present invention includes all such possible N-oxides.

The present invention also covers useful forms of the compounds of the present invention, such as metabolites, hydrates, solvates, prodrugs, salts, in particular pharmaceutically acceptable salts, and/or co-precipitates.

The compounds of the present invention can exist as a hydrate, or as a solvate, wherein the compounds of the present invention contain polar solvents, in particular water, methanol or ethanol for example, as structural element of the crystal lattice of the compounds. It is possible for the amount of polar solvents, in particular water, to exist in a stoichiometric or non-stoichiometric ratio. In the case of stoichiometric solvates, e.g. a hydrate, hemi-, (semi-), mono-, sesqui-, di-, tri-, tetra-, penta- etc. solvates or hydrates, respectively, are possible. The present invention includes all such hydrates or solvates.

Further, it is possible for the compounds of the present invention to exist in free form, e.g. as a free base, or as a free acid, or as a zwitterion, or to exist in the form of a salt. Said salt may be any salt, either an organic or inorganic addition salt, particularly any pharmaceutically acceptable organic or inorganic addition salt, which is customarily used in pharmacy, or which is used, for example, for isolating or purifying the compounds of the present invention.

The term "pharmaceutically acceptable salt" refers to an inorganic or organic acid addition salt of a compound of the present invention. For example, see S. M. Berge, *et al.* "Pharmaceutical Salts," J. Pharm. Sci. 1977, 66, 1-19.

A suitable pharmaceutically acceptable salt of the compounds of the present invention may be, for example, an acid-addition salt of a compound of the present invention bearing a nitrogen atom, in a chain or in a ring, for example, which is sufficiently basic, such as an acid-addition salt with an inorganic acid, or "mineral acid", such as hydrochloric, hydrobromic, hydroiodic, sulfuric, sulfamic, bisulfuric, phosphoric, or nitric acid, for example, or with an organic acid, such as formic, acetic, acetoacetic, pyruvic, trifluoroacetic, propionic, butyric, hexanoic, heptanoic, undecanoic, lauric, benzoic, salicylic, 2-(4-hydroxybenzoyl)-benzoic, camphoric, cinnamic, cyclopentanepropionic, digluconic, 3-hydroxy-2-naphthoic, nicotinic, pamoic, pectinic, 3-phenylpropionic, pivalic, 2-hydroxyethanesulfonic, itaconic, trifluoromethanesulfonic, dodecylsulfuric, ethanesulfonic, benzenesulfonic, para-toluenesulfonic, methanesulfonic, 2-naphthalenesulfonic, naphthalenedisulfonic, camphorsulfonic acid, citric, tartaric, stearic, lactic, oxalic, malonic, succinic, malic, adipic, alginic, maleic, fumaric, D-gluconic, mandelic, ascorbic, glucoheptanoic, glycerophosphoric, aspartic, sulfosalicylic, or thiocyanic acid, for example.

Further, another suitably pharmaceutically acceptable salt of a compound of the present invention which is sufficiently acidic, is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium, magnesium or strontium salt, or an aluminium or a zinc salt, or an ammonium salt derived from ammonia or from an organic primary, secondary or tertiary amine having 1 to 20 carbon atoms, such as ethylamine, diethylamine, triethylamine, ethyldiisopropylamine, monoethanolamine, diethanolamine,

triethanolamine, dicyclohexylamine, dimethylaminoethanol, diethylaminoethanol, tris(hydroxymethyl)aminomethane, procaine, dibenzylamine, *N*-methylmorpholine, arginine, lysine, 1,2-ethylenediamine, *N*-methylpiperidine, *N*-methyl-glucamine, *N,N*-dimethyl-glucamine, *N*-ethyl-glucamine, 1,6-hexanediamine, glucosamine, sarcosine, serinol, 2-amino-1,3-
5 propanediol, 3-amino-1,2-propanediol, 4-amino-1,2,3-butanetriol, or a salt with a quarternary ammonium ion having 1 to 20 carbon atoms, such as tetramethylammonium, tetraethylammonium, tetra(*n*-propyl)ammonium, tetra(*n*-butyl)ammonium, *N*-benzyl-*N,N,N*-trimethylammonium, choline or benzalkonium.

Those skilled in the art will further recognise that it is possible for acid addition salts of the
10 claimed compounds to be prepared by reaction of the compounds with the appropriate inorganic or organic acid via any of a number of known methods. Alternatively, alkali and alkaline earth metal salts of acidic compounds of the present invention are prepared by reacting the compounds of the present invention with the appropriate base via a variety of known methods.

15 The present invention includes all possible salts of the compounds of the present invention as single salts, or as any mixture of said salts, in any ratio.

In the present text, in particular in the Experimental Section, for the synthesis of intermediates and of examples of the present invention, when a compound is mentioned as a salt form with the corresponding base or acid, the exact stoichiometric composition of said salt form, as
20 obtained by the respective preparation and/or purification process, is, in most cases, unknown. Unless specified otherwise, suffixes to chemical names or structural formulae relating to salts, such as "hydrochloride", "trifluoroacetate", "sodium salt", or "x HCl", "x CF₃COOH", "x Na⁺", for example, mean a salt form, the stoichiometry of which salt form not being specified.

This applies analogously to cases in which synthesis intermediates or example compounds or
25 salts thereof have been obtained, by the preparation and/or purification processes described, as solvates, such as hydrates, with (if defined) unknown stoichiometric composition.

Furthermore, the present invention includes all possible crystalline forms, or polymorphs, of the compounds of the present invention, either as single polymorph, or as a mixture of more than one polymorph, in any ratio.

30 Moreover, the present invention also includes prodrugs of the compounds according to the invention. The term "prodrugs" here designates compounds which themselves can be biologically active or inactive, but are converted (for example metabolically or hydrolytically) into compounds according to the invention during their residence time in the body.


The invention further includes all possible crystallized and polymorphic forms of the inventive
35 compounds, whereby the polymorphs are existing either as a single polymorph form or are existing as a mixture of several polymorphs in all concentrations.

In accordance with a second embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

- A represents phenyl or monocyclic heteroaryl, wherein said phenyl is substituted with R⁴ and said monocyclic heteroaryl is optionally substituted one time with R^{6a}.
- 5 X represents CR^{4a} or N;
 Y represents CR³ or N;
 Z represents CH or N, wherein
 if X represents N, Y represents CR³ and Z represents CH, and
 if X represents CR^{4a}, Z represents CH and Y represents CR³ or N, and
 10 if Z represents N, Y represents N and X represents CH;
- R¹ represents C₂-C₄-alkyl substituted once with R⁷ and optionally once with hydroxy, or C₃-C₆-cycloalkyl substituted once with R⁷ and optionally once with hydroxy, or 5- to 6-membered heterocycloalkyl optionally substituted once with hydroxy;
- R² represents chloro, methyl, fluoromethyl, difluoromethyl or trifluoromethyl;
- 15 R³ represents hydrogen;
 R⁴ represents methyl, halogen or cyano;
 R^{4a} represents hydrogen;
 R⁵ represents hydrogen;
 R⁶ represents hydrogen;
- 20 R^{6a} represents C₁-C₄-alkyl, C₁-C₄-haloalkyl, halogen or cyano;
 R⁷ represents -SR¹⁰, -SO-R¹⁰, -SO₂-R¹⁰, -SO₂NR⁸R⁹ or -SO(NR⁸)-R¹⁰;
 R⁸ and R⁹ are the same or different and represent, independently from each other, hydrogen, C₁-C₃-alkyl or trifluoroacetyl;
 R¹⁰ represents C₁-C₃-alkyl;
- 25 their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.


In accordance with a third embodiment of the first aspect, the present invention covers
 30 compounds of general formula (I), *supra*, in which:

A represents a group selected from:



wherein * indicates the point of attachment of said group with the rest of the molecule;

- X represents CH;
 Y represents CH or N;
 Z represents CH;
 R¹ represents ethyl or propyl substituted once with R⁷, or
 5 a group selected from:



wherein * indicates the point of attachment of said group with the rest of the molecule;

- R² represents chloro or trifluoromethyl;
 R⁷ represents -SR¹⁰, -SO-R¹⁰, -SO₂-R¹⁰ or -SO(NR⁸)-R¹⁰;
 10 R⁸ represents hydrogen or trifluoroacetyl;
 R¹⁰ represents methyl;

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

15

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

- A represents phenyl or monocyclic heteroaryl, wherein said phenyl is substituted with R⁴, R⁵ and R⁶ and said monocyclic heteroaryl is optionally substituted one to three times,
 20 independently from each other, with R^{6a}.

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.


- 25 In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

- A represents phenyl or monocyclic heteroaryl, wherein said phenyl is substituted with R⁴ and said monocyclic heteroaryl is optionally substituted one time with R^{6a}.

- 30 their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

A represents a group selected from:



wherein * indicates the point of attachment of said group with the rest of the molecule;

5 their polymorphs, enantiomers, diastereomers, racemates, tautomers, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

10 X represents CR^{4a} or N;

Y represents CR^3 or N;

Z represents CH or N, wherein

if X represents N, Y represents CR^3 and Z represents CH, and

if X represents CR^{4a} , Z represents CH and Y represents CR^3 or N, and

15 if Z represents N, Y represents N and X represents CH;

their polymorphs, enantiomers, diastereomers, racemates, tautomers, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

20 In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

X represents CH;

Y represents CH or N;

Z represents CH;

25 their polymorphs, enantiomers, diastereomers, racemates, tautomers, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers 30 compounds of general formula (I), *supra*, in which:

R^1 represents C_2 - C_6 -alkyl substituted once with R^7 and optionally once with hydroxy and optionally one to three times with halogen, or

C₃-C₆-cycloalkyl substituted once with R⁷ and optionally once with hydroxy, or
 5- to 6-membered heterocycloalkyl optionally substituted once with hydroxy or oxo;
 their polymorphs, enantiomers, diastereomers, racemates, tautomers, N-oxides, hydrates
 and solvates, as well as their physiological acceptable salts and solvates of these salts, as well
 5 as mixtures of the same.


In accordance with a further embodiment of the first aspect, the present invention covers
 compounds of general formula (I), *supra*, in which:

R¹ represents C₂-C₄-alkyl substituted once with R⁷ and optionally once with hydroxy, or
 10 C₃-C₆-cycloalkyl substituted once with R⁷ and optionally once with hydroxy, or
 5- to 6-membered heterocycloalkyl optionally substituted once with hydroxy;
 their polymorphs, enantiomers, diastereomers, racemates, tautomers, N-oxides, hydrates
 and solvates, as well as their physiological acceptable salts and solvates of these salts, as well
 as mixtures of the same.

15

In accordance with a further embodiment of the first aspect, the present invention covers
 compounds of general formula (I), *supra*, in which:

R¹ represents ethyl or propyl substituted once with R⁷, or
 a group selected from:



wherein * indicates the point of attachment of said group with the rest of the molecule;

their polymorphs, enantiomers, diastereomers, racemates, tautomers, N-oxides, hydrates
 and solvates, as well as their physiological acceptable salts and solvates of these salts, as well
 as mixtures of the same.

25

In accordance with a further embodiment of the first aspect, the present invention covers
 compounds of general formula (I), *supra*, in which:

R² represents chloro, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, methoxy,
 difluoromethoxy, trifluoromethoxy or -NR⁸R⁹;

30 their polymorphs, enantiomers, diastereomers, racemates, tautomers, N-oxides, hydrates
 and solvates, as well as their physiological acceptable salts and solvates of these salts, as well
 as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R² represents chloro, methyl, fluoromethyl, difluoromethyl or trifluoromethyl;

5 their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

10 R² represents chloro or trifluoromethyl;

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

15 In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R³ represents hydrogen, halogen or methyl;

20 their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R³ represents hydrogen;

25 their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

30 R⁴ represents hydrogen, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, halogen or cyano;

35 their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R⁴ represents methyl, halogen or cyano;
their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

5

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R⁴ represents halogen;
their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

10

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R⁴ represents fluoro;
their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

15

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R⁵ represents hydrogen or halogen;
their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

20

25

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R⁵ represents hydrogen;
their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

30

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R⁶ represents hydrogen or halogen;

35

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

5 In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R⁶ represents hydrogen;

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well

10 as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R^{6a} represents C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₆-cycloalkyl, C₁-C₄-alkoxy, halogen or

15 cyano;

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

20 In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R^{6a} represents C₁-C₄-alkyl, C₁-C₄-haloalkyl, halogen or cyano;

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well

25 as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R^{6a} represents methyl;

30 their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

35 R⁷ represents -SR¹⁰, -SO-R¹⁰, -SO₂-R¹⁰, -SO₂NR⁸R⁹ or -SO(NR⁸)-R¹⁰;

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

5 In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R⁷ represents -SR¹⁰, -SO-R¹⁰, -SO₂-R¹⁰ or -SO(NR⁸)-R¹⁰;

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well

10 as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R⁸ and R⁹ are the same or different and represent, independently from each other, hydrogen,

15 C₁-C₃-alkyl or trifluoroacetyl, or

together with the nitrogen atom to which they are attached form a 4- to 6-membered nitrogen containing heterocyclic ring, said ring optionally containing one additional

heteroatom selected from O, S, NH, NR^a in which R^a represents a C₁-C₄-alkyl group;

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates

20 and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

25 R⁸ and R⁹ are the same or different and represent, independently from each other, hydrogen,

C₁-C₃-alkyl or trifluoroacetyl;

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates

and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

30

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R⁸ represents hydrogen or trifluoroacetyl;

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates

35 and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R¹⁰ represents C₁-C₄-alkyl;

5 their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

10 R¹⁰ represents C₁-C₃-alkyl;

their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

15 In accordance with a further embodiment of the first aspect, the present invention covers compounds of general formula (I), *supra*, in which:

R¹⁰ represents methyl;

20 their polymorphs, enantiomeres, diastereomeres, racemates, tautomeres, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

In a particular further embodiment of the first aspect, the present invention covers combinations of two or more of the above mentioned embodiments under the heading "further embodiments of the first aspect of the present invention".

25 The present invention covers any sub-combination within any embodiment or aspect of the present invention of compounds of general formula (I), *supra*.


The present invention covers any sub-combination within any embodiment or aspect of the present invention of intermediate compounds of general formula (VII). The present invention covers the compounds of general formula (I) which are disclosed in the Example Section of
30 this text, *infra*.

The compounds according to the invention of general formula (I) can be prepared according to the following scheme 1. The scheme and procedures described below illustrate synthetic routes to the compounds of general formula (I) of the invention and are not intended to be
35 limiting. It is clear to the person skilled in the art that the order of transformations as exemplified in scheme 1 can be modified in various ways. The order of transformations exemplified in this scheme is therefore not intended to be limiting. In addition, interconversion of any of the substituents R¹, R², R³, R⁴, R⁵, R⁶ or R⁷ can be achieved before and/or after the

exemplified transformations. These modifications can be such as the introduction of protecting groups, cleavage of protecting groups, reduction or oxidation of functional groups, halogenation, metallation, metal-catalysed coupling reactions, substitution or other reactions known to the person skilled in the art. These transformations include those which introduce a
5 functionality which allows for further interconversion of substituents. Appropriate protecting groups and their introduction and cleavage are well-known to the person skilled in the art. Specific examples are described in the subsequent paragraphs.

Scheme 1 shows a route for the preparation of compounds of general formula (I) in which R¹,
10 R², A, X, Y and Z have the meaning as given for general formula (I), *supra*. Ketomalonates according to formula (III) are in some instances commercially available or can be synthesized from alpha-halo-acetophenones (II) according to procedures known to the person skilled in the art. Related alpha-halo-acetophenones are usually commercially available. Conversion of such
15 alpha-halo-acetophenones with malonic acid esters according to formula (IIa) in the presence of a suitable base in a suitable solvent results in the formation of ketomalonates according to formula (III). R in formula (IIa), (III), (V) and (VI) represents a suitable alkyl group such as methyl, ethyl, propyl or other homologous groups or isomers thereof. A suitable solvent can be, but should not be restricted to, acetone, acetonitrile, DMF, DMA, DMSO or THF, or mixtures of these or other solvents. A suitable base can be, but should not be restricted to, potassium
20 carbonate, sodium hydride, cesium carbonate or potassium hexamethyldisilazide.

Formation of dihydropyridazinones according to formula (V) from intermediates (III) and suitable aryl-hydrazines (IV), which are in many cases commercially available, can be
accomplished by reaction of these components in a suitable solvent at elevated temperature. A
25 suitable solvent can be, but should not be restricted to, ethanol or acetic acid.



Scheme 1: Route for the preparation of compounds of general formula (I) in which R¹, R², A, X, Y and Z have the meaning as given for general formula (I), *supra*, and Hal represents halogen and R represents C₁-C₄-alkyl.

5

Dihydropyridazinones according to formula (V) can be transferred to pyridazinones according to formula (VI). This can be accomplished by the use of a suitable reagents such as copper dichloride at elevated temperature (Bioorg. Med. Chem. Lett., **21**, (2011), P. 6362 ff.; Synthesis, (2003), p. 436 ff.; J. Med. Chem., **46**, (2003), p. 349 ff.).


10

The resulting pyridazinones according to formula (VI) with an ester functional group can be converted by methods known to the person skilled in the art, for example by basic hydrolysis with, for example, aqueous alkali metal hydroxides, or by acidic hydrolysis using, for example, hydrogen chloride in dioxane or trifluoroacetic acid, into the pyridazinone carboxylic acids (VII).

15

These can be converted by coupling with amines of the formula (VIII) in which R¹ is as defined for the general formula (I), *supra*. Coupling agents and methods for such syntheses of carboxamides from carboxylic acids and amines are known to the person skilled in the art.

Examples which may be mentioned here include the use of HATU, HBTU, PyBOB or T3P with the addition of a suitable base. The conversion of the carboxylic acids to their amides is described in general terms in reference books such as "Compendium of Organic Synthetic Methods", volume I-VI (Wiley Interscience) or "The Practice of Peptide Synthesis", Bodansky (Springer Verlag).



Scheme 2: Additional route for the preparation of intermediate (VI) in which R², A, X, Y and Z have the meaning as given for general formula (I), *supra*, R represents C₁-C₄-alkyl, R' and R'' represent simultaneously H or C₁-C₄-alkyl or form together a C₂-C₇-alkylene group as part of a 1,2- or 1,3-diol boronic ester or a -CO-CH₂-(NCH₃)-CH₂-CO- group.

Ketomalonates according to formula (III) are in some instances commercially available or can be synthesized as described above. R in formula (III), (IX), (X) and (VI) represents a suitable alkyl group such as methyl, ethyl, propyl or other homologous groups or isomers thereof.


Formation of dihydropyridazinones according to formula (IX) from intermediates (III) and hydrazine can be accomplished by reaction in a suitable solvent at elevated temperature. A suitable solvent can be, but should not be restricted to, ethanol or acetic acid. R in formula (III), (IX), (X) and (VI) represents a suitable alkyl group such as C₁-C₄ alkyl.

Dihydropyridazinones (IX) can be transferred to pyridazinones (X). This can be accomplished by the use of a suitable reagent. A suitable reagent can be, but should not be restricted to, copper dichloride at elevated temperature.

Substituted pyridazinones according to formula (VI) can be prepared by Chan-Lam coupling reactions of pyridazinones according to formula (X) using boron derivatives as boronic acids, boronic acid esters, and mida boronates with suitable solvents at room temperature or

elevated temperatures. A suitable solvent can be, but should not be restricted to, acetonitrile, dichloromethane, pyridine or DMF. A suitable catalyst can be, but should not be restricted to copper (II) acetate. Suitable basic additives can be, but should not be restricted to, trialkylamine, 2,2'-bipyridine, sodium carbonate, cesium carbonate, and cesium hydrogen carbonate.


5



Scheme 3: Additional route for the preparation of intermediate (X) in which R², X, Y and Z have the meaning as given for general formula (I), *supra*, and R represents C₁-C₄-alkyl.

10 Acetophenones according to formula (XII) are in many cases commercially available. R in formula (XIII), (XIV), and (X) represents a suitable alkyl group such as methyl, ethyl, propyl or other homologous groups or isomers thereof. The formation of ketohydroxymalonates according to formula (XIV) can be accomplished by reaction at elevated temperatures.


15 Formation of pyridazinones according to formula (X) from intermediates (XIV) and hydrazine can be accomplished by reaction in a suitable solvent at elevated temperature. A suitable solvent can be, but should not be restricted to, ethanol or acetic acid.



Scheme 4: Additional route for the preparation of intermediate (VI) in which R², A, X, Y and Z have the meaning as given for general formula (I), *supra*, and R represents C₁-C₄-alkyl.

- 5 Acetophenones according to formula (XII) are in many cases commercially available. R in formula (XIII), (XIV), and (VI) represents a suitable alkyl group such as methyl, ethyl, propyl or other homologous groups or isomers thereof. The formation of ketohydroxymalonates according to formula (XIV) can be accomplished by reaction at elevated temperatures.

10 Formation of pyridazinones according to formula (VI) from intermediates (XIV) and hydrazine (IV) can be accomplished by reaction in a suitable solvent at elevated temperature. A suitable solvent can be, but should not be restricted to, ethanol or acetic acid.



Scheme 5: Route for the preparation of compounds of general formulae (la), (lb), (lc) and (ld) in which R², A, X, Y and Z have the meaning as given for general formula (I), *supra* and X¹-S-X², X¹-SO-X², X¹-SO₂-X² and X¹-SO(NR⁸)-X² have the meaning as given for R¹ in the general formula (I), *supra*.

The acids according to formula (VII) are converted by coupling with amines according to formulae (VIIIa), (VIIIb) and, (VIIIc) in which R¹ is as defined for the general formula (I), *supra*.

10 Coupling agents and methods for such syntheses of carboxamides from carboxylic acids and amines are known to the person skilled in the art. Examples which may be mentioned here include the use of HATU, HBTU, PyBOB or T3P with the addition of a suitable base. The conversion of carboxylic acids to their amides is described in general terms in reference books. Amines for the synthesis of sulfides according to formula (la) are commercially available and can be coupled as described, *supra*. Amines for the synthesis of sulfones according to formula (lb) are in some cases commercially available and can be coupled as described, *supra*. Sulfones according to formula (lb) can be also prepared starting from sulfides of formula (la) by methods known to the person skilled in the art, for example, by oxidation of the sulfides with peracids, for example, meta-chloroperoxybenzoic acid, in a suitable solvent, for example, chloroform.

15
20

The sulfides according to formula (Ia) and the sulfoxides according to formula (Ic) can be oxidized to the sulfones according to formula (Ib) by methods known to the person skilled in the art, for example, by oxidation of the sulfides or sulfoxides with peracids, for example, meta-chloroperbenzoic acid, in a suitable solvent, for example, chloroform or dichloromethane at 5 0°C to room temperature, especially the peracid is added between 0-10°C and stirred for 1-2 at 0-10 °C and then stirring is continued at room temperature.


Sulfides according to formula (Ia) can be converted to the sulfoxides according to formula (Ic) by methods known to the person skilled in the art, for example, with sodium metaperiodate in a 10 suitable solvent or a mixture of solvents, for example, a mixture of water, methanol or ethanol, and an organic solvent, e.g. acetone, dioxane, or tetrahydrofuran, at 0-50°C, in particular at room temperature.

The sulfoxides can be converted to the sulfoximines according to formula (Id) by methods 15 known to the person skilled in the art. The synthesis starts with converting the sulfoxide (Ic) to the trifluoroacetates (Ie). To the compound in a suitable solvent, for example, dichloromethane, are added 2,2,2-trifluoroacetamide, (diacetoxyiodo)benzene, rhodium(II)acetate dimer, and magnesium oxide and stirred at room temperature. The reaction mixture can be diluted with dichloromethane, washed with water, dried, e.g. over magnesium sulfate, and concentrated to 20 dryness. The intermediate (Ie) can be purified, e.g. by HPLC or flash chromatography. The trifluoroacetamide (Ie) is then hydrolysed under basic conditions to the sulfoximines (Id). The trifluoroacetate (Ie) is dissolved in an alcohol, e.g. in methanol, and hydrolysed under basic conditions with an alkali metal hydroxide or alkali metal carbonate, in particular with alkali metal carbonate, e.g. potassium carbonate, under room temperature. This two step procedure 25 can be performed without isolating and purifying the trifluoroacetate (Ie). The crude trifluoroacetate (Ie) is subjected to the hydrolysis, described, *supra*, leading to the sulfoximes (Id).

Alternatively, the sulfides according to formula (Ia) can be converted to the sulfoximines according to formula (Id). To sodium tert-butoxide in a suitable solvent, e.g. tetrahydrofuran, 30 at reduced temperature, e.g. 10 °C, is added 2,2,2-trifluoroacetamide in a suitable solvent, e.g. tetrahydrofuran, and 1,3-dibromo-5,5-dimethylhydantoin in tetrahydrofuran. The sulfide (Ia) dissolved in suitable solvent, e.g. tetrahydrofuran and dioxane, is added under reduced temperature, e.g. 10 °C, and stirred at this temperature and later at room temperature. In the next step, the crude material is dissolved in suitable solvent, e.g. methanol, oxone is added 35 and the pH is maintained at 10-11 with an aqueous solution of an alkali metal hydroxide, e.g. potassium hydroxide. The reaction mixture is stirred at room temperature.

The compounds are either commercially available or can be prepared according to procedures available from the public domain, as understandable to the person skilled in the art. Specific examples are described in the Experimental Section.

- 5 In accordance with a second aspect, the present invention covers methods of preparing compounds of general formula (I) as defined *supra*, said methods comprising the step of allowing an intermediate compound of general formula (VII):



(VII),


10 in which

- A represents phenyl or monocyclic heteroaryl, wherein said phenyl is substituted with R⁴, R⁵ and R⁶ and said monocyclic heteroaryl is optionally substituted one to three times, independently from each other, with R^{6a}.
- X represents CR^{4a} or N;
- 15 Y represents CR³ or N;
- Z represents CH or N, wherein
 if X represents N, Y represents CR³ and Z represents CH, and
 if X represents CR^{4a}, Z represents CH and Y represents CR³ or N, and
 if Z represents N, Y represents N and X represents CH;
- 20 R² represents chloro, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy or -NR⁸R⁹;
- R³ represents hydrogen, halogen or methyl;
- R⁴ represents hydrogen, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, halogen or cyano;
- 25 R^{4a} represents hydrogen or halogen;
- R⁵ represents hydrogen or halogen;
- R⁶ represents hydrogen or halogen;
- R^{6a} represents C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₆-cycloalkyl, C₁-C₄-alkoxy, halogen or cyano;
- 30 R⁸ and R⁹ are the same or different and represent, independently from each other, hydrogen, C₁-C₃-alkyl or trifluoroacetyl, or together with the nitrogen atom to which they are attached form a 4- to 6-membered

nitrogen containing heterocyclic ring, said ring optionally containing one additional heteroatom selected from O, S, NH, NR^a in which R^a represents a C₁-C₄-alkyl group;

to react with a compound of general formula (VIII) :

5



in which

R¹ represents C₂-C₆-alkyl substituted once with R⁷ and optionally once with hydroxy and optionally one to three times with halogen, or

10

C₃-C₆-cycloalkyl substituted once with R⁷ and optionally once with hydroxy, or

5- to 6-membered heterocycloalkyl optionally substituted once with hydroxy or oxo;

R⁷ represents -SR¹⁰, -SO-R¹⁰, -SO₂-R¹⁰, -SO₂NR⁸R⁹ or -SO(NR⁸)-R¹⁰;

R⁸ and R⁹ are the same or different and represent, independently from each other, hydrogen,

15


C₁-C₃-alkyl or trifluoroacetyl, or

together with the nitrogen atom to which they are attached form a 4- to 6-membered nitrogen containing heterocyclic ring, said ring optionally containing one additional heteroatom selected from O, S, NH, NR^a in which R^a represents a C₁-C₄-alkyl group;

R¹⁰ represents C₁-C₄-alkyl;

20

thereby giving a compound of general formula (I) :




(I),

in which R¹, R², A, X, Y and Z are as defined *supra*.

25 The present invention covers methods of preparing compounds of the present invention of general formula (I), said methods comprising the steps as described in the Experimental Section herein.

In accordance with a third aspect, the present invention covers intermediate compounds which are useful for the preparation of the compounds of general formula (I), *supra*.

30 Particularly, the inventions covers the intermediate compounds of general formula (VII) :




(VII),

in which

- 5 A represents phenyl or monocyclic heteroaryl, wherein said phenyl is substituted with R⁴, R⁵ and R⁶ and said monocyclic heteroaryl is optionally substituted one to three times, independently from each other, with R^{6a}.
- X represents CR^{4a} or N;
- Y represents CR³ or N;
- Z represents CH or N, wherein
- 10 if X represents N, Y represents CR³ and Z represents CH, and
 if X represents CR^{4a}, Z represents CH and Y represents CR³ or N, and
 if Z represents N, Y represents N and X represents CH;
- R² represents chloro, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy or -NR⁸R⁹;
- 15 R³ represents hydrogen, halogen or methyl;
- R⁴ represents hydrogen, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, halogen or cyano;
- R^{4a} represents hydrogen or halogen;
- R⁵ represents hydrogen or halogen;
- 20 R⁶ represents hydrogen or halogen;
- R^{6a} represents C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₆-cycloalkyl, C₁-C₄-alkoxy, halogen or cyano;
- R⁸ and R⁹ are the same or different and represent, independently from each other, hydrogen, C₁-C₃-alkyl or trifluoroacetyl, or
- 25 together with the nitrogen atom to which they are attached form a 4- to 6-membered nitrogen containing heterocyclic ring, said ring optionally containing one additional heteroatom selected from O, S, NH, NR^a in which R^a represents a C₁-C₄-alkyl group.

30 In accordance with a forth aspect, the present invention covers the use of said intermediate compounds for the preparation of a compound of general formula (I) as defined *supra*.

Particularly, the inventions covers the use of intermediate compounds of general formula (VII) :



(VII),

in which

- A represents phenyl or monocyclic heteroaryl, wherein said phenyl is substituted with R⁴,
 5 R⁵ and R⁶ and said monocyclic heteroaryl is optionally substituted one to three times, independently from each other, with R^{6a}.
- X represents CR^{4a} or N;
 Y represents CR³ or N;
 Z represents CH or N, wherein
 10 if X represents N, Y represents CR³ and Z represents CH, and
 if X represents CR^{4a}, Z represents CH and Y represents CR³ or N, and
 if Z represents N, Y represents N and X represents CH;
- R² represents chloro, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy or -NR⁸R⁹;
- 15 R³ represents hydrogen, halogen or methyl;
 R⁴ represents hydrogen, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, halogen or cyano;
- R^{4a} represents hydrogen or halogen;
 R⁵ represents hydrogen or halogen;
- 20 R⁶ represents hydrogen or halogen;
 R^{6a} represents C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₆-cycloalkyl, C₁-C₄-alkoxy, halogen or cyano;
- R⁸ and R⁹ are the same or different and represent, independently from each other, hydrogen, C₁-C₃-alkyl or trifluoroacetyl, or
 25 together with the nitrogen atom to which they are attached form a 4- to 6-membered nitrogen containing heterocyclic ring, said ring optionally containing one additional heteroatom selected from O, S, NH, NR^a in which R^a represents a C₁-C₄-alkyl group;
- for the preparation of a compound of general formula (I) as defined *supra*.
- 30 The present invention covers the intermediate compounds which are disclosed in the Example Section of this text, *infra*.
- The present invention covers any sub-combination within any embodiment or aspect of the present invention of intermediate compounds of general formula (VII), *supra*.

The compounds of general formula (I) of the present invention can be converted to any salt, preferably pharmaceutically acceptable salts, as described herein, by any method which is known to the person skilled in the art. Similarly, any salt of a compound of general formula (I) of the present invention can be converted into the free compound, by any method which is known to the person skilled in the art.

Compounds of general formula (I) of the present invention demonstrate a valuable pharmacological spectrum of action, which could not have been predicted. Compounds of the present invention have surprisingly been found to effectively inhibit AHR and it is possible therefore that said compounds be used for the treatment or prophylaxis of diseases, preferably cancer or conditions with dysregulated immune responses or other disorders associated with aberrant AHR signaling, in humans and animals.

Disorders and conditions particularly suitable for treatment with an AHR inhibitor of the present invention are liquid and solid tumours, such as cancers of the breast, respiratory tract, brain, reproductive organs, digestive tract, urinary tract, eye, liver, skin, head and neck, thyroid, parathyroid and their distant metastases. Those disorders also include lymphomas, sarcomas, and leukaemias.

Examples of breast cancers include, but are not limited to, triple negative breast cancer, invasive ductal carcinoma, invasive lobular carcinoma, ductal carcinoma *in situ*, and lobular carcinoma *in situ*.

Examples of cancers of the respiratory tract include, but are not limited to, small-cell and non-small-cell lung carcinoma, as well as bronchial adenoma and pleuropulmonary blastoma.

Examples of brain cancers include, but are not limited to, brain stem and hypophtalmic glioma, cerebellar and cerebral astrocytoma, glioblastoma, medulloblastoma, ependymoma, as well as neuroectodermal and pineal tumour.

Tumours of the male reproductive organs include, but are not limited to, prostate and testicular cancer.

Tumours of the female reproductive organs include, but are not limited to, endometrial, cervical, ovarian, vaginal, and vulvar cancer, as well as sarcoma of the uterus.

Examples of ovarian cancer include, but are not limited to serous tumour, endometrioid tumour, mucinous cystadenocarcinoma, granulosa cell tumour, Sertoli-Leydig cell tumour and arrhenoblastoma.

Examples of cervical cancer include, but are not limited to squamous cell carcinoma, adenocarcinoma, adenosquamous carcinoma, small cell carcinoma, neuroendocrine tumour, glassy cell carcinoma and villoglandular adenocarcinoma.

Tumours of the digestive tract include, but are not limited to, anal, colon, colorectal, esophageal, gallbladder, gastric, pancreatic, rectal, small-intestine, and salivary gland cancers.

Examples of esophageal cancer include, but are not limited to esophageal cell carcinomas and adenocarcinomas, as well as squamous cell carcinomas, leiomyosarcoma, malignant melanoma, rhabdomyosarcoma and lymphoma,.

5 Examples of gastric cancer include, but are not limited to intestinal type and diffuse type gastric adenocarcinoma.

Examples of pancreatic cancer include, but are not limited to ductal adenocarcinoma, adenosquamous carcinomas and pancreatic endocrine tumours.

Tumours of the urinary tract include, but are not limited to, bladder, penile, kidney, renal pelvis, ureter, urethral and human papillary renal cancers.

10 Examples of kidney cancer include, but are not limited to renal cell carcinoma, urothelial cell carcinoma, juxtaglomerular cell tumour (reninoma), angiomyolipoma, renal oncocytoma, Bellini duct carcinoma, clear-cell sarcoma of the kidney, mesoblastic nephroma and Wilms' tumour.

Examples of bladder cancer include, but are not limited to transitional cell carcinoma, squamous cell carcinoma, adenocarcinoma, sarcoma and small cell carcinoma.

15 Eye cancers include, but are not limited to, intraocular melanoma and retinoblastoma.

Examples of liver cancers include, but are not limited to, hepatocellular carcinoma (liver cell carcinomas with or without fibrolamellar variant), cholangiocarcinoma (intrahepatic bile duct carcinoma), and mixed hepatocellular cholangiocarcinoma.

20 Skin cancers include, but are not limited to, squamous cell carcinoma, Kaposi's sarcoma, malignant melanoma, Merkel cell skin cancer, and non-melanoma skin cancer.

Head-and-neck cancers include, but are not limited to, squamous cell cancer of the head and neck, laryngeal, hypopharyngeal, nasopharyngeal, oropharyngeal cancer, salivary gland cancer, lip and oral cavity cancer and squamous cell.

25 Lymphomas include, but are not limited to, AIDS-related lymphoma, non-Hodgkin's lymphoma, cutaneous T-cell lymphoma, Burkitt lymphoma, Hodgkin's disease, and lymphoma of the central nervous system.

Sarcomas include, but are not limited to, sarcoma of the soft tissue, osteosarcoma, malignant fibrous histiocytoma, lymphosarcoma, and rhabdomyosarcoma.

30 Leukemias include, but are not limited to, acute myeloid leukemia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, chronic myelogenous leukemia, and hairy cell leukemia.

The term "treating" or "treatment" as stated throughout this document is used conventionally, for example the management or care of a subject for the purpose of combating, alleviating, reducing, relieving, improving the condition of a disease or disorder, such as a carcinoma.

35 The compounds of the present invention can be used in particular in therapy and prevention, *i.e.* prophylaxis, of tumour growth and metastases, especially in solid tumours of all indications and stages with or without pre-treatment of the tumour growth.

Generally, the use of chemotherapeutic agents and/or anti-cancer agents in combination with a compound or pharmaceutical composition of the present invention will serve to:

yield better efficacy in reducing the growth of a tumour or even eliminate the tumour as compared to administration of either agent alone,

- 5 provide for the administration of lesser amounts of the administered chemotherapeutic agents, provide for a chemotherapeutic treatment that is well tolerated in the patient with fewer deleterious pharmacological complications than observed with single agent chemotherapies and certain other combined therapies,
- 10 provide for treating a broader spectrum of different cancer types in mammals, especially humans, provide for a higher response rate among treated patients, provide for a longer survival time among treated patients compared to standard chemotherapy treatments, provide a longer time for tumour progression, and/or
- 15 yield efficacy and tolerability results at least as good as those of the agents used alone, compared to known instances where other cancer agent combinations produce antagonistic effects.

In addition, the compounds of general formula (I) of the present invention can also be used in combination with radiotherapy and/or surgical intervention.

- 20 In a further embodiment of the present invention, the compounds of general formula (I) of the present invention may be used to sensitize a cell to radiation, *i.e.* treatment of a cell with a compound of the present invention prior to radiation treatment of the cell renders the cell more susceptible to DNA damage and cell death than the cell would be in the absence of any treatment with a compound of the present invention. In one aspect, the cell is treated with at
- 25 least one compound of general formula (I) of the present invention.

Thus, the present invention also provides a method of killing a cell, wherein a cell is administered one or more compounds of the present invention in combination with conventional radiation therapy.

- 30 The present invention also provides a method of rendering a cell more susceptible to cell death, wherein the cell is treated with one or more compounds of general formula (I) of the present invention prior to the treatment of the cell to cause or induce cell death. In one aspect, after the cell is treated with one or more compounds of general formula (I) of the present invention, the cell is treated with at least one compound, or at least one method, or a combination thereof, in order to cause DNA damage for the purpose of inhibiting the function of
- 35 the normal cell or killing the cell.

In other embodiments of the present invention, a cell is killed by treating the cell with at least one DNA damaging agent, *i.e.* after treating a cell with one or more compounds of general formula (I) of the present invention to sensitize the cell to cell death, the cell is treated with at

least one DNA damaging agent to kill the cell. DNA damaging agents useful in the present invention include, but are not limited to, chemotherapeutic agents (e.g. cis platin), ionizing radiation (X-rays, ultraviolet radiation), carcinogenic agents, and mutagenic agents.

In other embodiments, a cell is killed by treating the cell with at least one method to cause or induce DNA damage. Such methods include, but are not limited to, activation of a cell signalling pathway that results in DNA damage when the pathway is activated, inhibiting of a cell signalling pathway that results in DNA damage when the pathway is inhibited, and inducing a biochemical change in a cell, wherein the change results in DNA damage. By way of a non-limiting example, a DNA repair pathway in a cell can be inhibited, thereby preventing the repair of DNA damage and resulting in an abnormal accumulation of DNA damage in a cell.

In one aspect of the invention, a compound of general formula (I) of the present invention is administered to a cell prior to the radiation or other induction of DNA damage in the cell. In another aspect of the invention, a compound of general formula (I) of the present invention is administered to a cell concomitantly with the radiation or other induction of DNA damage in the cell. In yet another aspect of the invention, a compound of general formula (I) of the present invention is administered to a cell immediately after radiation or other induction of DNA damage in the cell has begun.

In another aspect, the cell is in vitro. In another embodiment, the cell is in vivo.

The compounds of the present invention can be administered as the sole pharmaceutical agent or in combination with one or more other pharmaceutically active ingredients where the combination causes no unacceptable adverse effects. The present invention also covers such pharmaceutical combinations. For example, the compounds of the present invention can be combined with: 131I-chTNT, abarelix, abiraterone, aclarubicin, adalimumab, ado-trastuzumab emtansine, afatinib, aflibercept, aldesleukin, alectinib, alemtuzumab, alendronic acid, alitretinoin, altretamine, amifostine, aminoglutethimide, hexyl aminolevulinate, amrubicin, amsacrine, anastrozole, anecstim, anethole dithiolethione, anetumab ravtansine, angiotensin II, antithrombin III, aprepitant, arcitumomab, arglabin, arsenic trioxide, asparaginase, atezolizumab, axitinib, azacitidine, basiliximab, belotecan, bendamustine, besilesomab, belinostat, bevacizumab, bexarotene, bicalutamide, bisantrene, bleomycin, blinatumomab, bortezomib, buserelin, bosutinib, brentuximab vedotin, busulfan, cabazitaxel, cabozantinib, calcitonine, calcium folinate, calcium levofolate, capecitabine, capromab, carbamazepine carboplatin, carboquone, carfilzomib, carmofur, carmustine, catumaxomab, celecoxib, celmoleukin, ceritinib, cetuximab, chlorambucil, chlormadinone, chlormethine, cidofovir, cinacalcet, cisplatin, cladribine, clodronic acid, clofarabine, cobimetinib, copanlisib, crisantaspase, crizotinib, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, daratumumab, darbeopetin alfa, dabrafenib, dasatinib, daunorubicin, decitabine,

degarelix, denileukin diftitox, denosumab, depreotide, deslorelin, dianhydrogalactitol, dexrazoxane, dibrospidium chloride, dianhydrogalactitol, diclofenac, dinutuximab, docetaxel, dolasetron, doxifluridine, doxorubicin, doxorubicin + estrone, dronabinol, eculizumab, edrecolomab, elliptinium acetate, elotuzumab, eltrombopag, endostatin, enocitabine, 5 enzalutamide, epirubicin, epitiostanol, epoetin alfa, epoetin beta, epoetin zeta, eptaplatin, eribulin, erlotinib, esomeprazole, estradiol, estramustine, ethinylestradiol, etoposide, everolimus, exemestane, fadrozole, fentanyl, filgrastim, fluoxymesterone, floxuridine, fludarabine, fluorouracil, flutamide, folinic acid, formestane, fosaprepitant, fotemustine, fulvestrant, gadobutrol, gadoteridol, gadoteric acid meglumine, gadoversetamide, gadoxetic 10 acid, gallium nitrate, ganirelix, gefitinib, gemcitabine, gemtuzumab, Glucarpidase, glutoxim, GM-CSF, goserelin, granisetron, granulocyte colony stimulating factor, histamine dihydrochloride, histrelin, hydroxycarbamide, I-125 seeds, lansoprazole, ibandronic acid, ibritumomab tiuxetan, ibrutinib, idarubicin, ifosfamide, imatinib, imiquimod, improsulfan, indisetron, incadronic acid, ingenol mebutate, interferon alfa, interferon beta, interferon 15 gamma, iobitridol, iobenguane (123I), iomeprol, ipilimumab, irinotecan, Itraconazole, ixabepilone, ixazomib, lanreotide, lansoprazole, lapatinib, lasocholine, lenalidomide, lenvatinib, lenograstim, lentinan, letrozole, leuprorelin, levamisole, levonorgestrel, levothyroxine sodium, lisuride, lobaplatin, lomustine, lonidamine, masoprocol, medroxyprogesterone, megestrol, melarsoprol, melphalan, mepitiostane, mercaptopurine, mesna, methadone, methotrexate, 20 methoxsalen, methylaminolevulinate, methylprednisolone, methyltestosterone, metirosine, mifamurtide, miltefosine, miriplatin, mitobronitol, mitoguazone, mitolactol, mitomycin, mitotane, mitoxantrone, mogamulizumab, molgramostim, mopidamol, morphine hydrochloride, morphine sulfate, nabilone, nabiximols, nafarelin, naloxone + pentazocine, naltrexone, nartograstim, necitumumab, nedaplatin, nelarabine, neridronic acid, netupitant/palonosetron, nivolumab, 25 pentetretotide, nilotinib, nilutamide, nimorazole, nimotuzumab, nimustine, nintedanib, nitracrine, nivolumab, obinutuzumab, octreotide, ofatumumab, olaparib, olaratumab, omacetaxine mepesuccinate, omeprazole, ondansetron, oprelvekin, orgotein, orlitimod, osimertinib, oxaliplatin, oxycodone, oxymetholone, ozogamicine, p53 gene therapy, paclitaxel, palbociclib, palifermin, palladium-103 seed, palonosetron, pamidronic acid, panitumumab, panobinostat, 30 pantoprazole, pazopanib, pegaspargase, PEG-epoetin beta (methoxy PEG-epoetin beta), pembrolizumab, pegfilgrastim, peginterferon alfa-2b, pembrolizumab, pemetrexed, pentazocine, pentostatin, peplomycin, Perflubutane, perfosfamide, Pertuzumab, picibanil, pilocarpine, pirarubicin, pixantrone, plerixafor, plicamycin, poliglusam, polyestradiol phosphate, polyvinylpyrrolidone + sodium hyaluronate, polysaccharide-K, pomalidomide, ponatinib, 35 porfimer sodium, pralatrexate, prednimustine, prednisone, procarbazine, procodazole, propranolol, quinagolide, rabeprazole, racotumomab, radium-223 chloride, radotinib, raloxifene, raltitrexed, ramosetron, ramucirumab, ranimustine, rasburicase, razoxane, refametinib , regorafenib, risedronic acid, rhenium-186 etidronate, rituximab, rolapitant,

romidepsin, romiplostim, romurtide, roniciclib, samarium (153Sm) lexidronam, sargramostim, satumomab, secretin, siltuximab, sipuleucel-T, sizofiran, sobuzoxane, sodium glycididazole, sonidegib, sorafenib, stanozolol, streptozocin, sunitinib, talaporfin, talimogene laherparepvec, tamibarotene, tamoxifen, tapentadol, tasonermin, teceleukin, technetium (99mTc) 5 nofetumomab merpentan, 99mTc-HYNIC-[Tyr3]-octreotide, tegafur, tegafur + gimeracil + oteracil, temoporfin, temozolomide, temsirolimus, teniposide, testosterone, tetrafosmin, thalidomide, thiotepa, thymalfasin, thyrotropin alfa, tioguanine, tocilizumab, topotecan, toremifene, tositumomab, trabectedin, trametinib, tramadol, trastuzumab, trastuzumab emtansine, treosulfan, tretinoin, trifluridine + tipiracil, trilostane, triptorelin, trametinib, 10 trofosfamide, thrombopoietin, tryptophan, ubenimex, valatinib, valrubicin, vandetanib, vapreotide, vemurafenib, vinblastine, vincristine, vindesine, vinflunine, vinorelbine, vismodegib, vorinostat, vorozole, yttrium-90 glass microspheres, zinostatin, zinostatin stimalamer, zoledronic acid, zorubicin.

15 The compounds of the invention can further be combined with other reagents targeting the immune system, such as immune checkpoint inhibitors. Compositions comprising a PD-1/-L1 axis antagonist and an AHR antagonist and methods of using the same are provided herein. Data presented herein demonstrate that a combination of AHR inhibition and blockade of the PD-1/-L1 axis reduces the growth of tumor cells in more than an additive manner. PD-1, along 20 with its ligands PD-L1 and PD-L2, function as negative regulators of T cell activation. AHR suppresses immune cell function while increasing cancer cell proliferation and motility. PD-L1 is overexpressed in many cancers and overexpression of PD-1 often occurs concomitantly in tumor infiltrating T cells. Thus results in attenuation of T cell activation and evasion of immune surveillance, which contributes to impaired antitumor immune responses. (Keir M E et al. 25 (2008) Annu. Rev. Immunol. 26:677). Simultaneously targeting both the PD-1/-L1 axis and AHR enhances antitumor immune responses in more than an additive manner, leading to reduction of tumor growth that is unexpected. In some experiments, the resulting effect is greater than the expected or calculated additive effect of the individual components given separately. Thus, compositions comprising a PD-1/-L1 axis antagonist and an AHR antagonist 30 are surprisingly effective in enhancing an immune response and in the treatment of cancer.

In addition, the inventive compounds can also be used as a therapeutic in a variety of other disorders wherein AHR is involved such as, cardiovascular and lung diseases.

Accordingly, the compounds according to the invention are suitable for the treatment and/or prophylaxis in particular of cardiovascular, inflammatory and fibrotic disorders and of renal 35 disorders, in particular of acute and chronic renal insufficiency, and also of acute and chronic renal failure.

Accordingly, the compounds according to the invention can be used in medicaments for the treatment and/or prophylaxis of cardiovascular, inflammatory and fibrotic disorders, renal disorders, in particular of acute and chronic renal insufficiency, and also of acute and chronic renal failure.

- 5 For the purpose of the present invention the term renal insufficiency comprises both acute and chronic manifestations of renal insufficiency, and also underlying or related renal disorders such as diabetic and non-diabetic nephropathies, hypertensive nephropathies, ischaemic renal disorders, renal hypoperfusion, intradialytic hypotension, obstructive uropathy, renal stenoses, glomerulopathies, glomerulonephritis (such as, for example, primary glomerulonephritides;
10 minimal change glomerulonephritis (lipoidnephrosis); membranous glomerulonephritis; focal segmental glomerulosclerosis (FSGS); membrane-proliferative glomerulonephritis; crescentic glomerulonephritis; mesangioproliferative glomerulonephritis (IgA nephritis, Berger's disease); post-infectious glomerulonephritis; secondary glomerulonephritides: diabetes mellitus, lupus erythematosus, amyloidosis, Goodpasture syndrome, Wegener granulomatosis, Henoch-Schönlein purpura, microscopic polyangiitis, acute glomerulonephritis, pyelonephritis (for
15 example as a result of: urolithiasis, benign prostate hyperplasia, diabetes, malformations, abuse of analgesics, Crohn's disease), glomerulosclerosis, arteriolonecrose of the kidney, tubulointerstitial diseases, nephropathic disorders such as primary and congenital or aquired renal disorder, Alport syndrome, nephritis, immunological kidney disorders such as kidney
20 transplant rejection and immunocomplex-induced renal disorders, nephropathy induced by toxic substances, nephropathy induced by contrast agents, diabetic and non-diabetic nephropathy, renal cysts, nephrosclerosis, hypertensive nephrosclerosis and nephrotic syndrome which can be characterized diagnostically, for example by abnormally reduced creatinine and/or water excretion, abnormally elevated blood concentrations of urea, nitrogen,
25 potassium and/or creatinine, altered activity of renal enzymes, for example glutamyl synthetase, altered urine osmolarity or urine volume, elevated microalbuminuria, macroalbuminuria, lesions on glomerulae and arterioles, tubular dilatation, hyperphosphataemia and/or the need for dialysis. The present invention also comprises the use of the compounds according to the invention for the treatment and/or prophylaxis of
30 sequelae of renal insufficiency, for example pulmonary oedema, heart failure, uremia, anemia, electrolyte disturbances (for example hypercalemia, hyponatremia) and disturbances in bone and carbohydrate metabolism.

The present invention also comprises the use of the compounds according to the invention for the treatment and/or prevention of sequelae of renal insufficiency, for example pulmonary
35 oedema, heart failure, uraemia, anaemia, electrolyte disturbances (for example hyperkalaemia, hyponatraemia) and disturbances in bone and carbohydrate metabolism.

The compounds according to the invention are further suitable for the treatment and/or prevention of polycystic kidney disease (PCKD) and of the syndrome of inappropriate ADH secretion (SIADH).

5 Furthermore, the compounds according to the invention are also suitable for the treatment and/or prophylaxis of metabolic syndrome, hypertension, resistant hypertension, acute and chronic heart failure, coronary heart disease, stable and unstable angina pectoris, peripheral and cardiac vascular disorders, arrhythmias, atrial and ventricular arrhythmias and impaired
10 conduction, for example atrioventricular blocks degrees I-III (AB block I-III), supraventricular tachyarrhythmia, atrial fibrillation, atrial flutter, ventricular fibrillation, ventricular flutter, ventricular tachyarrhythmia, Torsade de pointes tachycardia, atrial and ventricular extrasystoles, AV-junctional extrasystoles, sick sinus syndrome, syncope, AV-nodal re-entry tachycardia, Wolff-Parkinson-White syndrome, of acute coronary syndrome (ACS), autoimmune cardiac disorders (pericarditis, endocarditis, valvulitis, aortitis, cardiomyopathies), shock such as cardiogenic shock, septic shock and anaphylactic shock, aneurysms, boxer
15 cardiomyopathy (premature ventricular contraction (PVC)), for treatment and/or prophylaxis of thromboembolic disorders and ischaemias such as myocardial ischaemia, myocardial infarction, stroke, cardiac hypertrophy, transient and ischaemic attacks, preeclampsia, inflammatory cardiovascular disorders, spasms of the coronary arteries and peripheral arteries, oedema formation, for example pulmonary oedema, cerebral oedema, renal oedema or
20 oedema caused by heart failure, peripheral circulatory disturbances, reperfusion damage, arterial and venous thromboses, myocardial insufficiency, endothelial dysfunction, to prevent restenoses, for example after thrombolysis therapies, percutaneous transluminal angioplasties (PTA), transluminal coronary angioplasties (PTCA), heart transplants and bypass operations, and also micro- and macrovascular damage (vasculitis), increased levels of fibrinogen and of
25 low-density lipoprotein (LDL) and increased concentrations of plasminogen activator inhibitor 1 (PAI-1), and also for treatment and/or prophylaxis of erectile dysfunction and female sexual dysfunction.

In addition, the compounds according to the invention are also suitable for treatment and/or prophylaxis of asthmatic disorders, pulmonary arterial hypertension (PAH) and other forms of
30 pulmonary hypertension (PH) including left-heart disease, HIV, sickle cell anaemia, thromboembolisms (CTEPH), sarcoidosis, COPD or pulmonary fibrosis-associated pulmonary hypertension, chronic-obstructive pulmonary disease (COPD), acute respiratory distress syndrome (ARDS), acute lung injury (ALI), alpha-1-antitrypsin deficiency (AATD), pulmonary fibrosis, pulmonary emphysema (for example pulmonary emphysema induced by cigarette
35 smoke) and cystic fibrosis (CF).

The compounds described in the present invention are also active compounds for control of central nervous system disorders characterized by disturbances of the NO/cGMP system. They are suitable in particular for improving perception, concentration, learning or memory

after cognitive impairments like those occurring in particular in association with situations/diseases/syndromes such as mild cognitive impairment, age-associated learning and memory impairments, age-associated memory losses, vascular dementia, craniocerebral trauma, stroke, dementia occurring after strokes (post stroke dementia), post-traumatic
5 craniocerebral trauma, general concentration impairments, concentration impairments in children with learning and memory problems, Alzheimer's disease, Lewy body dementia, dementia with degeneration of the frontal lobes including Pick's syndrome, Parkinson's disease, progressive dementia with corticobasal degeneration, amyolateral sclerosis (ALS), Huntington's disease, demyelination, multiple sclerosis, thalamic degeneration, Creutzfeld-
10 Jacob dementia, HIV dementia, schizophrenia with dementia or Korsakoff's psychosis. They are also suitable for treatment and/or prophylaxis of central nervous system disorders such as states of anxiety, tension and depression, CNS-related sexual dysfunctions and sleep disturbances, and for controlling pathological disturbances of the intake of food, stimulants and addictive substances.

15 The compounds according to the invention are furthermore also suitable for controlling cerebral blood flow and thus represent effective agents for controlling migraines. They are also suitable for the prophylaxis and control of sequelae of cerebral infarction (cerebral apoplexy) such as stroke, cerebral ischaemia and craniocerebral trauma. The compounds according to the invention can likewise be used for controlling states of pain and tinnitus.

20

The compounds according to the invention are also suitable for treatment and/or prophylaxis of fibrotic disorders of the internal organs, for example the lung, the heart, the kidney, the bone marrow and in particular the liver, and also dermatological fibroses and fibrotic eye disorders. In the context of the present invention, the term fibrotic disorders includes in particular the
25 following terms: hepatic fibrosis, cirrhosis of the liver, pulmonary fibrosis, endomyocardial fibrosis, nephropathy, glomerulonephritis, interstitial renal fibrosis, fibrotic damage resulting from diabetes, bone marrow fibrosis and similar fibrotic disorders, scleroderma, morphea, keloids, hypertrophic scarring (also following surgical procedures), naevi, diabetic retinopathy, proliferative vitreoretinopathy and disorders of the connective tissue (for example sarcoidosis).

30 The compounds according to the invention are also suitable for controlling postoperative scarring, for example as a result of glaucoma operations.

The compounds according to the invention can also be used cosmetically for ageing and keratinized skin.

Moreover, the compounds according to the invention are suitable for treatment and/or
35 prophylaxis of hepatitis, neoplasms, osteoporosis, glaucoma and gastroparesis.

The present invention further provides for the use of the compounds according to the invention for treatment and/or prophylaxis of disorders, especially the disorders mentioned above.

The present invention further provides for the use of the compounds according to the invention for the treatment and/or prophylaxis of chronic renal disorders, acute and chronic renal insufficiency, diabetic, inflammatory or hypertensive nephropathies, fibrotic disorders, cardiac insufficiency, angina pectoris, hypertension, pulmonary hypertension, ischemias, vascular disorders, thromboembolic disorders, arteriosclerosis, sickle cell anemia, erectile dysfunction, benign prostate hyperplasia, dysuria associated with benign prostate hyperplasia, Huntington, dementia, Alzheimer and Creutzfeld-Jakob.

The present invention further provides a method for treatment and/or prophylaxis of disorders, in particular the disorders mentioned above, using an effective amount of at least one of the compounds according to the invention.

The present invention further provides a method for the treatment and/or prophylaxis of chronic renal disorders, acute and chronic renal insufficiency, diabetic, inflammatory or hypertensive nephropathies, fibrotic disorders, cardiac insufficiency, angina pectoris, hypertension, pulmonary hypertension, ischemias, vascular disorders, thromboembolic disorders, arteriosclerosis, sickle cell anemia, erectile dysfunction, benign prostate hyperplasia, dysuria associated with benign prostate hyperplasia, Huntington, dementia, Alzheimer and Creutzfeld-Jakob.

In another embodiment, the inventive compounds can also be used to treat or to prevent uterine fibroids (uterine leiomyoma or uterine myoma) in women.

Uterine fibroids are benign tumors of the myometrium, the smooth muscle layer of the uterus. Uterine fibroids grow slowly during a women's life, and their growth is dependent on the female sexual hormones estradiol and progesterone [Kawaguchi K et al. Immunohistochemical analysis of oestrogen receptors, progesterone receptors and Ki-67 in leiomyoma and myometrium during the menstrual cycle and pregnancy *Virchows Arch A Pathol Anat Histopathol.* 1991;419(4):309-15.], therefore the highest prevalence of uterine fibroids with approx. 70% and >80% in white and afro-american women, respectively, is found from 35 years of age onwards to menopause, when they shrink due to reduced hormone levels [Baird DD et al. High cumulative incidence of uterine leiomyoma in black and white women: Ultrasound evidence *Am J Obstet Gynecol.* 2003 Jan;188(1):100-7.]. Approx 30% and 45% of white and afro-american women, respectively, do show clinically relevant symptoms due to their fibroids, which are heavy menstrual bleeding and pain, which is related to the menstrual cycle [David M et al. Myoma-associated pain frequency and intensity: a retrospective evaluation of 1548 myoma patients. *Eur J Obstet Gynecol Reprod Biol.* 2016 Apr;199:137-40]. Heavy menstrual bleeding in this respect is defined by a blood loss of more than 80 mL in a menstrual bleeding period [Fraser IS et al. The FIGO Recommendations on Terminologies and Definitions for Normal and Abnormal Uterine Bleeding, *Semin Reprod Med* 2011; 29(5): 383-390]. Submucosal position of the uterine fibroids, e.g. those located directly below the

endometrium, seems to have an even more severe effect on uterine bleeding, which may result in anemia in affected women [Yang JH et al. Impact of submucous myoma on the severity of anemia. *Fertil Steril*. 2011 Apr;95(5):1769-72]. Furthermore, uterine fibroids, due to their symptoms, do severely affect the quality of life of affected women [Downes E et al. The
5 burden of uterine fibroids in five European countries. *Eur J Obstet Gynecol Reprod Biol*. 2010 Sep;152(1):96-102].

So far, it is not understood how uterine fibroids do cause heavy menstrual bleeding. Disregulated genes in uterine fibroids, in comparison to normal myometrium, can give a hint to understand the underlying mechanisms. In published and internal studies, we found TDO2,
10 Tryptophan 2,3-dioxygenase, being highly upregulated [Tsibris JC et al. Insights from gene arrays on the development and growth regulation of uterine leiomyomata. *Fertil Steril*. 2002 Jul;78(1):114-21.]. TDO2 metabolizes the substrate L-Tryptophan to L-Kynurenine, which can be further metabolized to kynurenic acid. Both, L-Kynurenine and Kynurenic acid are physiological ligands and activators for the arylhydrocarbon receptor AHR [Opitz CA et al. An
15 endogenous tumour-promoting ligand of the human aryl hydrocarbon receptor *Nature*. 2011 Oct 5;478(7368):197-203].

L-Kynurenine controls at least two physiological processes which are dysregulated in uterine fibroids. L-Kynurenine, synthesized by an upregulation of IDO (Indoleamine-2,3-dioxygenase) or TDO2, and acting via the AHR receptor, suppresses the immune system and thus prevents
20 immune cells from recognizing and clearing the tumor cells [Munn DH Blocking IDO activity to enhance anti-tumor immunity. *Front Biosci (Elite Ed)*. 2012 Jan 1;4:734-45]. Furthermore, an upregulation of L-Kynurenine leads to a vasodilation of vessels, and thus can directly increase blood loss and bleeding [Wang Y et al. Kynurenine is an endothelium-derived relaxing factor produced during inflammation *Nature Medicine* 16, 279–285 (2010)].

25 In summary, the upregulation of L-Kynurenine through activation of its physiological receptor AHR seems to support uterine fibroid growth by local suppression of the immune system, and might cause heavy menstrual bleeding by vasodilation of endometrial vessels in proximity to the tumor.

Therefore, a systemic or local application of compounds from the present invention inhibiting
30 activation of the AHR and thus blocking the effect of uterine fibroid derived L-Kynurenine presents a new and valid treatment option for uterine fibroids.

Compounds of the present invention can be utilized to inhibit, block, reduce or decrease AHR activation by exogenous and/or endogenous ligands for the reduction of tumour growth and the
35 modulation of dysregulated immune responses e.g. to block immunosuppression and increase immune cell activation and infiltration in the context of cancer and cancer immunotherapy; This method comprises administering to a mammal in need thereof, including a human, an amount

of a compound of this invention, or a pharmaceutically acceptable salt, isomer, polymorph, metabolite, hydrate, solvate or ester thereof; which is effective to treat the disorder.

The present invention also provides methods of treating a variety of other disorders wherein AHR is involved such as, but not limited to, inflammation, vaccination for infection & cancer, viral infections, obesity and diet-induced obesity, adiposity, metabolic disorders, hepatic steatosis and uterine fibroids.

These disorders have been well characterized in humans, but also exist with a similar etiology in other mammals, and can be treated by administering pharmaceutical compositions of the present invention.

10 The term "treating" or "treatment" as used in the present text is used conventionally, e.g., the management or care of a subject for the purpose of combating, alleviating, reducing, relieving, improving the condition of a disease or disorder, such as liquid and solid tumours.

In accordance with a further aspect, the present invention covers compounds of general formula (I), as described *supra*, or stereoisomers, tautomers, N-oxides, hydrates, solvates, and salts thereof, particularly pharmaceutically acceptable salts thereof, or mixtures of same, for use in the treatment or prophylaxis of diseases, in particular cancer or conditions with dysregulated immune responses or other disorders associated with aberrant AHR signaling.

The pharmaceutical activity of the compounds according to the invention can be explained by their activity as AHR inhibitors.

20 In accordance with a further aspect, the present invention covers the use of compounds of general formula (I), as described *supra*, or stereoisomers, tautomers, N-oxides, hydrates, solvates, and salts thereof, particularly pharmaceutically acceptable salts thereof, or mixtures of same, for the treatment or prophylaxis of diseases, in particular cancer or conditions with dysregulated immune responses or other disorders associated with aberrant AHR signaling, particularly liquid and solid tumours.

In accordance with a further aspect, the present invention covers the use of a compound of formula (I), described *supra*, or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, particularly a pharmaceutically acceptable salt thereof, or a mixture of same, for the prophylaxis or treatment of diseases, in particular cancer or conditions with dysregulated immune responses or other disorders associated with aberrant AHR signaling, particularly liquid and solid tumours.

30 In accordance with a further aspect, the present invention covers the use of compounds of general formula (I), as described *supra*, or stereoisomers, tautomers, N-oxides, hydrates, solvates, and salts thereof, particularly pharmaceutically acceptable salts thereof, or mixtures of same, in a method of treatment or prophylaxis of diseases, in particular cancer or conditions with dysregulated immune responses or other disorders associated with aberrant AHR signaling, particularly liquid and solid tumours.

In accordance with a further aspect, the present invention covers the use of compounds of general formula (I), as described *supra*, or stereoisomers, tautomers, N-oxides, hydrates, solvates, and salts thereof, particularly pharmaceutically acceptable salts thereof, or mixtures of same, in a method of treatment or prophylaxis of diseases, in particular cancer or conditions with dysregulated immune responses or other disorders associated with aberrant AHR signaling, particularly liquid and solid tumours.

In accordance with a further aspect, the present invention covers use of a compound of general formula (I), as described *supra*, or stereoisomers, tautomers, N-oxides, hydrates, solvates, and salts thereof, particularly pharmaceutically acceptable salts thereof, or mixtures of same, for the preparation of a pharmaceutical composition, preferably a medicament, for the prophylaxis or treatment of diseases, in particular cancer or conditions with dysregulated immune responses or other disorders associated with aberrant AHR signaling, particularly liquid and solid tumours.

In accordance with a further aspect, the present invention covers a method of treatment or prophylaxis of diseases, in particular cancer or conditions with dysregulated immune responses or other disorders associated with aberrant AHR signaling, particularly liquid and solid tumours, using an effective amount of a compound of general formula (I), as described *supra*, or stereoisomers, tautomers, N-oxides, hydrates, solvates, and salts thereof, particularly pharmaceutically acceptable salts thereof, or mixtures of same.

In accordance with a further aspect, the present invention covers pharmaceutical compositions, in particular a medicament, comprising a compound of general formula (I), as described *supra*, or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, a salt thereof, particularly a pharmaceutically acceptable salt, or a mixture of same, and one or more excipients), in particular one or more pharmaceutically acceptable excipient(s). Conventional procedures for preparing such pharmaceutical compositions in appropriate dosage forms can be utilized.

The present invention furthermore covers pharmaceutical compositions, in particular medicaments, which comprise at least one compound according to the invention, conventionally together with one or more pharmaceutically suitable excipients, and to their use for the above mentioned purposes.

It is possible for the compounds according to the invention to have systemic and/or local activity. For this purpose, they can be administered in a suitable manner, such as, for example, via the oral, parenteral, pulmonary, nasal, sublingual, lingual, buccal, rectal, vaginal, dermal, transdermal, conjunctival, otic route or as an implant or stent.

For these administration routes, it is possible for the compounds according to the invention to be administered in suitable administration forms.

For oral administration, it is possible to formulate the compounds according to the invention to dosage forms known in the art that deliver the compounds of the invention rapidly and/or in a

modified manner, such as, for example, tablets (uncoated or coated tablets, for example with enteric or controlled release coatings that dissolve with a delay or are insoluble), orally-disintegrating tablets, films/wafers, films/lyophilisates, capsules (for example hard or soft gelatine capsules), sugar-coated tablets, granules, pellets, powders, emulsions, suspensions, aerosols or solutions. It is possible to incorporate the compounds according to the invention in crystalline and/or amorphised and/or dissolved form into said dosage forms.

Parenteral administration can be effected with avoidance of an absorption step (for example intravenous, intraarterial, intracardial, intraspinal or intralumbal) or with inclusion of absorption (for example intramuscular, subcutaneous, intracutaneous, percutaneous or intraperitoneal).

Administration forms which are suitable for parenteral administration are, inter alia, preparations for injection and infusion in the form of solutions, suspensions, emulsions, lyophilisates or sterile powders.

Examples which are suitable for other administration routes are pharmaceutical forms for inhalation [inter alia powder inhalers, nebulizers], nasal drops, nasal solutions, nasal sprays; tablets/films/wafers/capsules for lingual, sublingual or buccal administration; suppositories; eye drops, eye ointments, eye baths, ocular inserts, ear drops, ear sprays, ear powders, ear-rinses, ear tampons; vaginal capsules, aqueous suspensions (lotions, mixturae agitandae), lipophilic suspensions, emulsions, ointments, creams, transdermal therapeutic systems (such as, for example, patches), milk, pastes, foams, dusting powders, implants or stents.

The compounds according to the invention can be incorporated into the stated administration forms. This can be effected in a manner known per se by mixing with pharmaceutically suitable excipients. Pharmaceutically suitable excipients include, inter alia,

fillers and carriers (for example cellulose, microcrystalline cellulose (such as, for example, Avicel[®]), lactose, mannitol, starch, calcium phosphate (such as, for example, Di-Cafos[®]),

ointment bases (for example petroleum jelly, paraffins, triglycerides, waxes, wool wax, wool wax alcohols, lanolin, hydrophilic ointment, polyethylene glycols),

bases for suppositories (for example polyethylene glycols, cacao butter, hard fat),

solvents (for example water, ethanol, isopropanol, glycerol, propylene glycol, medium chain-length triglycerides fatty oils, liquid polyethylene glycols, paraffins),

surfactants, emulsifiers, dispersants or wetters (for example sodium dodecyl sulfate), lecithin, phospholipids, fatty alcohols (such as, for example, Lanette[®]), sorbitan fatty acid esters (such

as, for example, Span[®]), polyoxyethylene sorbitan fatty acid esters (such as, for example, Tween[®]), polyoxyethylene fatty acid glycerides (such as, for example, Cremophor[®]),

polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, glycerol fatty acid esters,

poloxamers (such as, for example, Pluronic[®]),

- buffers, acids and bases (for example phosphates, carbonates, citric acid, acetic acid, hydrochloric acid, sodium hydroxide solution, ammonium carbonate, trometamol, triethanolamine),
- isotonicity agents (for example glucose, sodium chloride),
- 5 adsorbents (for example highly-disperse silicas),
- viscosity-increasing agents, gel formers, thickeners and/or binders (for example polyvinylpyrrolidone, methylcellulose, hydroxypropylmethylcellulose, hydroxypropylcellulose, carboxymethylcellulose-sodium, starch, carbomers, polyacrylic acids (such as, for example, Carbopol[®]); alginates, gelatine),
- 10 disintegrants (for example modified starch, carboxymethylcellulose-sodium, sodium starch glycolate (such as, for example, Explotab[®]), cross-linked polyvinylpyrrolidone, croscarmellose-sodium (such as, for example, AcDiSol[®]),
- flow regulators, lubricants, glidants and mould release agents (for example magnesium stearate, stearic acid, talc, highly-disperse silicas (such as, for example, Aerosil[®]),
- 15 coating materials (for example sugar, shellac) and film formers for films or diffusion membranes which dissolve rapidly or in a modified manner (for example polyvinylpyrrolidones (such as, for example, Kollidon[®]), polyvinyl alcohol, hydroxypropylmethylcellulose, hydroxypropylcellulose, ethylcellulose, hydroxypropylmethylcellulose phthalate, cellulose acetate, cellulose acetate phthalate, polyacrylates, polymethacrylates such as, for example,
- 20 Eudragit[®]),
- capsule materials (for example gelatine, hydroxypropylmethylcellulose),
- synthetic polymers (for example polylactides, polyglycolides, polyacrylates, polymethacrylates (such as, for example, Eudragit[®]), polyvinylpyrrolidones (such as, for example, Kollidon[®]), polyvinyl alcohols, polyvinyl acetates, polyethylene oxides, polyethylene glycols and their
- 25 copolymers and blockcopolymers),
- plasticizers (for example polyethylene glycols, propylene glycol, glycerol, triacetine, triacetyl citrate, dibutyl phthalate),
- penetration enhancers,
- stabilisers (for example antioxidants such as, for example, ascorbic acid, ascorbyl palmitate,
- 30 sodium ascorbate, butylhydroxyanisole, butylhydroxytoluene, propyl gallate),
- preservatives (for example parabens, sorbic acid, thiomersal, benzalkonium chloride, chlorhexidine acetate, sodium benzoate),
- colourants (for example inorganic pigments such as, for example, iron oxides, titanium dioxide),
- 35 flavourings, sweeteners, flavour- and/or odour-masking agents.

The present invention furthermore relates to a pharmaceutical composition which comprise at least one compound according to the invention, conventionally together with one or more pharmaceutically suitable excipient(s), and to their use according to the present invention.

5 In accordance with another aspect, the present invention covers pharmaceutical combinations, in particular medicaments, comprising at least one compound of general formula (I) of the present invention and at least one or more further active ingredients, in particular for the treatment and/or prophylaxis of cancer or conditions with dysregulated immune responses or other disorders associated with aberrant AHR signalinggeneric name disorders, particularly liquid and solid tumours.

10 The term "combination" in the present invention is used as known to persons skilled in the art, it being possible for said combination to be a fixed combination, a non-fixed combination or a kit-of-parts.

A "fixed combination" in the present invention is used as known to persons skilled in the art and is defined as a combination wherein, for example, a first active ingredient, such as one or
15 more compounds of general formula (I) of the present invention, and a further active ingredient are present together in one unit dosage or in one single entity. One example of a "fixed combination" is a pharmaceutical composition wherein a first active ingredient and a further active ingredient are present in admixture for simultaneous administration, such as in a formulation. Another example of a "fixed combination" is a pharmaceutical combination
20 wherein a first active ingredient and a further active ingredient are present in one unit without being in admixture.

A non-fixed combination or "kit-of-parts" in the present invention is used as known to persons skilled in the art and is defined as a combination wherein a first active ingredient and a further active ingredient are present in more than one unit. One example of a non-fixed combination or
25 kit-of-parts is a combination wherein the first active ingredient and the further active ingredient are present separately. It is possible for the components of the non-fixed combination or kit-of-parts to be administered separately, sequentially, simultaneously, concurrently or chronologically staggered.

Based upon standard laboratory techniques known to evaluate compounds useful for the
30 treatment of cancer or conditions with dysregulated immune responses or other disorders associated with aberrant AHR signaling, by standard toxicity tests and by standard pharmacological assays for the determination of treatment of the conditions identified above in mammals, and by comparison of these results with the results of known active ingredients or medicaments that are used to treat these conditions, the effective dosage of the compounds of
35 the present invention can readily be determined for treatment of each desired indication. The amount of the active ingredient to be administered in the treatment of one of these conditions can vary widely according to such considerations as the particular compound and dosage unit

employed, the mode of administration, the period of treatment, the age and sex of the patient treated, and the nature and extent of the condition treated.

The total amount of the active ingredient to be administered will generally range from about 0.001 mg/kg to about 200 mg/kg body weight per day, and preferably from about 0.01 mg/kg to about 20 mg/kg body weight per day. Clinically useful dosing schedules will range from one to three times a day dosing to once every four weeks dosing. In addition, it is possible for "drug holidays", in which a patient is not dosed with a drug for a certain period of time, to be beneficial to the overall balance between pharmacological effect and tolerability. It is possible for a unit dosage to contain from about 0.5 mg to about 1500 mg of active ingredient, and can be administered one or more times per day or less than once a day. The average daily dosage for administration by injection, including intravenous, intramuscular, subcutaneous and parenteral injections, and use of infusion techniques will preferably be from 0.01 to 200 mg/kg of total body weight. The average daily rectal dosage regimen will preferably be from 0.01 to 200 mg/kg of total body weight. The average daily vaginal dosage regimen will preferably be from 0.01 to 200 mg/kg of total body weight. The average daily topical dosage regimen will preferably be from 0.1 to 200 mg administered between one to four times daily. The transdermal concentration will preferably be that required to maintain a daily dose of from 0.01 to 200 mg/kg. The average daily inhalation dosage regimen will preferably be from 0.01 to 100 mg/kg of total body weight.

Of course the specific initial and continuing dosage regimen for each patient will vary according to the nature and severity of the condition as determined by the attending diagnostician, the activity of the specific compound employed, the age and general condition of the patient, time of administration, route of administration, rate of excretion of the drug, drug combinations, and the like. The desired mode of treatment and number of doses of a compound of the present invention or a pharmaceutically acceptable salt or ester or composition thereof can be ascertained by those skilled in the art using conventional treatment tests.

EXPERIMENTAL SECTION

NMR peak forms are stated as they appear in the spectra, possible higher order effects have not been considered. The multiplicities are stated according to the signal form which appears in the spectrum, NMR-spectroscopic effects of a higher order were not taken into consideration. Multiplicity of the NMR signals: s = singlet, d = doublet, t = triplet, q = quartet, qi = quintet, b = broad signal, m = multiplet. NMR signals: shift in ppm. Combinations of multiplicity could be e.g. dd = doublet from doublet.

Chemical names were generated using the ACD/Name software from ACD/Labs. In some cases generally accepted names of commercially available reagents were used in place of ACD/Name generated names.

Table 1 lists the abbreviations used in this paragraph and in the Examples section as far as they are not explained within the text body. Other abbreviations have their meanings customary *per se* to the skilled person.

Table 1: Abbreviations

	ACN	acetonitrile
10	AcOH	acetic acid
	BPR	Back Pressure Regulator
	CDCl ₃	deuteriochloroform
	DAD	diode array detector
	DEA	diethylamine
15	DMF	N,N-dimethylformamide
	DMSO-d ₆	deuterated dimethyl sulfoxide
	DMSO	dimethyl sulfoxide
	ELSD	evaporative light scattering detector
	ESIpos	electrospray ionization positive
20	Expl.	example
	HATU	(7-aza-1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
	HBTU	O-benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate
	HCl	hydrochloric acid
25	HPLC	high-pressure liquid chromatography
	KA	kynurenic acid
	LCMS	liquid chromatography coupled with mass spectrometry
	LPS	lipopolysaccharide
	μL	microliter
30	mL	milliliter
	min	minute(s)
	MTBE	methyl tert-butyl ether
	p	pressure
	PBMC	peripheral blood mononuclear cells
35	PyBOB	(benzotriazol-1-yl)oxytripyrrolidinophosphonium hexafluorophosphate
	RP-HPLC	reverse-phase high-pressure liquid chromatography
	Rt	retention time

rt	room temperature
sat.	saturated
T3P	2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphorinane 2,4,6-trioxide
THF	tetrahydrofuran
5 TFA	trifluoroacetic acid
TLC	thin layer chromatography
TNF α	tumour necrosis factor alpha
μ M	micromolar
UPLC	Ultra high performance chromatography

10

The various aspects of the invention described in this application are illustrated by the following examples which are not meant to limit the invention in any way.

The example testing experiments described herein serve to illustrate the present invention and the invention is not limited to the examples given.

15

EXPERIMENTAL SECTION - GENERAL PART

All reagents, for which the synthesis is not described in the experimental part, are either commercially available, or are known compounds or may be formed from known compounds by known methods by a person skilled in the art.

20

The compounds and intermediates produced according to the methods of the invention may require purification. Purification of organic compounds is well known to the person skilled in the art and there may be several ways of purifying the same compound. In some cases, no purification may be necessary. In some cases, the compounds may be purified by
25 crystallization. In some cases, impurities may be stirred out using a suitable solvent. In some cases, the compounds may be purified by chromatography, particularly flash column chromatography, using for example prepacked silica gel cartridges, e.g. Biotage SNAP cartridges KP-Sil[®] or KP-NH[®] in combination with a Biotage autopurifier system (SP4[®] or Isolera Four[®]) and eluents such as gradients of hexane/ethyl acetate or
30 dichloromethane/methanol. In some cases, the compounds may be purified by preparative HPLC using for example a Waters autopurifier equipped with a diode array detector and/or on-line electrospray ionization mass spectrometer in combination with a suitable prepacked reverse phase column and eluents such as gradients of water and acetonitrile which may contain additives such as trifluoroacetic acid, formic acid or aqueous ammonia.

35

In some cases, purification methods as described above can provide those compounds of the present invention which possess a sufficiently basic or acidic functionality in the form of a salt, such as, in the case of a compound of the present invention which is sufficiently basic, a trifluoroacetate or formate salt for example, or, in the case of a compound of the present

invention which is sufficiently acidic, an ammonium salt for example. A salt of this type can either be transformed into its free base or free acid form, respectively, by various methods known to the person skilled in the art, or be used as salts in subsequent biological assays. It is to be understood that the specific form (e.g. salt, free base etc.) of a compound of the present invention as isolated and as described herein is not necessarily the only form in which said compound can be applied to a biological assay in order to quantify the specific biological activity.

UPLC methods

10

Method 1: Instrument: Waters Acquity UPLC-MS SQD 3001; column: Acquity UPLC BEH C18 1.7 μm , 50x2.1mm; eluent A: water + 0.1vol% formic acid (99%), eluent B: acetonitrile; gradient: 0-1.6 min 1-99% B, 1.6-2.0 min 99% B; flow 0.8 mL/min; temperature: 60°C; injection: 2 μL ; DAD scan: 210-400 nm; ELSD.

15

Method 2: Instrument: Waters Acquity UPLC-MS SQD 3001; column: Acquity UPLC BEH C18 1.7 μm , 50x2.1mm; eluent A: water + 0.2vol% aqueous ammonia (32%), eluent B: acetonitrile; gradient: 0-1.6 min 1-99% B, 1.6-2.0 min 99% B; flow 0.8 mL/min; temperature: 60°C; injection: 2 μL ; DAD scan: 210-400 nm; ELSD.

20

Preparative HPLC methods

Instrument: pump: Labomatic HD-3000, head HDK 280, lowpressure gradient module ND-B1000; manual injection valve: Rheodyne 3725i038; detector: Knauer Azura UVD 2.15; collector: Labomatic Labocol Vario-4000; column: Chromatorex RP C-18 10 μm , 125x30mm; eluent; gradient; UV-Detection.

25

Eluent acidic: solvent A: water + 0.1 vol% formic acid (99%), solvent B: acetonitrile; flow 150 mL/min.

30

Eluent basic: solvent A: water + 0.2 vol% ammonia (32%), solvent B: acetonitrile; flow 150 mL/min.

Method A: 0.00-0.50 min 1% B, 0.50–6.00 min 1-25% B, 6.00-6.10 min 25-100% B, 6.10-8.00 min 100% B

35

Method B: 0.00-0.50 min 10% B, 0.50–6.00 min 10-50% B, 6.00-6.10 min 50-100% B, 6.10-8.00 min 100% B


Method C: 0.00-0.50 min 15% B, 0.50–6.00 min 15-55% B, 6.00-6.10 min 55-100% B, 6.10-8.00 min 100% B

Method D: 0.00-0.50 min 30% B, 0.50–6.00 min 30-70% B, 6.00-6.10 min 70-100% B, 6.10-8.00 min 100% B

EXPERIMENTAL SECTION - INTERMEDIATES

Intermediate 1

Dimethyl [2-(4-chlorophenyl)-2-oxoethyl]malonate



10

A mixture of 15.0 g 2-bromo-1-(4-chlorophenyl)ethanone, 59 mL dimethyl malonate and 13.3 g potassium carbonate in 600 mL acetone was stirred at rt for 14 hours. After full conversion (TLC) the reaction mixture was poured into water, the organic phase was separated and washed with water and brine. After evaporation of the solvent in vacuo, the residue was purified by column chromatography (hexanes / ethyl acetate gradient to 50% ethyl acetate) to yield 17.0 g dimethyl [2-(4-chlorophenyl)-2-oxoethyl]propanedioate.


15

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ [ppm] = 3.63 (d, 2H); 3.68 (s, 6H); 3.97 (t, 1H); 7.58-7.64 (m, 2H); 7.98-8.04 (m, 2H).

20

Intermediate 2

Methyl 6-(4-chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3,4,5-tetrahydropyridazine-4-carboxylate



25


A mixture of 5.0 g of dimethyl [2-(4-chlorophenyl)-2-oxoethyl]malonate and 2.44 g (3-fluorophenyl)hydrazine in 100 mL of AcOH was stirred at 130°C for 5 hours. Then, the solvent

was removed in vacuo. The residue was purified by column chromatography (hexanes / ethyl acetate gradient with up to 40% ethyl acetate) to yield 3.2 g methyl 6-(4-chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3,4,5-tetrahydropyridazine-4-carboxylate.

- 5 $^1\text{H-NMR}$ (400 MHz, 25°C, DMSO- d_6): δ = 3.36-3.53 (m, 2H); 3.71 (s, 3H); 4.07 (dd, 1H); 7.17 (ddt, 1H); 7.38-7.57 (m, 5H); 7.85-7.90 (m, 2H).

Intermediate 3

Methyl 6-(4-chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylate



10


A mixture of 3.2 g of methyl 6-(4-chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3,4,5-tetrahydropyridazine-4-carboxylate and 3.58 g copper(II) chloride in 100 mL of acetonitrile was stirred at 90°C for 3 hours. After evaporation in vacuo, the residue was purified by column chromatography (hexanes / ethyl acetate gradient with up to 100% ethyl acetate) to yield 1.9 g

15 methyl 6-(4-chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylate.

$^1\text{H-NMR}$ (400 MHz, 25°C, DMSO- d_6): δ = 3.88 (s, 3H); 7.35 (ddt, 1H); 7.52-7.64 (m, 5H); 7.95-8.01 (m, 2H); 8.51 (s, 1H).

20 Intermediate 4

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid



A mixture of 1.9 g of methyl 6-(4-chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylate in 60 mL of acetonitrile was treated with 0.38 g lithium hydroxide, dissolved in


25 4.3 mL of water. The reaction mixture was stirred at room temperature for 5 hours. Then the pH value was adjusted to 6 with hydrochloric acid (10%). The solids were collected by filtration,

washed three times with water and dried in an oven to yield 1.5 g 6-(4-chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid.

¹H-NMR (300 MHz, 25°C, Methanol-d₄): δ = 7.32 (ddt, 1H); 7.49-7.62 (m, 5H); 7.92-7.97 (m, 2H); 8.02 (s, 1H).

Intermediate 5

Dimethyl {2-oxo-2-[4-(trifluoromethyl)phenyl]ethyl}malonate




10 Dimethyl malonate (9.894 g, 74.89 mmol) and potassium carbonate (7.763 g, 56.17 mmol) were added to acetone (140 mL). Under cooling (0-5 °C) a solution of 2-bromo-1-[4-(trifluoromethyl)phenyl]ethanone (10 g, 37.4 mmol) in acetone (60 mL) was added dropwise. It was stirred 2 h at 0-5 °C and at rt overnight. The volatile compounds were removed on a rotavap. Water and ethyl acetate were added, the layers were separated and the aqueous
15 phase was extracted twice with ethyl acetate. The combined organic layers were washed with concentrated aqueous sodium chloride solution, dried over magnesium sulfate and concentrated. The crude product was purified by flash chromatography (hexane / ethyl acetate) affording 8.03 g (67%) of the title product.

20 ¹H-NMR (400MHz, CHLOROFORM-d₃): δ [ppm] = 3.65 (d, 2H), 3.79 (s, 6H), 4.10 (t, 1H), 7.73 - 7.77 (m, 2H), 8.07 - 8.11 (m, 2H).

Intermediate 6

25 Methyl 2-(3-fluorophenyl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate



Step 1: Dimethyl {2-oxo-2-[4-(trifluoromethyl)phenyl]ethyl}malonate (4.00 g, 12.57 mmol) and (3-fluorophenyl)hydrazine hydrochloride (1:1) (3.065 g, 18.85 mmol) in acetic acid (50 mL) was stirred 8 h at 80 °C. Two of such batches were combined and concentrated on a rotavap. Hexane was added and it was removed on a rotavap. Water and ethyl acetate were added. The layers were separated and the aqueous phase was extracted four times with ethyl acetate. The combined organic layers were washed twice with water, dried over magnesium sulfate and concentrated to dryness affording 9.9 g (99.9%) of methyl 2-(3-fluorophenyl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3,4,5-tetrahydropyridazine-4-carboxylate which was used without further purification in the next step.


Step 2: Methyl 2-(3-fluorophenyl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3,4,5-tetrahydropyridazine-4-carboxylate (4.9 g, 7.46 mmol) was dissolved in acetonitrile (100 mL). Copper (II) chloride (3.007 g, 22.37 mmol) was added and it was stirred 9 h at 90 °C. The reaction mixture was allowed to reach rt. Two of such batches and a small batch (215 mg, 0.327 mmol) were combined and silica gel (60 g) was added. The volatiles were removed under vacuum. It was purified by flash chromatography (hexane / ethyl acetate) obtaining 2.3 g (23%) of the title compound.

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.89 (s, 3H), 7.33 - 7.39 (m, 1H), 7.54 - 7.58 (m, 1H), 7.58 - 7.64 (m, 2H), 7.88 (d, 2H), 8.17 (d, 2H), 8.57 (s, 1H).

20

Intermediate 7

2-(3-Fluorophenyl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid




Methyl 2-(3-fluorophenyl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate (2.3 g, 5.57 mmol) was dissolved in acetonitrile (57 mL). A solution of lithium hydroxide (400 mg, 16.71 mmol) in water (5.7 mL) was added at rt. It was stirred 24 h at rt. Water (10 mL) was added. 2N hydrochloric acid (9.56 mL) was added to adjust the pH to 4. It was stirred 1 h at rt. The precipitate was filtered off under suction, washed with water four times and dried under vacuum at 50 °C for 24 h yielding 1.89 g (85%) of the title compound which was used without further purification in the next step.

30

$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 7.31 - 7.37 (m, 1H), 7.52 - 7.56 (m, 1H), 7.57 - 7.63 (m, 2H), 7.85 (d, 2H), 8.16 (d, 2H), 8.24 (s, 1H).

5 Intermediate 8

Dimethyl [2-(4-chlorophenyl)-2-oxoethyl]malonate




2-Chloro-1-(4-chlorophenyl)ethanone (25 g, 107.1 mmol) was dissolved in acetone (500 mL). Then, dimethyl malonate (31.1 g, 235.4 mmol) and potassium carbonate (22.2 g, 160.6 mmol) were added at rt. It was stirred at rt overnight. The reaction mixture was reduced under vacuum to half its volume. Then, the residue was poured into water. The layers were separated and the aqueous phase was extracted twice with ethyl acetate. The combined organic layers were washed with water and concentrated aqueous sodium chloride solution, dried over sodium sulfate and concentrated. The crude product was purified by flash chromatography (silica gel, hexane / ethyl acetate, gradient) yielding 12.21 g (36%) of the title product.

$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 3.63 (d, 2H), 3.68 (s, 6H), 3.97 (t, 1H), 7.59 - 7.64 (m, 2H), 7.99 - 8.03 (m, 2H).

20 Intermediate 9

Methyl 6-(4-chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-2,3,4,5-tetrahydropyridazine-4-carboxylate




Dimethyl [2-(4-chlorophenyl)-2-oxoethyl]malonate (1360 mg, 4.78 mmol) and sodium acetate (1037 mg, 12.65 mmol) were dissolved in acetic acid (40 mL). Then, 4-hydrazino-1-methyl-1H-pyrazole dihydrochloride (780 mg, 4.22 mmol) was added portion wise. It was stirred for 1 h at rt and 20 h at 50°C. The reaction mixture was cooled down and concentrated on a rotary evaporator under reduced pressure. Ethyl acetate and water were added to dissolve the residue. Concentrated aqueous sodium hydrogen carbonate solution was added, the phases were separated, and the aqueous layer was extracted with ethyl acetate (four times with 80 mL). The combined organic layers were washed twice with water, dried over magnesium sulfate, and concentrated. The residue was purified by flash chromatography (silica gel, hexane / ethyl acetate, gradient) to afford 530 mg (36%) of the title product.

$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 3.35 - 3.46 (m, 2H), 3.68 (s, 3H), 3.85 (s, 3H), 4.03 (dd, 1H), 7.52 - 7.57 (m, 2H), 7.75 (d, 1H), 7.92 - 7.96 (m, 2H), 8.08 (s, 1H).

Intermediate 10

15 **Methyl 6-(4-chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-2,3-dihydropyridazine-4-carboxylate**




Methyl 6-(4-chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-2,3,4,5-tetrahydropyridazine-4-carboxylate (600 mg, 1.730 mmol) was dissolved in acetonitrile (40 mL). Copper dichloride (698 mg, 5.191 mmol) was added. It was stirred for 4 h at 90°C. It was cooled down and concentrated on a rotary evaporator. Water was added, the remaining solid was filtered by suction, washed five times with water, and dried under vacuum at 50°C to yield 741 mg of the title compound which was used without further purification in the next step.

25 $^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 3.88 (s, 3H), 3.91 (s, 3H), 7.59 (d, 2H), 8.05 - 8.13 (m, 3H), 8.44 (s, 1H), 8.52 (br s, 1H).

Intermediate 11

6-(4-Chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid




Methyl 6-(4-chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-2,3-dihydropyridazine-4-carboxylate (600 mg, 1.74 mmol) was dissolved in acetonitrile (60 mL). A solution of lithium hydroxide (125 mg, 5.221 mmol) in water (1.90 mL) was added at rt. It was stirred for 10 h at 40°C. Water was added and the pH was adjusted to 4 with 2N HCl. The precipitate was filtered off under suction, washed three times with water and dried under vacuum at 50°C obtaining 520 mg (90%) of the title compound which was used without further purification in the next step.

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.91 (s, 3H), 7.54 (br d, 2H), 7.78 (s, 1H), 8.00 - 8.07 (m, 3H), 8.41 (s, 1H).

Intermediate 12

15 Methyl 3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3,4,5-tetrahydropyridazine-4-carboxylate




Dimethyl {2-oxo-2-[4-(trifluoromethyl)phenyl]ethyl}malonate (5.68 g, 17.55 mmol) was dissolved in acetic acid (64 mL). A solution of hydrazine in THF (35 mL, 1.0M, 35 mmol) was added at rt. It was stirred for 3.5 h at 75°C. Then, a solution of hydrazine in THF (3.5 mL, 1.0M, 3.5 mmol) was added and stirring at 75°C was continued for 1 h. The reaction mixture was cooled down and water (0.6 L) was added. The precipitate was filtered off under suction, washed with water and dried under vacuum at 50°C yielding 4.06 g (76%) of the title compound which was used without further purification in the next step.

$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 3.23 (dd, 1H), 3.28 - 3.36 (m, 1H and water signal), 3.68 (s, 3H), 3.79 (dd, 1H), 7.80 (d, 2H), 7.96 (d, 2H), 11.43 (s, 1H).

Intermediate 13

5 Methyl 3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate




Methyl 3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3,4,5-tetrahydropyridazine-4-carboxylate (4.06 g, 13.52 mmol) was dissolved in acetonitrile (180 mL). Copper dichloride (4.55 g, 33.81 mmol) was added and it was stirred for 2.5 h at 90°C. The reaction mixture was cooled down and concentrated on a rotary evaporator to half its volume. Water was added (350 mL) and the reaction mixture was stirred for 10 min. The precipitate was filtered by suction, washed three times with water and dried at 50°C under vacuum to afford 3.67 g (91%) of the title compound.

$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 3.86 (s, 3H), 7.86 (d, 2H), 8.11 (d, 2H), 8.45 (s, 1H), 13.83 (s, 1H).

15

Intermediate 14

Methyl 2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate




Methyl 3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate (0.5 g, 1.68 mmol) was dissolved in DMF (26.6 mL). 1-Methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (698 mg, 3.53 mmol), 2,2'-bipyridine (655 mg, 4.19 mmol), cesium hydrogen carbonate (390 mg, 2.01 mmol), and anhydrous copper diacetate (380.7 mg, 2.10 mmol) were added. It was stirred for 21 h at rt. 1-Methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (349 mg, 1.68 mmol) was added and stirring was continued at rt overnight. Water (5 mL) was added and the pH was adjusted to 3 with 2N HCl (3.5 mL). The precipitate was filtered, washed three times with water, and dried at 50°C under vacuum to afford 594 mg (63%) of the title compound.

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.89 (s, 3H), 3.92 (s, 3H), 7.88 (br d, 2H), 8.11 (s, 1H), 8.28 (br d, 2H), 8.52 (s, 1H), 8.52 (s, 1H).

Intermediate 15

2-(1-Methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid




Methyl 2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate (590 mg, 1.56 mmol) was dissolved in acetonitrile (54 mL). A solution of lithium hydroxide (112 mg, 4.70 mmol) in water (1.7 mL) was added at rt. It was stirred for 3 h at rt. Water (100 mL) was added and the pH was adjusted to 6 with 2N HCl. The precipitate was filtered off under suction, washed with water and dried under vacuum at 50°C yielding 345 mg (45%) of the title compound which was used without further purification in the next step.

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.93 (br s, 3H), 7.47 - 8.39 (m, 7H)

25 Intermediate 16

Methyl 3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate




Methyl 3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate (2 g, 6.71 mmol) was dissolved in DMF (90 mL). 3-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (2.75 g, 13.41 mmol), 2,2'-bipyridine (2.62 g, 16.77 mmol), sodium carbonate (0.85 g, 8.02 mmol), and anhydrous copper diacetate (1.52 g, 8.37 mmol) were added. It was stirred for 3 h at 60°C. The reaction mixture was cooled down with an ice bath, water (240 mL) was added and the pH was adjusted to 3 with 2N HCl (20 mL). The precipitate was filtered, washed with water, and dried at 50°C under vacuum to afford 1.8 g (72%) of the title compound.

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.89 (s, 3H), 7.62 (dd, 1H), 7.88 (d, 2H), 8.15 - 8.21 (m, 3H), 8.60 (s, 1H), 8.68 (br d, 1H), 8.93 (br s, 1H).

Intermediate 17

3-Oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid




Methyl 3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate (1.80 g, 4.80 mmol) was dissolved in THF (28 mL). A solution of lithium hydroxide (345 mg, 23.95 mmol) in water (5 mL) was added at rt. It was stirred at rt overnight. Water (100 mL) was

added and the pH was adjusted to 6 with 2N HCl (4.5 mL). To the reaction mixture were added methylene chloride (50 mL) and chloroform (50 mL). The organic layer was separated and discarded. The precipitate was filtered off under suction, washed with water and dried under vacuum at 50°C affording 1036 mg (60%) of the title compound which was used without further purification in the next step.

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 7.60 (dd, 1H), 7.84 (d, 2H), 8.02 (s, 1H), 8.10 - 8.18 (m, 3H), 8.64 (d, 1H), 8.89 (d, 1H).

Intermediate 18

10 Diethyl hydroxy{2-oxo-2-[4-(trifluoromethyl)phenyl]ethyl}malonate




1-[4-(Trifluoromethyl)phenyl]ethanone (50 g, 0.266 mol) and diethyl oxomalonate (50.9 g, 0.292 mol) were stirred at 120 °C for 48 h. The reaction mixture was cooled to rt and the solid was filtered and washed with petrol ether (300 mL) affording 70 g (77%) of the title compound.

15 ¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.19 (t, 6H), 3.76 (s, 2H), 4.18 (q, 4H), 6.47 (s, 1H), 7.91 (d, 2H), 8.15 (d, 2H).

Intermediate 19

Ethyl 3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate



20


Diethyl hydroxy{2-oxo-2-[4-(trifluoromethyl)phenyl]ethyl}malonate (70 g, 0.193 mol) and hydrazine dihydrochloride (22.3 g, 0.212 mol) in ethanol (600 mL) were heated at 70 °C for 24

h. After completion, the reaction mixture was cooled to rt and concentrated under reduced pressure. The crude product was purified by column chromatography (silica gel, hexane / ethyl acetate 50%) to yield 35.0 g (58%) of the title compound.

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.31 (t, 3H), 4.32 (q, 2H), 7.86 (d, 2H), 8.11 (d, 2H), 8.42 (s, 1H), 13.81 (s, 1H).

Intermediate 20

Ethyl 3-oxo-2-(1,2-thiazol-4-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate



10

Ethyl 3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate (1.20 g, 3.84 mmol) was suspended in acetonitrile (24 mL). 4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2-thiazole (1.05 g, 5.00 mmol), pyridine (622 μL, 7.69 mmol), N,N-diethylethanamine (1.07 mL, 7.69 mmol), and anhydrous copper diacetate (907 mg, 5.00 mmol) were added. It was stirred for 28 h at rt. Water was added and the pH was adjusted to 3 with 2N HCl. The precipitate was filtered, washed three times with water, and dried at 50°C under vacuum to give 1.915 g of the title compound which was used without further purification in the next step.


15

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.34 (t, 3H), 4.37 (q, 2H), 7.90 (d, 2H), 8.27 (d, 2H), 8.56 (s, 1H), 9.15 (s, 1H), 9.60 (s, 1H).

20

Intermediate 21

3-Oxo-2-(1,2-thiazol-4-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid



To ethyl 3-oxo-2-(1,2-thiazol-4-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylate (1.91 g, 3.88 mmol) in acetonitrile (35 mL) was added lithium hydroxide (278 mg, 11.63 mmol) in water (4.2 mL). It was stirred for 2 h at rt. Water (5 mL) was added and the pH was adjusted to 3 with hydrochloric acid (3 mL, 2N). The precipitate was filtered, washed with water and dried under vacuum at 50 °C affording 1.3 g of the title compound and starting material.


The precipitate (465 mg) was stirred at 60 °C in aqueous sodium hydroxide solution. The solid material was filtered warm and washed with water. The residue was dried, suspended in water (20 mL) and the pH was adjusted to 3 with 2M hydrochloric acid. The solid material was collected, washed with water and dried under vacuum at 50 °C yielding 195 mg (11%) of the title compound. The first filtrate was acidified with 2M hydrochloric acid to pH 4, the precipitate was collected, washed with water and dried under vacuum at 50 °C to obtain 180 mg (10%) of the title compound.

The remaining impure material (720 mg) was stirred in aqueous sodium hydroxide solution for 1 h at rt. The pH was adjusted to 3 with hydrochloric acid (2 mL, 2M) and it was stirred for 0.5 h at rt. The solid was filtered, washed three times with water, dried under vacuum at 50 °C to give 660 mg (37%) of the title compound.

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 7.88 (br d, 2H), 8.28 (br d, 2H), 8.59 (br s, 1H), 9.16 (br s, 1H), 9.62 (br s, 1H), 13.94 (br s, 1H).

Intermediate 22


Diethyl hydroxy{2-oxo-2-[6-(trifluoromethyl)pyridin-3-yl]ethyl}malonate



Into a 100-mL round-bottom flask, was placed 1-[6-(trifluoromethyl)-3-pyridyl]ethanone (10 g, 52.87 mmol) and 1,3-diethyl 2-oxopropanedioate (15.65 g, 89.9 mmol). The resulting solution was stirred for 24 h at 130°C followed by the addition of more 1,3-diethyl 2-oxopropanedioate (13.81 g, 79.30 mmol) and heating for another 13 h at 130°C. The resulting mixture was cooled
 5 down to rt and poured into pentane. The precipitate was filtered off, washed with pentane and water yielding 22.6 g (crude) of diethyl 2-hydroxy-2-[2-oxo-2-[6-(trifluoromethyl)-3-pyridyl]ethyl]propanedioate which was used without further purification in the next step.

Intermediate 23

10 Ethyl 6-oxo-3-[6-(trifluoromethyl)-3-pyridyl]-1H-pyridazine-5-carboxylate




To a solution of diethyl 2-hydroxy-2-[2-oxo-2-[6-(trifluoromethyl)-3-pyridyl]ethyl]propanedioate (22.6 g, 62.2 mmol) in ethanol (255 mL) was added hydrazine hydrochloride (7.2 g, 68.5 mmol). The resulting solution was stirred for 24 h at 80°C. The reaction was then quenched by
 15 the addition of water. The resulting precipitate was filtered off and dried in vacuum to give 13.26 g (68%) of ethyl 6-oxo-3-[6-(trifluoromethyl)-3-pyridyl]-1H-pyridazine-5-carboxylate.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm = 1.32 (t, 3 H), 4.33 (d, 2 H), 8.03 (d, 1 H), 8.49 (s, 1 H), 8.55 (dd, 1 H), 9.25 (d, 1 H).

20 Intermediate 24

Ethyl 3-oxo-2-(1,2-thiazol-4-yl)-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxylate




Ethyl 3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxylate (475 mg, 1.52 mmol) was suspended in acetonitrile (10 mL). 4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2-thiazole (480 mg, 2.28 mmol), pyridine (0.245 mL, 3.03 mmol), N,N-diethylethanamine (0.423 mL, 3.03 mmol), and anhydrous copper diacetate (358 mg, 1.97 mmol) were added. It was stirred for 24 h at rt. 4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2-thiazole (100 mg, 0.474 mmol) was added and stirred for 24 h at rt. Buffer solution pH 7 (50 mL) was added and stirred for a short period. The precipitate was filtered, washed twice with water and dried under vacuum at 45 °C affording 630 mg of the title compound which was used without further purification in the next step.

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.34 (t, 3H), 4.38 (q, 2H), 8.07 (d, 1H), 8.63 (s, 1H), 8.73 (dd, 1H), 9.19 (s, 1H), 9.42 (d, 1H), 9.64 (s, 1H).

Intermediate 25

3-Oxo-2-(1,2-thiazol-4-yl)-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxylic acid



Ethyl 3-oxo-2-(1,2-thiazol-4-yl)-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxylate (625 mg, 1.58 mmol) was suspended in THF (19 mL). Lithium hydroxide (113 mg, 4.73 mmol) in water (2.3 mL) was added and stirred at rt for 24 h. Water (100 mL) was added and the pH was adjusted to 4 with hydrochloric acid (0.5N). It was stirred for a short period, the precipitate was filtered, washed three times with water and dried at 45 °C under vacuum to obtain 585 mg of the title compound which was used without further purification in the next step.


¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 7.99 (br s, 1H), 8.47 (br s, 1H), 9.14 (br s, 1H).

LC-MS (Instrument: Waters Acquity UPLC-MS SQD 3001; column: Acquity UPLC BEH C18 1.7 50x2.1mm; eluent A: water + 0.2 vol% aqueous ammonia (32%), eluent B: acetonitrile; gradient: 0-1.6 min 1-99% B, 1.6-2.0 min 99% B; flow 0.8 mL/min; temperature: 60°C;

injection: 2 μ L; DAD scan: 210-400 nm; ELSD): R_t = 0.59 min; MS (ESIpos): m/z = 369.1
[M+H]⁺

Intermediate 26


- 5 **Ethyl 2-(3-fluorophenyl)-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxylate**



- Diethyl 2-hydroxy-2-[2-oxo-2-[6-(trifluoromethyl)-3-pyridyl]ethyl]propanedioate (300 mg, 0.83 mmol) and (3-fluorophenyl)hydrazine hydrochloride (201 mg, 1.24 mmol) in ethanol (5 mL)
10 were stirred under reflux overnight. The reaction mixture was cooled down and concentrated on a rotavap. Water was added and it was extracted three times with dichloromethane. The combined organic phases were washed twice with water, dried over magnesium sulfate and concentrated. The residue was purified by HPLC affording 188 mg (56%) of the title compound.
- 15 ¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.33 (t, 3H), 4.36 (q, 2H), 7.34 - 7.40 (m, 1H), 7.56 - 7.67 (m, 3H), 8.05 (d, 1H), 8.59 - 8.63 (m, 2H), 9.32 (d, 1H).

Intermediate 27

- 20 **2-(3-Fluorophenyl)-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxylic acid**




Ethyl 2-(3-fluorophenyl)-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxylate (174 mg, 0.43 mmol) was suspended in THF (5 mL). Aqueous sodium hydroxide solution (2N, 0.641 mL, 1.28 mmol) was added and stirred at rt overnight. Water was added and the pH was adjusted to 3 with hydrochloric acid (2.0N). The precipitate was filtered, washed two times with water and dried at 50 °C under vacuum to give 150 mg (93%) of the title compound which was used without further purification in the next step.

¹H-NMR (400MHz, DMSO-*d*₆): δ [ppm] = 7.35 - 7.41 (m, 1H), 7.57 - 7.69 (m, 3H), 8.05 (d, 1H), 8.62 - 8.66 (m, 2H), 9.34 (d, 1H), 13.87 (br s, 1H).

10

Intermediate 28

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[2-[S-methyl-N-(trifluoroacetyl)sulfonimidoyl]ethyl]-3-oxo-2,3-dihydropyridazine-4-carboxamide




15 6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[2-(methylsulfinyl)ethyl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (80 mg, 0.18 mmol) was dissolved in dichloromethane (2 mL) followed by the addition of 2,2,2-trifluoroacetamide (104.2 mg, 0.92 mmol), diacetoxyiodo benzenel (223 mg, 0.69 mmol), rhodium(II) acetate dimer (5.2 mg, 0.01 mmol) and magnesium oxide (74 mg, 1.84 mmol). The reaction mixture was stirred for 48 h at rt, filtered and purified by HPLC (method D, basic) to yield 29 mg (29%) of the title compound.

20

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm = 3.63 (s, 3 H), 3.88 - 3.97 (m, 2 H), 3.97 - 4.03 (m, 2 H), 7.37 - 7.41 (m, 1 H), 7.54 - 7.65 (m, 5 H), 7.98 - 8.03 (m, 2 H), 8.65 (s, 1 H), 9.61 - 9.67 (m, 1 H).

Intermediate 29

6-(4-Chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-[S-methyl-N-(trifluoroacetyl)sulfonimidoyl]propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide




5

6-(4-Chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (33 mg, 0.077 mmol) was dissolved in dichloromethane (1 mL) followed by the addition of 2,2,2-trifluoroacetamide (45 mg, 0.38 mmol), diacetoxyiodo benzene (95 mg, 0.29 mmol), rhodium(II) acetate dimer (8.5 mg, 0.02 mmol) and magnesium oxide (31.1 mg, 0.77 mmol). The reaction mixture was stirred at rt for 48 h, then diluted with dichloromethane and washed with water. The organic phase was dried over sodium sulfate, filtered off and concentrated in vacuum. The residue was purified by HPLC (method D, basic) to yield 11 mg (26%) of the title compound.

15 ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm = 1.38 - 1.43 (m, 3 H), 3.61 (m, 3 H), 3.92 - 3.94 (m, 3 H), 3.94 - 4.01 (m, 1 H), 4.20 - 4.28 (m, 1 H), 4.70 - 4.80 (m, 1 H), 7.58 - 7.62 (m, 2 H), 8.10 (d, 2 H), 8.13 (d, 1 H), 8.52 (s, 1 H), 8.57 (d, 1 H), 9.64 - 9.69 (m, 1 H).

Intermediate 30

20 **N-[(cis)-4-Hydroxytetrahydrothiophen-3-yl]-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide**







2-(1-Methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid (100 mg, 0.28 mmol) was dissolved in anhydrous DMF (2.5 mL). Cis-4-Aminotetrahydrothiophene-3-ol hydrochloride (85 mg, 0.55 mmol), N-ethyl-N-isopropylpropan-2-amine (311 μ L, 1.78 mmol), and propane phosphonic acid anhydride (T3P, 240 μ L, 50% in DMF, 0.412 mmol) were successively added. It was stirred at rt for 4 h. The reaction mixture was concentrated under reduce pressure. The crude product was purified by HPLC yielding 38 mg (28%) of the title compound.



¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 2.71 - 2.79 (m, 2H), 3.04 (dd, 1H), 3.13 (dd, 1H), 3.94 (s, 3H), 4.31 - 4.41 (m, 2H), 5.68 (d, 1H), 7.89 (d, 2H), 8.13 (s, 1H), 8.30 (d, 2H), 8.57 (s, 1H), 8.68 (s, 1H), 9.86 (d, 1H).

The following intermediates were prepared from the starting materials stated in the table using the procedure described for intermediate 30. Enantiomers/diastereomers were separated from their racemate/diastereomeric mixture by chiral HPLC using the column and solvent conditions stated.

Intermediate	Structure	IUPAC name	Starting materials	Analytics

31		N-[(cis)-4-hydroxytetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide	3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol hydrochloride	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 2.71 (t, 2H), 3.03 (dd, 1H), 3.10 (dd, 1H), 4.30 - 4.39 (m, 2H), 5.67 (d, 1H), 7.64 (dd, 1H), 7.89 (d, 2H), 8.18 (ddd, 1H), 8.22 (d, 2H), 8.71 (dd, 1H), 8.76 (s, 1H), 8.92 (d, 1H), 9.76 (d, 1H).
32		(+) -N-[(cis)-4-hydroxytetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, enantiomer 1, [α] _D ²⁰ = +7.3° (c = 1.00, DMSO)	3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol hydrochloride, Chiralpak ID 5μ 250x30mm, eluent A: CO ₂ , eluent B: ethanol, isocratic: 30% B, 100 mL/min, 40 °C, BPR: 150 bar, 254 nm	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 2.69 - 2.76 (m, 2H), 3.03 (dd, 1H), 3.11 (dd, 1H), 4.30 - 4.39 (m, 2H), 5.67 (d, 1H), 7.64 (ddd, 1H), 7.89 (d, 2H), 8.18 (ddd, 1H), 8.22 (d, 2H), 8.71 (dd, 1H), 8.77 (s, 1H), 8.92 (dd, 1H), 9.76 (d, 1H). Rt = 2.22 min, Chiralpak ID 5μ 100x4.6mm, eluent A: CO ₂ , eluent B: ethanol, isocratic: 30% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 254 nm

33		<p>(-)-N-[(cis)-4-hydroxytetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, enantiomer 2, $[\alpha]_D^{20} = -2.0^\circ$ (c = 1.00, DMSO)</p>	<p>3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol hydrochloride, Chiralpak ID 5μ 250x30mm, eluent A: CO₂, eluent B: ethanol, isocratic: 30% B, 100 mL/min, 40 °C, BPR: 150 bar, 254 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 2.69 - 2.76 (m, 2H), 3.03 (dd, 1H), 3.11 (dd, 1H), 4.30 - 4.39 (m, 2H), 5.67 (d, 1H), 7.65 (ddd, 1H), 7.89 (d, 2H), 8.18 (ddd, 1H), 8.22 (d, 2H), 8.71 (dd, 1H), 8.77 (s, 1H), 8.92 (dd, 1H), 9.76 (d, 1H). Rt = 5.53 min, Chiralpak ID 5μ 100x4.6mm, eluent A: CO₂, eluent B: ethanol, isocratic: 30% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 254 nm</p>
34		<p>6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxytetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide</p>	<p>6-(4-chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol hydrochloride</p>	<p>¹H-NMR (600MHz, DMSO-d₆): δ [ppm] = 2.69 - 2.74 (m, 2H), 3.03 (dd, 1H), 3.10 (dd, 1H), 4.30 - 4.36 (m, 2H), 5.66 (br d, 1H), 7.39 (td, 1H), 7.55 (d, 1H), 7.58 (d, 2H), 7.60 - 7.65 (m, 2H), 8.00 (d, 2H), 8.68 (s, 1H), 9.82 (d, 1H).</p>


35		<p>(+)-6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxytetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, enantiomer 1, $[\alpha]_D^{20} = +5.9^\circ$ (c = 1.00, methanol)</p>	<p>6-(4-chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol hydrochloride, Chiralpak IA 5μ 250x30mm, eluent A: CO₂, eluent B: ethanol, isocratic: 45% B, 100 mL/min, 40 °C, BPR: 150 bar, 254 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 2.68 - 2.75 (m, 2H), 3.03 (dd, 1H), 3.10 (dd, 1H), 4.29 - 4.37 (m, 2H), 5.65 (d, 1H), 7.36 - 7.42 (m, 1H), 7.53 - 7.65 (m, 5H), 7.98 - 8.02 (m, 2H), 8.68 (s, 1H), 9.81 (d, 1H). Rt = 2.16 min, Chiralpak IA 5μ 100x4.6mm, eluent A: CO₂, eluent B: ethanol, isocratic: 45% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 254 nm</p>
36		<p>(-)-6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxytetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, enantiomer 2, $[\alpha]_D^{20} = -6.6^\circ$ (c = 1.00, methanol)</p>	<p>6-(4-chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol hydrochloride, Chiralpak IA 5μ 250x30mm, eluent A: CO₂, eluent B: ethanol, isocratic: 45% B, 100 mL/min, 40 °C, BPR: 150 bar, 254 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 2.68 - 2.75 (m, 2H), 3.03 (m, 1H), 3.10 (dd, 1H), 4.29 - 4.37 (m, 2H), 5.65 (d, 1H), 7.36 - 7.42 (m, 1H), 7.53 - 7.65 (m, 5H), 7.98 - 8.02 (m, 2H), 8.68 (s, 1H), 9.81 (d, 1H). Rt = 3.84 min, Chiralpak IA 5μ 100x4.6mm, eluent A: CO₂, eluent B: ethanol, isocratic: 45% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 254 nm</p>

EXPERIMENTAL SECTION – EXAMPLES

The following examples describe the embodiment of the instant invention, not restricting the invention to these examples only.

Example 1

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfanyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide




6-(4-Chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid (700 mg, 2.03 mmol) was dissolved in anhydrous DMF (21 mL). (2S)-1-(Methylsulfonyl)propan-2-amine (427 mg, 4.06 mmol), N-ethyl-N-isopropylpropan-2-amine (1.59 mL, 9.14 mmol), and propane phosphonic acid anhydride (T3P, 1.78 mL, 50% in DMF, 3.05 mmol) were successively added. It was stirred at rt for 4 h. The reaction mixture was concentrated under reduce pressure and combined with a batch (100 mg acid) which was synthesized in analogous manner. The crude product was purified by flash chromatography (silica gel, hexane / ethyl acetate, gradient) yielding 775 mg (77%) of the title compound.

10

$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 1.26 (d, 3H), 2.10 (s, 3H), 2.65 - 2.76 (m, 2H), 4.17 - 4.28 (m, 1H), 7.36 - 7.41 (m, 1H), 7.53 - 7.66 (m, 5H), 7.97 - 8.02 (m, 2H), 8.65 (s, 1H), 9.42 (d, 1H).

15 **Example 2**

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfonyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide



6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfonyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (50 mg, 116 μmol) was dissolved in chloroform (434 μL). Between 0-10 $^\circ\text{C}$ 3-chlorobenzenecarboxylic acid (60 mg, 348 μmol) was added portionwise. It was stirred at rt for 1.5 h. This batch and a second batch prepared from 20 mg sulfide in analogous manner were combined and diluted with dichloromethane (30 mL). The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution (10


mL), water (10 mL), dried over sodium sulfate and concentrated to dryness. The residue was purified by RP-HPLC (column: X-Bridge C18 5 μ m 100x30 mm, mobile phase: (water + 0.2 vol% aqueous ammonia (32%)) / acetonitrile, gradient and X-Bridge C18 5 μ m 100x30 mm, mobile phase: (water + 0.1 vol% formic acid (99%)) / methanol, gradient) to afford 21.5 mg (29%) of the title compound.

$^1\text{H-NMR}$ (400MHz, DMSO- d_6): δ [ppm] = 1.34 (d, 3H), 3.02 (s, 3H), 3.38 (dd, 1H), 3.60 (dd, 1H), 4.53 - 4.63 (m, 1H), 7.36 - 7.41 (m, 1H), 7.53 - 7.66 (m, 5H), 7.98 - 8.03 (m, 2H), 8.64 (s, 1H), 9.52 (d, 1H).

$[\alpha]_D^{20} = +51.3^\circ$ (c = 1.00, methanol).

Example 3

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide




6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (150 mg, 347 μ mol) was dissolved in a mixture of acetone (796 μ L), methanol (464 μ L), and water (164 μ L). Sodium periodate (74.3 mg, 347 μ mol) was added and stirred overnight at rt. The reaction mixture was concentrated under reduced pressure and combined with a second batch (20 mg of the sulfide) which was prepared under analogous conditions. The residue was dissolved in dichloromethane (30 mL), washed twice with water (50 mL) dried over magnesium sulfate and concentrated. The residue was purified by RP-HPLC (column: X-Bridge C18 5 μ m 100x30 mm, mobile phase: (water + 0.2 vol% aqueous ammonia (32%)) / acetonitrile, gradient) to give 80 mg (45%) of the title compound as diastereomeric mixture of ca. 1:1.

$^1\text{H-NMR}$ (400MHz, DMSO- d_6): δ [ppm] = 1.32 - 1.35 (m, 6H), 2.57 (s, 3H), 2.61 (s, 3H), 2.88 - 3.17 (m, 4H), 4.37 - 4.54 (m, 2H), 7.35 - 7.42 (m, 2H), 7.53 - 7.66 (m, 10H), 8.00 (d, 4H), 8.64 (s, 1H), 8.65 (s, 1H), 9.48 - 9.55 (m, 2H).

Example 4

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, diastereomer 1



- 5 6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (77 mg) was separated by chiral HPLC (column: Chiralpak IA 5 μ m 250x30 mm, mobile phase: eluent A: CO₂, eluent B: isopropanol, isocratic: 45% B, 100 mL/min, 40 °C, BPR: 150 bar, 254 nm) to yield 18 mg of the title compound.

- 10 ¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.34 (d, 3H), 2.61 (s, 3H), 2.98 - 3.08 (m, 2H), 4.37 - 4.48 (m, 1H), 7.36 - 7.41 (m, 1H), 7.54 - 7.66 (m, 5H), 7.98 - 8.03 (m, 2H), 8.64 (s, 1H), 9.50 (d, 1H).


Chiral HPLC: Rt = 1.28 min

- 15 Chiralpak IA 5 μ m 100x4.6 mm; eluent A: CO₂, eluent B: isopropanol, isocratic: 45% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 254 nm.

$[\alpha]_D^{20} = +17.7^\circ$ (c = 1.00, methanol).

20 **Example 5**

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, diastereomer 2



6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (77 mg) was separated by chiral HPLC (column: Chiralpak IA 5 μ m 250x30 mm, mobile phase: eluent A: CO₂, eluent B: isopropanol, isocratic: 45% B, 100 mL/min, 40 °C, BPR: 150 bar, 254 nm) to yield 19 mg of the title compound.

5

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.33 (d, 3H), 2.57 (s, 3H), 2.91 (dd, 1H), 3.13 (dd, 1H), 4.42 - 4.54 (m, 1H), 7.35 - 7.42 (m, 1H), 7.53 - 7.65 (m, 5H), 7.97 - 8.03 (m, 2H), 8.65 (s, 1H), 9.52 (d, 1H).

10 Chiral HPLC: Rt = 2.70 min


Chiralpak IA 5 μ m 100x4.6 mm; eluent A: CO₂, eluent B: isopropanol, isocratic: 45% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 254 nm.

$[\alpha]_D^{20} = +95.6^\circ$ (c = 1.00, methanol).

15

Example 6

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(S-methylsulfonyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide



20 **Step 1:** 6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (290 mg, 647 μ mol) was dissolved in dichloromethane (7 mL). 2,2,2-Trifluoroacetamide (365 mg, 3.24 mmol), (diacetoxyiodo)benzene (782 mg, 2.43 mmol), rhodium(II) acetate dimer (71.5 mg, 162 μ mol), and magnesium oxide (261 mg, 6.47 mmol) were added. It was stirred at rt overnight. The reaction mixture was diluted with
25 dichloromethane (40 mL), washed with water (15 mL) dried over magnesium sulfate and concentrated.


Step 2: The residue of step 1 was dissolved in methanol (12.7 mL) and potassium carbonate (161.5 mg, 1.17 mmol) was added. It was stirred overnight at rt. The reaction mixture was diluted with dichloromethane (25 mL), washed with water (10 mL) dried over magnesium
30 sulfate and concentrated. The residue was purified by RP-HPLC (column: X-Bridge C18 5 μ m

100x30 mm, mobile phase: (water + 0.2 vol% aqueous ammonia (32%)) / acetonitrile, gradient) to give 107 mg (36%) of the title compound as diastereomeric mixture of ca. 1:1.

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.34 (d, 6H), 2.94 (s, 6H), 3.21 - 3.31 (m, 2H), 3.49 - 3.57 (m, 2H), 3.75 (d, 2H), 4.49 - 4.61 (m, 2H), 7.35 - 7.42 (m, 2H), 7.53 - 7.66 (m, 10H), 7.97 - 8.03 (m, 4H), 8.63 (s, 1H), 8.63 (s, 1H), 9.47 (d, 1H), 9.52 (d, 1H).

Example 7

10 **6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(S-methylsulfonimidoyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, diastereomer 1**



15 6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(S-methylsulfonimidoyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (100 mg) was separated by chiral HPLC (column: Chiralpak IB 5 μm 250x30 mm, mobile phase: eluent A: CO₂, eluent B: ethanol, isocratic: 32% B, 100 mL/min, 40 °C, BPR: 150 bar, 254 nm) to give 45 mg of the title compound.

¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.34 (d, 3H), 2.94 (s, 3H), 3.24 - 3.31 (m, 1H), 3.53 (dd, 1H), 3.73 (s, 1H), 4.49 - 4.61 (m, 1H), 7.35 - 7.41 (m, 1H), 7.53 - 7.66 (m, 5H), 8.00 (d, 2H), 8.63 (s, 1H), 9.52 (d, 1H).

20


Chiral HPLC: Rt = 1.73 min

Chiralpak IB 5 μm 100x4.6 mm; eluent A: CO₂, eluent B: ethanol, isocratic: 32% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 254 nm.

25 $[\alpha]_D^{20} = +64.1^\circ$ (c = 1.00, DMSO).

Example 8

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(S-methylsulfonimidoyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, diastereomer 2



6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(S-methylsulfonimidoyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (100 mg) was separated by chiral HPLC (column: Chiralpak IB 5 μm 250x30 mm, mobile phase: eluent A: CO_2 , eluent B: ethanol, isocratic: 32% B, 100 mL/min, 40 $^\circ\text{C}$, BPR: 150 bar, 254 nm) to give 40 mg of the title compound.

$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 1.34 (d, 3H), 2.93 (s, 3H), 3.24 (dd, 1H), 3.52 (dd, 1H), 3.77 (s, 1H), 4.49 - 4.61 (m, 1H), 7.35 - 7.42 (m, 1H), 7.53 - 7.65 (m, 5H), 7.98 - 8.02 (m, 2H), 8.63 (s, 1H), 9.47 (d, 1H).

10


Chiral HPLC: R_t = 2.21 min

Chiralpak IB 5 μm 100x4.6 mm; eluent A: CO_2 , eluent B: ethanol, isocratic: 32% B, 4 mL/min, 37.5 $^\circ\text{C}$, BPR: 100 bar, 254 nm.

15 $[\alpha]_D^{20}$ = +58.8 $^\circ$ (c = 1.00, DMSO).

Example 9

2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfanyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide



20

2-(1-Methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid (800 mg, 2.20 mmol) was dissolved in anhydrous DMF (23 mL). (2S)-1-(Methylsulfanyl)propan-2-amine (462 mg, 4.39 mmol), N-ethyl-N-isopropylpropan-2-amine (1.72 mL, 9.88 mmol), and propane phosphonic acid anhydride (T3P, 1.92 mL, 50% in DMF,

3.29 mmol) were successively added. It was stirred at rt for 4 h. The reaction mixture was concentrated under reduce pressure. The crude product was purified by flash chromatography (silica gel, hexane / ethyl acetate, gradient) affording 875 mg (88%) of the title compound.

- 5 ¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.29 (d, 3H), 2.12 (s, 3H), 2.68 - 2.79 (m, 2H), 3.94 (s, 3H), 4.19 - 4.30 (m, 1H), 7.89 (d, 2H), 8.13 (d, 1H), 8.30 (d, 2H), 8.58 (s, 1H), 8.65 (s, 1H), 9.50 (d, 1H).



[α]_D²⁰ = +49.9° (c = 1.00, DMSO).



10



The following examples were prepared from the starting materials stated in the table using the procedure described in this example. Enantiomers/diastereomers were separated from their racemate/diastereomeric mixture by chiral HPLC using the column and solvent conditions stated.


15 **Table 2: Examples 10 – 21**



Expl	Structure	IUPAC name	Starting materials	Analytics
10		N-[(cis)-4-hydroxy-1,1-dioxido-2,3-dihydrothiophene-3-yl]-3-oxo-2-(1,2-thiazol-4-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide	3-oxo-2-(1,2-thiazol-4-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol 1,1-dioxide hydrochloride	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 3.17 (t, 1H), 3.27 - 3.32 (m, 1H), 3.51 - 3.59 (m, 2H), 4.53 - 4.59 (m, 1H), 4.73 - 4.83 (m, 1H), 6.38 (d, 1H), 7.90 (d, 2H), 8.30 (d, 2H), 8.75 (s, 1H), 9.14 (s, 1H), 9.62 (s, 1H), 9.91 (d, 1H).



11		<p>(-)-N-[(cis)-4-hydroxy-1,1-dioxido-2,3-dihydro-1,2-thiazol-4-yl]-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, enantiomer 1, $[\alpha]_D^{20} = -6.6^\circ$ (c = 1.00, DMSO)</p>	<p>3-oxo-2-(1,2-thiazol-4-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol 1,1-dioxide hydrochloride, YMC Cellulose SB 5μ 250x30mm, eluent A: methanol + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 50% B, 40 mL/min, 254 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.17 (t, 1H), 3.26 - 3.32 (m, 1H and water signal), 3.51 - 3.58 (m, 2H), 4.56 (br s, 1H), 4.75 - 4.82 (m, 1H), 6.38 (s, 1H), 7.90 (d, 2H), 8.30 (d, 2H), 8.75 (s, 1H), 9.14 (s, 1H), 9.62 (s, 1H), 9.91 (d, 1H). Rt = 3.36 min, YMC Cellulose SB 3μm 100x4.6mm, eluent A: methanol + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 50% B, 1.4 mL/min, 25 °C, 254 nm</p>
12		<p>(+)-N-[(cis)-4-hydroxy-1,1-dioxido-2,3-dihydro-1,2-thiazol-4-yl]-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, enantiomer 2, $[\alpha]_D^{20} = +7.6^\circ$ (c = 1.00, DMSO)</p>	<p>3-oxo-2-(1,2-thiazol-4-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol 1,1-dioxide hydrochloride, YMC Cellulose SB 5μ 250x30mm, eluent A: methanol + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 50% B, 40 mL/min, 254 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.17 (t, 1H), 3.27 - 3.32 (m, 1H and water signal), 3.51 - 3.58 (m, 2H), 4.56 (br s, 1H), 4.74 - 4.83 (m, 1H), 6.38 (d, 1H), 7.90 (d, 2H), 8.30 (d, 2H), 8.75 (s, 1H), 9.14 (s, 1H), 9.62 (s, 1H), 9.91 (d, 1H). Rt = 4.15 min, YMC Cellulose SB 3μm 100x4.6mm, eluent A: methanol + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 50% B, 1.4 mL/min, 25 °C, 254 nm</p>

13		2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide	2-(3-fluorophenyl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol 1,1-dioxide	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 3.10 - 3.18 (m, 1H), 3.28 (d, 1H), 3.50 - 3.56 (m, 2H), 4.51 - 4.56 (m, 1H), 4.73 - 4.80 (m, 1H), 6.34 (d, 1H), 7.38 - 7.44 (m, 1H), 7.55 - 7.58 (m, 1H), 7.61 - 7.67 (m, 2H), 7.89 (d, 2H), 8.21 (d, 2H), 8.76 (s, 1H), 9.95 (d, 1H).
14		(+)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, enantiomer 1, [α] _D ²⁰ = +3.8° (c = 1.00, methanol)	2-(3-fluorophenyl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol 1,1-dioxide, Chiralpak IA 5μ 250x30mm, eluent A: methanol, eluent B: ethanol, isocratic: 50% B, 40 mL/min, 254 nm	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 3.10 - 3.18 (m, 1H), 3.25 - 3.30 (m, 1H), 3.50 - 3.56 (m, 2H), 4.54 (br s, 1H), 4.73 - 4.80 (m, 1H), 6.35 (br d, 1H), 7.38 - 7.44 (m, 1H), 7.55 - 7.58 (m, 1H), 7.60 - 7.67 (m, 2H), 7.89 (d, 2H), 8.21 (d, 2H), 8.76 (s, 1H), 9.95 (d, 1H). Rt = 3.83 min, Chiralpak IA 3μ 100x4.6mm, eluent A: methanol + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 50% B, 1.4 mL/min, 254 nm

15		<p>(-)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, enantiomer 2, $[\alpha]_D^{20} = -3.6^\circ$ (c = 1.00, methanol)</p>	<p>2-(3-fluorophenyl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol 1,1-dioxide, Chiralpak IA 5μ 250x30mm, eluent A: methanol, eluent B: ethanol, isocratic: 50% B, 40 mL/min, 254 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.10 - 3.18 (m, 1H), 3.28 (d, 1H), 3.50 - 3.56 (m, 2H), 4.54 (br s, 1H), 4.73 - 4.80 (m, 1H), 6.34 (br d, 1H), 7.38 - 7.44 (m, 1H), 7.54 - 7.58 (m, 1H), 7.61 - 7.67 (m, 2H), 7.89 (d, 2H), 8.21 (d, 2H), 8.76 (s, 1H), 9.95 (d, 1H). Rt = 5.19 min, Chiralpak IA 3μ 100x4.6mm, eluent A: methanol + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 50% B, 1.4 mL/min, 254 nm</p>
16		<p>N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide</p>	<p>3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol 1,1-dioxide hydrochloride (1:1)</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.14 (t, 1H), 3.28 (d, 1H), 3.50 - 3.56 (m, 2H), 4.54 (br d, 1H), 4.77 (dtd, 1H), 6.34 (d, 1H), 7.65 (dd, 1H), 7.89 (d, 2H), 8.18 (ddd, 1H), 8.23 (d, 2H), 8.71 (br d, 1H), 8.78 (s, 1H), 8.92 (br d, 1H), 9.92 (d, 1H).</p>


17		<p>(+)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, enantiomer 1, $[\alpha]_D^{20} = +7.1^\circ$ (c = 1.00, DMSO)</p>	<p>N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, Chiralpak ID 5μ 250x30mm, eluent A: CO₂, eluent B: ethanol + 0.2vol% aqueous ammonia (32%), isocratic: 34% B, 100 mL/min, 40 °C, BPR: 150 bar, 220 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.11 - 3.18 (m, 1H), 3.25 - 3.30 (m, 1H), 3.50 - 3.57 (m, 2H), 4.54 (br d, 1H), 4.77 (dtd, 1H), 6.34 (d, 1H), 7.63 - 7.67 (m, 1H), 7.89 (d, 2H), 8.18 (ddd, 1H), 8.23 (d, 2H), 8.71 (dd, 1H), 8.78 (s, 1H), 8.92 (d, 1H), 9.92 (d, 1H). Rt = 1.33 min, Chiralpak ID 5μ 100x4.6mm, eluent A: CO₂, eluent B: ethanol + 0.2vol% aqueous ammonia (32%), isocratic: 39% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 220 nm</p>
----	---	--	---	--

18		<p>(-)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, enantiomer 2, $[\alpha]_D^{20} = -3.7^\circ$ (c = 1.00, DMSO)</p>	<p>N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, Chiralpak ID 5μ 250x30mm, eluent A: CO₂, eluent B: ethanol + 0.2vol% aqueous ammonia (32%), isocratic: 34% B, 100 mL/min, 40 °C, BPR: 150 bar, 220 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.11 - 3.18 (m, 1H), 3.25 - 3.30 (m, 1H), 3.50 - 3.56 (m, 2H), 4.54 (br d, 1H), 4.77 (dtd, 1H), 6.34 (d, 1H), 7.65 (ddd, 1H), 7.89 (d, 2H), 8.18 (ddd, 1H), 8.23 (d, 2H), 8.71 (dd, 1H), 8.78 (s, 1H), 8.91 - 8.93 (m, 1H), 9.92 (d, 1H). Rt = 2.27 min, Chiralpak ID 5μ 100x4.6mm, eluent A: CO₂, eluent B: ethanol + 0.2vol% aqueous ammonia (32%), isocratic: 39% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 220 nm</p>
19		<p>2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxamide</p>	<p>2-(3-fluorophenyl)-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxylic acid, cis-4-aminotetrahydrothiophene-3-ol 1,1-dioxide hydrochloride (1:1)</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 3.11 - 3.18 (m, 1H), 3.26 - 3.30 (m, 1H), 3.50 - 3.56 (m, 2H), 4.54 (br d, 1H), 4.77 (dtd, 1H), 6.34 (d, 1H), 7.38 - 7.44 (m, 1H), 7.56 - 7.69 (m, 3H), 8.05 (d, 1H), 8.67 (dd, 1H), 8.85 (s, 1H), 9.36 (d, 1H), 9.92 (d, 1H).</p>

20		<p>(+)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxamide, enantiomer 1, $[\alpha]_D^{20} = +3.7^\circ$ (c = 1.00, DMSO)</p>	<p>2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxamide, Chiralpak IA 5μ 250x30mm, eluent A: methanol, eluent B: ethanol, isocratic: 50% B, 40 mL/min, 254 nm</p>	<p>$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 3.11 - 3.18 (m, 1H), 3.25 - 3.30 (m, 1H), 3.50 - 3.57 (m, 2H), 4.54 (br d, 1H), 4.76 (dtd, 1H), 6.34 (d, 1H), 7.38 - 7.44 (m, 1H), 7.57 - 7.69 (m, 3H), 8.05 (d, 1H), 8.67 (dd, 1H), 8.85 (s, 1H), 9.36 (d, 1H), 9.92 (d, 1H). Rt = 3.55 min, Chiralpak IA 3μ 100x4.6mm, eluent A: methanol + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 50% B, 1.4 mL/min, 25 °C, 254 nm</p>
21		<p>(-)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxamide, enantiomer 2, $[\alpha]_D^{20} = -0.7^\circ$ (c = 1.00, DMSO)</p>	<p>2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxamide, Chiralpak IA 5μ 250x30mm, eluent A: methanol, eluent B: ethanol, isocratic: 50% B, 40 mL/min, 254 nm</p>	<p>$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 3.11 - 3.18 (m, 1H), 3.26 - 3.30 (m, 1H), 3.50 - 3.57 (m, 2H), 4.54 (br d, 1H), 4.77 (dtd, 1H), 6.34 (d, 1H), 7.39 - 7.44 (m, 1H), 7.57 - 7.68 (m, 3H), 8.05 (d, 1H), 8.67 (dd, 1H), 8.85 (s, 1H), 9.36 (d, 1H), 9.92 (d, 1H). Rt = 5.41 min, Chiralpak IA 3μ 100x4.6mm, eluent A: methanol + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 50% B, 1.4 mL/min, 25 °C, 254 nm</p>

Example 22

2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfonyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide




2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfonyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (100 mg, 221 μmol) was dissolved in chloroform (830 μL). Between 0-10 $^{\circ}\text{C}$ 3-chlorobenzenecarboxylic acid (114 mg, 661 μmol) was added portionwise. It was stirred at rt for 3 h. The reaction mixture was diluted with dichloromethane (30 mL). The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution (10 mL), water (10 mL), dried over magnesium sulfate and concentrated to dryness. The residue was purified by RP-HPLC (column: X-Bridge C18 5 μm 100x30 mm, mobile phase: (water + 0.2 vol% aqueous ammonia (32%)) / acetonitrile, gradient) to obtain 73 mg (68%) of the title compound.

$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 1.37 (d, 3H), 3.04 (s, 3H), 3.42 (dd, 1H), 3.61 (dd, 1H), 3.94 (s, 3H), 4.54 - 4.65 (m, 1H), 7.89 (d, 2H), 8.14 (s, 1H), 8.30 (d, 2H), 8.55 (s, 1H), 8.63 (s, 1H), 9.58 (d, 1H).

$[\alpha]_{\text{D}}^{20} = +69.7^{\circ}$ ($c = 1.00$, DMSO).

20 Example 23

2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide




2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfanyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (270 mg, 598 μ mol) was dissolved in a mixture of acetone (2.75 mL), methanol (1.2 mL), and water (283 μ L). Sodium periodate (128 mg, 598 μ mol) was added and stirred at rt for 72 h. Sodium periodate (25.6 mg, 120 μ mol) was added and stirred at rt for 24 h. The reaction mixture was filtered and the solid was washed with acetone. The combined filtrate were concentrated and dissolved in dichloromethane (30 mL), washed twice with water (50 mL) dried over magnesium sulfate and concentrated. The residue was purified by RP-HPLC (column: X-Bridge C18 5 μ m 100x30 mm, mobile phase: (water + 0.2 vol% aqueous ammonia (32%)) / acetonitrile, gradient) yielding 206 mg (74%) of the title compound as diastereomeric mixture. of ca. 1:1.

1 H-NMR (400MHz, DMSO- d_6): δ [ppm] = 1.36 - 1.38 (m, 6H), 2.59 (s, 3H), 2.63 (s, 3H), 2.95 (dd, 1H), 3.00 - 3.11 (m, 2H), 3.16 (dd, 1H), 3.94 (s, 6H), 4.38 - 4.56 (m, 2H), 7.89 (d, 4H), 8.12 - 8.15 (m, 2H), 8.30 (d, 4H), 8.54 (s, 1H), 8.57 (s, 1H), 8.63 (s, 1H), 8.64 (s, 1H), 9.54 - 9.59 (m, 2H).

Example 24

2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, diastereomer 1



2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (203 mg) was separated by chiral HPLC (column: Chiralpak ID 5 μ m 250x30 mm, mobile phase: eluent A: acetonitrile and 0.1 vol% of diethylamine (99%), eluent B: ethanol, isocratic: 15% B, 60 mL/min, 254 nm) to yield 95 mg of the title compound.

1 H-NMR (400MHz, DMSO- d_6): δ [ppm] = 1.37 (d, 3H), 2.63 (s, 3H), 2.99 - 3.11 (m, 2H), 3.93 (s, 3H), 4.38 - 4.50 (m, 1H), 7.89 (d, 2H), 8.14 (s, 1H), 8.30 (d, 2H), 8.57 (s, 1H), 8.63 (s, 1H), 9.57 (d, 1H).

30

Chiral HPLC: Rt = 2.95 min


Chiralpak ID 3 μm 100x4.6 mm; eluent A: acetonitrile and 0.1vol% of diethylamine (99%),
eluent B: ethanol, isocratic: 15% B, 1.4 mL/min, 25 °C, 254 nm.

$[\alpha]_{\text{D}}^{20} = +28.7^\circ$ (c = 1.00, DMSO).

5

Example 25

2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, diastereomer 2



10 2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (203 mg) was separated by chiral HPLC (column: Chiralpak ID 5 μm 250x30 mm, mobile phase: eluent A: acetonitrile and 0.1vol% of diethylamine (99%), eluent B: ethanol, isocratic: 15% B, 60 mL/min, 254 nm) to yield 83 mg of the title compound.

15

$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 1.36 (d, 3H), 2.59 (s, 3H), 2.95 (dd, 1H), 3.16 (dd, 1H), 3.93 (s, 3H), 4.44 - 4.56 (m, 1H), 7.89 (d, 2H), 8.13 (s, 1H), 8.30 (d, 2H), 8.54 (s, 1H), 8.64 (s, 1H), 9.56 (d, 1H).

20 Chiral HPLC: $R_t = 4.80$ min


Chiralpak ID 3 μm 100x4.6 mm; eluent A: acetonitrile and 0.1vol% of diethylamine (99%),
eluent B: ethanol, isocratic: 15% B, 1.4 mL/min, 25 °C, 254 nm.



$[\alpha]_{\text{D}}^{20} = +104.1^\circ$ (c = 1.00, DMSO).



25


The following examples were prepared from the starting materials stated in the table using the procedure described in the examples above. Enantiomers/diastereomers were separated from their racemate/diastereomeric mixture by chiral HPLC using the column and solvent conditions stated.


30 **Table 3: Examples 26 – 37**


Expl	Structure	IUPAC name	Starting materials	Analytics
26		6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 1)	(+)-6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxytetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide enantiomer 1	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 2.70 - 2.80 (m, 2H, minor), 2.83 - 2.94 (m, 2H, major), 3.07 - 3.13 (m, 1H, major), 3.59 (dd, 1H, major), 3.81 (dd, 1H, minor), 4.40 - 4.43 (m, 1H, minor), 4.46 - 4.54 (m, 1H, minor), 4.57 - 4.60 (m, 1H, major), 4.97 - 5.07 (m, 1H, major), 5.84 (d, 1H, minor), 5.88 (d, 1H, major), 7.35 - 7.43 (m, 1H), 7.52 - 7.66 (m, 5H), 7.98 - 8.04 (m, 2H), 8.68 (s, 1H, minor), 8.70 (s, 1H, major), 9.87 (d, 1H, major), 9.99 (d, 1H, minor).


27		<p>(+)-6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, diastereomer 1, $[\alpha]_D^{20} = +27.7^\circ$ (c = 1.00, DMSO)</p>	<p>6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 1), Chiralpak IC 5μ 250x30mm, eluent A: acetonitrile + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 10% B, 50 mL/min, 254 nm</p>	<p>$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 2.70 - 2.80 (m, 2H), 3.35 - 3.38 (m, 1H and water signal), 3.81 (dd, 1H), 4.39 - 4.44 (m, 1H), 4.46 - 4.55 (m, 1H), 5.84 (d, 1H), 7.37 - 7.42 (m, 1H), 7.54 - 7.65 (m, 5H), 7.98 - 8.03 (m, 2H), 8.68 (s, 1H), 9.99 (d, 1H). Rt = 3.37 min, Chiralpak IC 3μ 100x4.6mm, eluent A: acetonitrile + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 10% B, 1.4 mL/min, 25 $^\circ\text{C}$, 254 nm</p>
28		<p>(-)-6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, diastereomer 2, $[\alpha]_D^{20} = -13.3^\circ$ (c = 1.00, DMSO)</p>	<p>6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 1), Chiralpak IC 5μ 250x30mm, eluent A: acetonitrile + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 10% B, 50 mL/min, 254 nm</p>	<p>$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 2.83 - 2.94 (m, 2H), 3.10 (dd, 1H), 3.59 (dd, 1H), 4.55 - 4.61 (m, 1H), 4.97 - 5.06 (m, 1H), 5.89 (d, 1H), 7.36 - 7.42 (m, 1H), 7.53 - 7.65 (m, 5H), 7.99 - 8.04 (m, 2H), 8.71 (s, 1H), 9.87 (d, 1H). Rt = 5.23 min, Chiralpak IC 3μ 100x4.6mm, eluent A: acetonitrile + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 10% B, 1.4 mL/min, 25 $^\circ\text{C}$, 254 nm</p>


29		6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 2)	(-)-6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxytetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, enantiomer 2	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 2.70 - 2.80 (m, 2H, minor), 2.83 - 2.94 (m, 2H, major), 3.10 (dd, 1H, major), 3.59 (dd, 1H, major), 3.81 (dd, 1H, minor), 4.38 - 4.43 (m, 1H, minor), 4.46 - 4.55 (m, 1H, minor), 4.56 - 4.61 (m, 1H, major), 4.97 - 5.06 (m, 1H, major), 5.84 (d, 1H, minor), 5.88 (d, 1H, major), 7.36 - 7.42 (m, 1H), 7.53 - 7.66 (m, 5H), 7.98 - 8.04 (m, 2H), 8.68 (s, 1H, minor), 8.71 (s, 1H, major), 9.87 (d, 1H, major), 9.99 (d, 1H, minor).
30		(-)-6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, diastereomer 1, $[\alpha]_D^{20} = -21.5^\circ$ (c = 1.00, DMSO)	6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 2), Chiralpak IC 5 μ 250x30mm, eluent A: acetonitrile + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 10% B, 50 mL/min, 254 nm	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 2.70 - 2.80 (m, 2H), 3.34 - 3.38 (m, 1H and water signal), 3.81 (dd, 1H), 4.39 - 4.44 (m, 1H), 4.46 - 4.55 (m, 1H), 5.84 (d, 1H), 7.37 - 7.42 (m, 1H), 7.54 - 7.66 (m, 5H), 7.98 - 8.03 (m, 2H), 8.68 (s, 1H), 9.99 (d, 1H). Rt = 2.12 min, Chiralpak IC 3 μ 100x4.6mm, eluent A: acetonitrile + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 10% B, 1.4 mL/min, 25 °C, 254 nm


31		<p>(+)-6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, diastereomer 2, $[\alpha]_D^{20} = +14.2^\circ$ (c = 1.00, DMSO)</p>	<p>6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 2), Chiralpak IC 5μ 250x30mm, eluent A: acetonitrile + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 10% B, 50 mL/min, 254 nm</p>	<p>$^1\text{H-NMR}$ (400MHz, DMSO-d_6): δ [ppm] = 2.83 - 2.94 (m, 2H), 3.10 (dd, 1H), 3.59 (dd, 1H), 4.55 - 4.61 (m, 1H), 4.97 - 5.06 (m, 1H), 5.89 (d, 1H), 7.36 - 7.42 (m, 1H), 7.53 - 7.65 (m, 5H), 7.99 - 8.04 (m, 2H), 8.70 (s, 1H), 9.87 (d, 1H). Rt = 4.56 min, Chiralpak IC 3μ 100x4.6mm, eluent A: acetonitrile + 0.1vol% diethylamine (99%), eluent B: ethanol, isocratic: 10% B, 1.4 mL/min, 25 °C, 254 nm</p>
----	---	--	--	--


32		N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture)	N-[(cis)-4-hydroxytetrahydrothiophen-3-yl]-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 2.72 - 2.84 (m, 2H, minor), 2.87 - 2.97 (m, 2H, major), 3.12 (dd, 1H, major), 3.62 (dd, 1H, major), 3.84 (dd, 1H, minor), 3.93 - 3.94 (m, 3H), 4.43 - 4.48 (m, 1H, minor), 4.49 - 4.57 (m, 1H, minor), 4.59 - 4.64 (m, 1H, major), 5.00 - 5.10 (m, 1H, major), 5.88 (d, 1H, minor), 5.92 (d, 1H, major), 7.90 (d, 2H), 8.12 - 8.14 (m, 1H), 8.29 - 8.34 (m, 2H), 8.57 (s, 1H, major), 8.59 (s, 1H, minor), 8.68 (s, 1H, minor), 8.70 (s, 1H, major), 9.92 (d, 1H, major), 10.05 (d, 1H, minor).
----	---	--	---	--


33		N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture)	N-[(cis)-4-hydroxytetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 2.71 - 2.80 (m, 2H, minor), 2.85 - 2.95 (m, 2H, major), 3.11 (dd, 1H, major), 3.59 (d, 1H, major), 3.82 (dd, 1H, minor), 4.40 - 4.45 (m, 1H, minor), 4.48 - 4.55 (m, 1H, minor), 4.59 (t, 1H, major), 4.98 - 5.07 (m, 1H, major), 5.91 (br s, 1H), 7.64 (dd, 1H), 7.89 (d, 2H), 8.18 (ddd, 1H), 8.23 (d, 2H), 8.71 (dd, 1H), 8.76 (s, 1H, minor), 8.79 (s, 1H, major), 8.90 - 8.94 (m, 1H), 9.83 (d, 1H, major), 9.94 (d, 1H, minor).
----	---	--	---	--

34		<p>N-[(cis)-4-hydroxy-1-oxido-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, isomer 1, $[\alpha]_D^{20} = -10.3^\circ$ (c = 1.00, DMSO)</p>	<p>N-[(cis)-4-hydroxy-1-oxido-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture), Chiralpak ID 5μ 250x30mm, eluent A: tert-butylmethylether + 0.1vol% diethylamine (99%), eluent B: methanol, isocratic: 50% B, 40 mL/min, 254 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 2.71 - 2.80 (m, 2H), 3.34 - 3.39 (m, 1H and water signal), 3.82 (dd, 1H), 4.40 - 4.45 (m, 1H), 4.48 - 4.56 (m, 1H), 5.85 (d, 1H), 7.65 (ddd, 1H), 7.89 (d, 2H), 8.18 (ddd, 1H), 8.22 (d, 2H), 8.71 (dd, 1H), 8.76 (s, 1H), 8.93 (d, 1H), 9.94 (d, 1H). Rt = 2.65 min, Chiralpak ID 3μ 100x4.6mm, eluent A: tert.-butylmethylether + 0.1vol% diethylamine (99%), eluent B: methanol, isocratic: 50% B, 1.4 mL/min, 25 °C, 254 nm</p>
----	---	--	--	---

35		<p>N-[(cis)-4-hydroxy-1-oxido-2-(pyridin-3-yl)-6-(4-(trifluoromethyl)phenyl)-2,3-dihydropyridazin-4-yl]-3-oxo-2-(pyridin-3-yl)-6-(4-(trifluoromethyl)phenyl)-2,3-dihydropyridazin-4-carboxamide, isomer 2, $[\alpha]_D^{20} = +15.6^\circ$ (c = 1.00, DMSO)</p>	<p>N-[(cis)-4-hydroxy-1-oxido-2-(pyridin-3-yl)-6-(4-(trifluoromethyl)phenyl)-2,3-dihydropyridazin-4-yl]-3-oxo-2-(pyridin-3-yl)-6-(4-(trifluoromethyl)phenyl)-2,3-dihydropyridazin-4-carboxamide (diastereomeric mixture), Chiralpak ID 5μ 250x30mm, eluent A: tert-butylmethylether + 0.1vol% diethylamine (99%), eluent B: methanol, isocratic: 50% B, 40 mL/min, 254 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 2.71 - 2.80 (m, 2H), 3.34 - 3.39 (m, 1H and water signal), 3.82 (dd, 1H), 4.40 - 4.45 (m, 1H), 4.48 - 4.56 (m, 1H), 5.85 (d, 1H), 7.65 (ddd, 1H), 7.89 (d, 2H), 8.18 (ddd, 1H), 8.22 (d, 2H), 8.71 (dd, 1H), 8.76 (s, 1H), 8.93 (d, 1H), 9.94 (d, 1H). Rt = 3.78 min, Chiralpak ID 3μ 100x4.6mm, eluent A: tert.-butylmethylether + 0.1vol% diethylamine (99%), eluent B: methanol, isocratic: 50% B, 1.4 mL/min, 25 °C, 254 nm</p>
----	---	---	---	---

36		<p>N-[(cis)-4-hydroxy-1-oxido-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, isomer 3,</p>	<p>N-[(cis)-4-hydroxy-1-oxido-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture), Chiralpak ID 5μ 250x30mm, eluent A: tert-butylmethylether + 0.1vol% diethylamine (99%), eluent B: methanol, isocratic: 50% B, 40 mL/min, 254 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 2.84 - 2.95 (m, 2H), 3.11 (dd, 1H), 3.59 (dd, 1H), 4.57 - 4.62 (m, 1H), 4.98 - 5.07 (m, 1H), 5.90 (d, 1H), 7.65 (ddd, 1H), 7.89 (d, 2H), 8.17 (ddd, 1H), 8.23 (d, 2H), 8.71 (dd, 1H), 8.79 (s, 1H), 8.92 (d, 1H), 9.82 (d, 1H). Rt = 4.40 min, Chiralpak ID 3μ 100x4.6mm, eluent A: tert-butylmethylether + 0.1vol% diethylamine (99%), eluent B: methanol, isocratic: 50% B, 1.4 mL/min, 25 °C, 254 nm</p>
----	---	---	---	---

37		<p>N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, isomer 4,</p>	<p>N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture), Chiralpak ID 5μ 250x30mm, eluent A: tert-butylmethylether + 0.1vol% diethylamine (99%), eluent B: methanol, isocratic: 50% B, 40 mL/min, 254 nm</p>	<p>¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 2.84 - 2.95 (m, 2H), 3.11 (dd, 1H), 3.59 (dd, 1H), 4.59 (q, 1H), 4.98 - 5.07 (m, 1H), 5.90 (d, 1H), 7.64 (ddd, 1H), 7.89 (d, 2H), 8.17 (ddd, 1H), 8.23 (d, 2H), 8.71 (dd, 1H), 8.79 (s, 1H), 8.92 (d, 1H), 9.82 (d, 1H). Rt = 5.20 min, Chiralpak ID 3μ 100x4.6mm, eluent A: tert-butylmethylether + 0.1vol% diethylamine (99%), eluent B: methanol, isocratic: 50% B, 1.4 mL/min, 25 °C, 254 nm</p>
----	---	---	---	--

Example 38**2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(S-methylsulfonimidoyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide**

5


Step 1: 2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (342 mg, 732 μ mol) was suspended in dichloromethane (7.9 mL). 2,2,2-Trifluoroacetamide (413 mg, 3.66 mmol), (diacetoxyiodo)benzene (884 mg, 2.74 mmol), rhodium(II) acetate dimer (80.8 mg, 183 μ mol), and magnesium oxide (295 mg, 7.32 mmol) were added. It was stirred at rt overnight. The reaction mixture was diluted with dichloromethane (40 mL), washed with water (15 mL) dried over magnesium sulfate and concentrated.

10

Step 2: The residue of step 1 was dissolved in methanol (16 mL) and potassium carbonate (202 mg, 1.46 mmol) was added. It was stirred overnight at rt. This reaction mixture was combined with a second batch (50 mg of starting material for step 1 and treated analogously) diluted with dichloromethane (50 mL), washed with water (20 mL) dried over magnesium sulfate and concentrated. The residue was purified by RP-HPLC (column: X-Bridge C18 5 μ m 100x30 mm, mobile phase: (water + 0.2 vol% aqueous ammonia (32%)) / acetonitrile, gradient) to give 143 mg (35%) of the title compound as diastereomeric mixture of ca. 1:1.

$^1\text{H-NMR}$ (400MHz, DMSO- d_6): δ [ppm] = 1.37 (d, 6H), 2.96 (s, 6H), 3.25 - 3.33 (m, 2H and water signal), 3.51 - 3.57 (m, 2H), 3.77 (d, 2H), 3.94 (s, 6H), 4.50 - 4.62 (m, 2H), 7.89 (d, 4H), 8.14 (s, 2H), 8.30 (d, 4H), 8.55 (s, 2H), 8.63 (s, 1H), 8.63 (s, 1H), 9.51 - 9.59 (m, 2H).

Example 392-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(S-methylsulfonimidoyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, diastereomer 1



2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(S-methylsulfonimidoyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (135 mg) was separated by chiral HPLC (column: Chiralpak IB 5 μ m 250x30 mm, mobile phase: eluent A: CO₂, eluent B: ethanol + 0.2vol% aqueous ammonia (32%), isocratic: 23% B, 100 mL/min, 40 °C, BPR: 150 bar, 220 nm) to obtain 58 mg of the title compound.


$^1\text{H-NMR}$ (400MHz, DMSO- d_6): δ [ppm] = 1.37 (d, 3H), 2.96 (s, 3H), 3.29 - 3.32 (m, 1H and water signal), 3.54 (dd, 1H), 3.75 (s, 1H), 3.94 (s, 3H), 4.50 - 4.62 (m, 1H), 7.89 (d, 2H), 8.13 (s, 1H), 8.29 (d, 2H), 8.55 (s, 1H), 8.63 (s, 1H), 9.57 (d, 1H).

Chiral HPLC: Rt = 2.12 min

Chiralpak IB 5 μ m 100x4.6 mm; eluent A: CO₂, eluent B: ethanol + 0.2vol% aqueous ammonia (32%), isocratic: 28% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 220 nm.

$[\alpha]_D^{20}$ = +55.4° (c = 1.00, DMSO).

Example 402-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(S-methylsulfonimidoyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide, diastereomer 2



2-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(S-methylsulfonimidoyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (135 mg) was separated by chiral HPLC (column: Chiralpak IB 5 μ m 250x30 mm, mobile phase: eluent A: CO₂, eluent B: ethanol + 0.2vol% aqueous ammonia (32%), isocratic: 23% B, 100 mL/min, 40 °C, BPR: 150 bar, 220 nm) to obtain 53 mg of the title compound.


10 ¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.37 (d, 3H), 2.96 (s, 3H), 3.28 (dd, 1H), 3.53 (dd, 1H), 3.79 (s, 1H), 3.94 (s, 3H), 4.50 - 4.62 (m, 1H), 7.89 (d, 2H), 8.14 (s, 1H), 8.30 (d, 2H), 8.55 (s, 1H), 8.62 (s, 1H), 9.53 (d, 1H).

Chiral HPLC: Rt = 2.96 min

15 Chiralpak IB 5 μ m 100x4.6 mm; eluent A: CO₂, eluent B: ethanol + 0.2vol% aqueous ammonia (32%), isocratic: 28% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 220 nm.

$[\alpha]_D^{20} = +65.1^\circ$ (c = 1.00, DMSO).

20 **Example 412-(1-Methyl-1H-pyrazol-4-yl)-N-[(2S)-1-[S-methyl-N-(trifluoroacetyl)sulfonimidoyl]propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide**







50 mg from step 1 of the synthesis of 2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(S-methylsulfonimidoyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide was separated by HPLC to obtain 8 mg of the title compound.

- 5 ¹H-NMR (400MHz, DMSO-d₆): δ [ppm] = 1.41 (2x d, 3H, 2 isomers), 3.62 (s, 3H, 1 isomer), 3.63 (s, 3H, 1 isomer), 3.92 - 4.01 (m, 4H), 4.25 (dd, 1H), 4.70 - 4.82 (m, 1H), 7.89 (d, 2H), 8.15 (2x s, 1H, 2 isomers), 8.30 (d, 2H), 8.54 (s, 1H), 8.64 (s, 1H, 1 isomer), 8.65 (s, 1H, 1 isomer), 9.65 (d, 1H).
- 10 The following examples were prepared from the starting materials stated in the table using the procedure described in the examples above. Enantiomers/diastereomers were separated from their racemate/diastereomeric mixture by chiral HPLC using the column and solvent conditions stated.

Table 4: Examples 42 – 47


Expl	Structure	IUPAC name	Starting materials	Analytics
42		6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 1)	6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 1)	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 3.06 - 3.25 (m, 2H), 3.39 - 3.51 (m, 2H), 4.15 (s, 1H, minor), 4.31 (s, 1H, major), 4.48 - 4.57 (m, 1H), 4.61 - 4.70 (m, 1H, minor), 4.74 - 4.83 (m, 1H, major), 6.12 (d, 1H, major), 6.22 (d, 1H, minor), 7.36 - 7.42 (m, 1H), 7.53 - 7.66 (m, 5H), 7.99 - 8.03 (m, 2H), 8.69 (s, 1H, minor), 8.69 (s, 1H, major), 9.96 (d, 1H).
43		6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, isomer 1, [α] _D ²⁰ = -12.7° (c = 1.00, DMSO)	6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 1), Chiralpak IA 5μ 250x30mm, eluent A: CO ₂ , eluent B: ethanol, isocratic: 59% B, 59% B, 100 mL/min, 40 °C, BPR: 150 bar, 254 nm	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 3.11 - 3.21 (m, 2H), 3.41 - 3.50 (m, 2H), 4.15 (s, 1H), 4.48 - 4.51 (m, 1H), 4.62 - 4.70 (m, 1H), 6.22 (d, 1H), 7.36 - 7.42 (m, 1H), 7.53 - 7.66 (m, 5H), 7.98 - 8.03 (m, 2H), 8.69 (s, 1H), 9.96 (d, 1H). Rt = 2.24 min, Chiralpak IA 5μ 100x4.6mm, eluent A: CO ₂ , eluent B: ethanol, isocratic: 59% B, 59% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 254 nm

44		6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 2)	6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 2)	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 3.06 - 3.24 (m, 2H), 3.40 - 3.50 (m, 2H), 4.15 (s, 1H, minor), 4.31 (s, 1H, major), 4.49 - 4.56 (m, 1H), 4.66 (dtd, 1H, minor), 4.78 (dtd, 1H, major), 6.12 (d, 1H, major), 6.22 (d, 1H, minor), 7.37 - 7.42 (m, 1H), 7.53 - 7.66 (m, 5H), 7.98 - 8.04 (m, 2H), 8.69 (s, 1H, minor), 8.69 (s, 1H, major), 9.96 (d, 1H).
45		6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, diastereomer 1, [α] _D ²⁰ = +16.5° (c = 1.00, DMSO)	6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 2), Chiralpak IC 5μ 250x30mm, eluent A: CO ₂ , eluent B: ethanol, isocratic: 42% B, 100 mL/min, 40 °C, BPR: 150 bar, 254 nm	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 3.11 - 3.21 (m, 2H), 3.42 - 3.50 (m, 2H), 4.15 (s, 1H), 4.48 - 4.53 (m, 1H), 4.61 - 4.70 (m, 1H), 6.22 (d, 1H), 7.36 - 7.42 (m, 1H), 7.53 - 7.66 (m, 5H), 7.99 - 8.03 (m, 2H), 8.69 (s, 1H), 9.96 (d, 1H). Rt = 2.18 min, Chiralpak IC 5μ 100x4.6mm, eluent A: CO ₂ , eluent B: ethanol, isocratic: 42% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 254 nm

46		6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, diastereomer 2, $[\alpha]_D^{20} = -5.1^\circ$ (c = 1.00, DMSO)	6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture 2), Chiralpak IC 5 μ 250x30mm, eluent A: CO ₂ , eluent B: ethanol, isocratic: 42% B, 100 mL/min, 40 °C, BPR: 150 bar, 254 nm	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 3.10 (t, 1H), 3.21 (dd, 1H), 3.39 - 3.49 (m, 2H), 4.31 (s, 1H), 4.50 - 4.56 (m, 1H), 4.78 (dtd, 1H), 6.12 (d, 1H), 7.36 - 7.42 (m, 1H), 7.53 - 7.65 (m, 5H), 7.98 - 8.03 (m, 2H), 8.69 (s, 1H), 9.96 (d, 1H). Rt = 3.41 min, Chiralpak IC 5 μ 100x4.6mm, eluent A: CO ₂ , eluent B: ethanol, isocratic: 42% B, 4 mL/min, 37.5 °C, BPR: 100 bar, 254 nm
47		N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture)	N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide (diastereomeric mixture)	¹ H-NMR (400MHz, DMSO-d ₆): δ [ppm] = 3.09 - 3.28 (m, 3H), 3.41 - 3.52 (m, 3H), 3.94 (s, 3H), 4.17 (s, 1H, minor), 4.33 (s, 1H, major), 4.52 - 4.60 (m, 1H), 4.64 - 4.74 (m, 1H, minor), 4.76 - 4.85 (m, 1H, major), 6.15 (d, 1H, major), 6.25 (d, 1H, minor), 7.90 (d, 2H), 8.13 (s, 1H), 8.31 (br d, 2H), 8.57 (s, 1H), 8.69 (s, 1H), 10.01 (d, 1H).

Example 48

6-(4-Chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfanyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide




6-(4-Chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid
 250 mg, 0.756 mmol) was dissolved in dimethylformamide (5 mL) and treated with (2S)-1-
 (methylsulfonyl)propan-2-amine (159 mg, 1.51 mmol), N-ethyl-N-diisopropylamine (0.39 mL,
 5 2.27 mmol), 4-dimethylaminopyridine (4.6 mg, 0.038 mmol) and HATU (575 mg, 1.5 mmol).
 The reaction mixture was stirred overnight at rt and then purified by HPLC (method D) to yield
 63 mg (19%) of the title compound.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm = 1.28 (d, 3 H), 2.12 (s, 3 H), 2.67 - 2.78 (m, 2 H), 3.93
 10 (s, 3 H), 4.19 - 4.29 (m, 1 H), 7.57 - 7.62 (m, 2 H), 8.07 - 8.14 (m, 3 H), 8.57 (d, 2 H), 9.52 (d, 1
 H).

Example 49

**6-(4-Chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfonyl)propan-2-yl]-3-
 15 oxo-2,3-dihydropyridazine-4-carboxamide**




6-(4-Chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfonyl)propan-2-yl]-3-oxo-
 2,3-dihydropyridazine-4-carboxamide (55 mg, 0.13 mmol) was dissolved in acetone (0.3 mL),
 methanol (0.18 mL) and water (0.6 mL) followed by the addition of sodium periodate (56 mg,
 20 0.26 mmol). The reaction mixture was stirred at rt overnight followed by the addition of water.
 After stirring overnight a precipitate formed which was filtered off, washed with water and dried
 in vacuum. The residue was purified by HPLC (method C) to yield 34 mg (59%) of the title
 compound.

^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ ppm = 1.33 - 1.40 (m, 3 H), 2.61 (d, 3 H), 2.89 - 3.19 (m, 2 H), 3.93 (s, 3 H), 4.39 - 4.55 (m, 1 H), 7.58 - 7.62 (m, 2 H), 8.07 - 8.15 (m, 3 H), 8.51 - 8.60 (m, 2 H), 9.55 - 9.61 (m, 1 H).

5 **Example 50**

6-(4-Chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfonyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide




6-(4-Chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-N-[1-(methylsulfonyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide (5 mg, 8%) was isolated as a byproduct in the synthesis of 6-(4-chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide.

^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ ppm = 1.37 (d, 3 H), 3.03 (s, 3 H), 3.38 - 3.46 (m, 1 H), 3.57 - 3.64 (m, 1 H), 3.93 (s, 3 H), 4.53 - 4.64 (m, 1 H), 7.60 (d, 2 H), 8.07 - 8.14 (m, 3 H), 8.55 (d, 2 H), 9.55 - 9.61 (m, 1 H).

Example 51

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[2-(methylsulfinyl)ethyl]-3-oxo-2,3-dihydropyridazine-4-carboxamide




6-(4-Chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid (200 mg, 0.58 mmol) was dissolved in dimethylformamide (4 mL) and treated with 2-(methylsulfinyl)ethanamine (124.36 mg, 1.16 mmol), N-ethyl-N-diisopropylamine (0.33 mL, 1.74 mmol), 4-dimethylaminopyridine (5.3 mg, 0.03 mmol) and HATU (441.2 mg, 1.16 mmol).

The reaction mixture was stirred overnight at rt and then purified by HPLC (method D) to yield 101 mg (38%) of the title compound.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm = 2.60 (s, 3 H), 2.88 - 2.97 (m, 1 H), 3.03 - 3.12 (m, 1 H),
 5 3.75 (q, 2 H), 7.34 - 7.42 (m, 1 H), 7.52 - 7.65 (m, 5 H), 7.99 (dd, 2 H), 8.64 (d, 1 H), 9.58 (t, 1 H).

Example 52

10 **6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[2-(S-methylsulfonimidoyl)ethyl]-3-oxo-2,3-dihydropyridazine-4-carboxamide**




15 6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-{2-[S-methyl-N-(trifluoroacetyl)sulfonimidoyl]ethyl}-3-oxo-2,3-dihydropyridazine-4-carboxamide (25 mg, 0.046 mmol) was dissolved in methanol (1 mL) and treated with potassium carbonate (12.6 mg, 0.92 mmol). The reaction mixture was stirred overnight and then diluted with dichloromethane and brine. The phases were separated and the organic phase was filtered (MN 617 WA filter paper) and concentrated to yield 18.6 mg (90%) of the title compound.

20 ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm = 2.95 (s, 3 H), 3.79 (q, 2 H), 3.86 (s, 1 H), 7.34 - 7.42 (m, 1 H), 7.52 - 7.66 (m, 5 H), 7.97 - 8.03 (m, 2 H), 8.64 (s, 1 H), 9.60 - 9.65 (m, 1 H).

Example 53

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[2-(S-methylsulfonimidoyl)ethyl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, Enantiomer 1



25


6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[2-(S-methylsulfonimidoyl)ethyl]-3-oxo-2,3-dihydropyridazine-4-carboxamide was separated by chiral HPLC (Instrument: Sepiatec: Prep SFC100; Säule: Chiralpak IA 5µm 250x30mm; eluent A: CO₂, eluent B: ethanol; isocratic: 60%B; flow 100.0 mL/min temperature: 40°C; BPR: 150bar; MWD @ 335) to yield 4.1 mg (40%) of the title compound.

Chiral HPLC: Rt = 6.22 min

Instrument: Agilent: 1260, Aurora SFC-Modul; column: Chiralpak IA 5µm 100x4.6mm; eluent A: CO₂, eluent B: ethanol; isocratic: 56%B; flow 4.0 mL/min; temperature: 37.5°C; BPR: 100bar; MWD @ 335nm.

Example 54

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[2-(S-methylsulfonimidoyl)ethyl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, Enantiomer 2



15

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[2-(S-methylsulfonimidoyl)ethyl]-3-oxo-2,3-dihydropyridazine-4-carboxamide was separated by chiral HPLC (Instrument: Sepiatec: Prep SFC100; Säule: Chiralpak IA 5µm 250x30mm; eluent A: CO₂, eluent B: ethanol; isocratic: 60%B; flow 100.0 mL/min temperature: 40°C; BPR: 150bar; MWD @ 335) to yield 3.4 mg (35%) of the title compound.

20


Chiral HPLC: Rt = 8.39 min

Instrument: Agilent: 1260, Aurora SFC-Modul; column: Chiralpak IA 5µm 100x4.6mm; eluent A: CO₂, eluent B: ethanol; isocratic: 56%B; flow 4.0 mL/min; temperature: 37.5°C; BPR: 100bar; MWD @ 335nm.

25

Example 55

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide




6-(4-Chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid (75 mg, 0.22 mmol) was dissolved in dimethylformamide (1.5 mL) and treated with *cis*-4-aminotetrahydrothiophene-3-ol 1,1-dioxide (69.2 mg, 0.43 mmol), *N*-ethyl-*N*-diisopropylamine (0.12 mL, 0.65 mmol), 4-dimethylaminopyridine (2.0 mg, 0.011 mmol) and HATU (165.4 mg, 0.43 mmol). The reaction mixture was stirred at rt overnight and then purified by HPLC (method D) to yield 43.6 mg (42%) of the title compound.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm = 3.08 - 3.18 (m, 1 H), 3.27 (br d, 1 H), 3.48 - 3.56 (m, 2 H), 4.53 (br d, 1 H), 4.75 (dtd, 1 H), 6.31 - 6.35 (m, 1 H), 7.35 - 7.43 (m, 1 H), 7.52 - 7.66 (m, 5 H), 7.97 - 8.04 (m, 2 H), 8.69 (s, 1 H), 9.94 - 10.00 (m, 1 H).

Example 56

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-*N*-[(*cis*)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, Enantiomer 1




6-(4-Chlorophenyl)-2-(3-fluorophenyl)-*N*-[(*cis*)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide was separated by chiral HPLC (Instrument: Labomatic HD5000, Labocord-5000; Gilson GX-241, Labcol Vario 4000, column: Chiralpak IB 5μ 250x30mm; eluent A: hexan + 0.1 vol-% diethylamine (99%); eluent B: ethanol; isocratic: 50%A+50%B; flow 50.0 mL/min; UV 254 nm) to yield 10.0 mg (9%) of the title compound.

Chiral HPLC: Rt = 4.18 min

Instrument: Agilent HPLC 1260; column: Chiralpak IB 3 μ 100x4,6mm; eluent A: hexan + 0.1 vol-% diethylamine (99%); eluent B: ethanol; isocratic: 50%A+50%B; flow 1.4 mL/min; temperature: 25 °C; DAD 254 nm.

5 **Example 57**

6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, Enantiomer 2




10 Rac-6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide was separated by chiral HPLC (Instrument: Labomatic HD5000, Labocord-5000; Gilson GX-241, Labcol Vario 4000, column: Chiralpak IB 5 μ 250x30mm; eluent A: hexan + 0.1 Vol-% diethylamine (99%); eluent B: ethanol; isocratic: 50%A+50%B; flow 50.0 mL/min; UV 254 nm) to yield 10.0 mg (9%) of the title compound.

15 Chiral HPLC: Rt = 5.28 min

Instrument: Agilent HPLC 1260; column: Chiralpak IB 3 μ 100x4,6mm; eluent A: hexan + 0.1 Vol-% diethylamine (99%); eluent B: ethanol; isocratic: 50%A+50%B; flow 1.4 mL/min; temperature: 25 °C; DAD 254 nm.

20 **Example 58**

6-(4-Chlorophenyl)-N-(1,1-dioxidotetrahydrothiophen-3-yl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxamide




25 6-(4-Chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid (75 mg, 0.22 mmol) was dissolved in dimethylformamide (1.5 mL) and treated with tetrahydrothiophen-3-amine 1,1-dioxide (58.8 mg, 0.43 mmol), N-ethyl-N-diisopropylamine (0.12 mL, 0.65 mmol),

4-dimethylaminopyridine (2.0 mg, 0.011 mmol) and HATU (165.4 mg, 0.43 mmol). The reaction mixture was stirred at rt overnight and then purified by HPLC (method D) to yield 45.8 mg (46%) of the title compound.

- 5 ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ ppm = 2.15 - 2.28 (m, 1 H), 3.14 - 3.24 (m, 2 H), 3.24 - 3.31 (m, 1 H), 3.46 - 3.54 (m, 1 H), 4.76 (sxt, 1 H), 7.35 - 7.42 (m, 1 H), 7.51 - 7.66 (m, 5 H), 7.97 - 8.03 (m, 2 H), 8.64 (s, 1 H), 9.67 (d, 1 H).

Example 59

- 10 **6-(4-Chlorophenyl)-2-(3-fluorophenyl)-N-[2-(methylsulfonyl)ethyl]-3-oxo-2,3-dihydropyridazine-4-carboxamide**



- 15 6-(4-Chlorophenyl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxylic acid (75 mg, 0.22 mmol) was dissolved in dimethylformamide (1.5 mL) and treated with 2-(methylsulfonyl)ethanamine (53.6 mg, 0.43 mmol), N-ethyl-N-diisopropylamine (0.12 mL, 0.65 mmol), 4-dimethylaminopyridine (2.0 mg, 0.011 mmol) and HATU (165.4 mg, 0.43 mmol). The reaction mixture was stirred at rt overnight and then purified by HPLC (method D) to yield 33.5 mg (34%) of the title compound.

- 20 ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ ppm = 3.04 (s, 3 H), 3.40 - 3.44 (m, 2 H), 3.80 (q, 2 H), 7.35 - 7.42 (m, 1 H), 7.52 - 7.66 (m, 5 H), 7.97 - 8.03 (m, 2 H), 8.64 (s, 1 H), 9.58 - 9.64 (m, 1 H).

EXPERIMENTAL SECTION – BIOLOGICAL ASSAYS

- 25 Examples were tested in selected biological assays one or more times. When tested more than once, data are reported as either average values or as median values, wherein

- the average value, also referred to as the arithmetic mean value, represents the sum of the values obtained divided by the number of times tested, and
 - the median value represents the middle number of the group of values when ranked in ascending or descending order. If the number of values in the data set is odd, the
- 30

median is the middle value. If the number of values in the data set is even, the median is the arithmetic mean of the two middle values.

Examples were synthesized one or more times. When synthesized more than once, data from biological assays represent average values or median values calculated utilizing data sets obtained from testing of one or more synthetic batch.

The *in vitro* activity of the compounds of the present invention can be demonstrated in the following assays:

Transactivation Assay in human cell line (in vitro assays 1 and 2)

Transactivation assays were carried out in U87 glioblastoma cells (ATCC) endogenously expressing AHR. In addition the cells were stably transfected with an AHR inducible firefly luciferase reporter gene construct that carried AHR-binding sites (DRE) in its promoter and a renilla reporter gene construct with constitutively active promoter. Kynurenic acid is an endogenous AHR activating ligand and was used to prestimulate test cells prior to testing the antagonistic properties of compounds.

***In vitro* assay 1: Antagonism in human cell line**

Cells in medium (tryptophan free RPMI, 1% FCS, 2 mM Glutamine) supplemented with 150 μ M kynurenic acid were grown for 20 hours in absence (negative control) or presence of increasing concentrations of test compounds (typical dilutions: 72 pmol/L, 0.25 nmol/L, 0.89 nmol/L; 3.1 nmol/L, 11 nmol/L, 38 nmol/L, 130 nmol/L, 470 nmol/L, 1.6 μ mol/L, 5.7 μ mol/L and 20 μ mol/L in duplicates). As positive inhibition control cells supplemented with 150 μ M kynurenic acid were incubated in presence of 5 μ M Staurosporin. Normalization was done by positive and negative controls.

Firefly luciferase and Renilla activity was determined by the DualGlo Luciferase Assay System (Promega, #2920). Renilla activity was used to assess toxic effects of compounds.

***In vitro* assay 2: Agonism in human cell line**

Cells in medium (tryptophan free RPMI, 1% FCS, 2 mM Glutamine) were grown for 20 hours in absence (negative control) or presence of increasing concentrations of test compounds (typical dilutions: 72 pmol/L, 0.25 nmol/L, 0.89 nmol/L; 3.1 nmol/L, 11 nmol/L, 38 nmol/L, 130 nmol/L, 470 nmol/L, 1.6 μ mol/L, 5.7 μ mol/L and 20 μ mol/L in duplicates). As positive activation control cells were incubated with 300 μ M kynurenic acid. Normalization was done by positive and negative controls.

Firefly luciferase activity was determined by the SteadyGlo Luciferase Assay System (Promega, #2520).

***In vitro* assay 3: AHR-regulated CYP1A1 expression in human cell line**

To assess the AHR inhibitory activity of the substances described in this application, the ability thereof to antagonise ligand-induced AHR gene regulation in a dose-dependent manner was quantified. For this purpose, quantitative PCR analysis was used to determine expression of the AHR-regulated gene CYP1A1 in a human monocytic U937 cell line upon stimulation with 200 μ M KA in the presence and absence of AHR inhibitor. U937 cells were sown at a concentration of 2×10^5 cells/well in 100 μ l of growth medium (RPMI 1640, 20% FCS) in 96-well microtitre plates. CYP1A1 expression was induced with 200 μ M KA (positive control) in the presence or absence of the substances for 6 hours. Human U937 cells were typically incubated with eight different concentrations of the substances (1 nM, 3 nM, 10 nM, 30 nM, 100 nM, 300 nM, 1 μ M and 3 μ M) and analyzed in duplicate on the same microtitre plate. After stimulation, cells were lysed with Nucleic Acid Lysis Solution (# 4305895, Applied Biosystems) and RNA was isolated using the 6100 Nucleic Acid Preparation Station (Applied Biosystems) and reverse-transcribed to cDNA using SuperScript VILO cDNA synthesis kit (# 11754-250, Invitrogen). Unstimulated cells were used as the negative control. Taqman probes for human CYP1A1 (Hs01054797_g1) and human HPRT (Hs02800695_m1) were used to analyze fold expression of CYP1A1 of HPRT. Quantitation was performed on a Taqman SDS7900HT.

Table 4: IC₅₀ values of examples in *in vitro* assays 1 - 3


Example	Assay 1: AHR-luc Hum Antagonism IC ₅₀ [M]	Assay 2: AHR-luc Hum Agonism IC ₅₀ [M]	Assay 3: Hum CYP1A1 Antagonism IC ₅₀ [M]
1	5,26 E-8	> 2,00 E-5	
2	1,47 E-7	> 2,00 E-5	
3	4,42 E-8	> 2,00 E-5	
4	4,02 E-8	> 2,00 E-5	
5	1,80 E-7	> 2,00 E-5	
6	2,32 E-7	> 2,00 E-5	
7	2,41 E-7	> 2,00 E-5	
8	1,40 E-7	> 2,00 E-5	
9	6,99 E-9	> 2,00 E-5	
10	2,41 E-9	> 2,90 E-5	
11	3,46 E-9	> 2,00 E-5	
12	1,11 E-9	5,41 E-6	
13	5,65 E-9	> 2,00 E-5	1,86 E-8

14	6,45 E-9	> 2,00 E-5	
15	1,77 E-8	> 2,00 E-5	3,69 E-8
16	2,92 E-8	> 2,00 E-5	
17	1,62 E-8	> 2,00 E-5	
18	3,82 E-8	> 2,00 E-5	
19	6,99 E-8	> 2,00 E-5	
20	5,43 E-8	1,24 E-5	
21	2,05 E-7	> 2,00 E-5	
22	2,19 E-8	> 2,00 E-5	
23	5,37 E-9	> 2,00 E-5	
24	7,54 E-9	> 2,00 E-5	7,56 E-9
25	4,14 E-8	> 2,00 E-5	
26	3,33 E-8	> 2,00 E-5	
27	9,35 E-8	> 2,00 E-5	
28	1,34 E-8	> 2,00 E-5	
29	3,01 E-7	> 2,00 E-5	
30	2,10 E-7	> 2,00 E-5	
31	1,45 E-7	> 2,00 E-5	
32	4,16 E-8	> 2,00 E-5	
33	1,39 E-7	> 2,00 E-5	
34	7,88 E-8	> 2,00 E-5	
35	2,72 E-7	> 2,00 E-5	
36	2,95 E-7	> 2,00 E-5	
37	1,50 E-7	> 2,00 E-5	
38	5,48 E-8	> 2,00 E-5	
39	1,33 E-7	> 2,00 E-5	
40	8,03 E-8	> 2,00 E-5	
41	3,92 E-8	> 2,00 E-5	

42	3,49 E-8	> 2,00 E-5	
43	4,38 E-8	> 2,00 E-5	
44	9,93 E-8	> 2,00 E-5	
45	5,69 E-8	> 2,00 E-5	7,56 E-7
46	1,03 E-7	> 2,00 E-5	
47	1,58 E-7	> 2,00 E-5	
48	1,55 E-8	> 2,00 E-5	2,99 E-8
49	2,64 E-8	> 2,00 E-5	
50	3,05 E-8	> 2,00 E-5	4,37 E-8
51	3,93 E-7	> 2,00 E-5	
52	7,00 E-7	> 2,00 E-5	
53	8,43 E-7	> 2,00 E-5	
54	5,81 E-7	> 2,00 E-5	
55	2,65 E-8	> 2,00 E-5	2,70 E-8
56	6,53 E-8	> 2,00 E-5	
57	5,77 E-9	> 2,00 E-5	1,71 E-8
58	1,45 E-7	> 2,00 E-5	
59	3,26 E-7	> 2,00 E-5	

CLAIMS

1. A compound of general formula (I):



(I),

5 in which

A represents phenyl or monocyclic heteroaryl, wherein said phenyl is substituted with R⁴, R⁵ and R⁶ and said monocyclic heteroaryl is optionally substituted one to three times, independently from each other, with R^{6a}.

X represents CR^{4a} or N;

10 Y represents CR³ or N;

Z represents CH or N, wherein

if X represents N, Y represents CR³ and Z represents CH, and

if X represents CR^{4a}, Z represents CH and Y represents CR³ or N, and

if Z represents N, Y represents N and X represents CH;

15 R¹ represents C₂-C₆-alkyl substituted once with R⁷ and optionally once with hydroxy and optionally one to three times with halogen, or

C₃-C₆-cycloalkyl substituted once with R⁷ and optionally once with hydroxy, or 5- to 6-membered heterocycloalkyl optionally substituted once with hydroxy or oxo;

20 R² represents chloro, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy or -NR⁸R⁹;

R³ represents hydrogen, halogen or methyl;

R⁴ represents hydrogen, methyl, fluoromethyl, difluoromethyl, trifluoromethyl, halogen or cyano;

R^{4a} represents hydrogen or halogen;

25 R⁵ represents hydrogen or halogen;

R⁶ represents hydrogen or halogen;

R^{6a} represents C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₆-cycloalkyl, C₁-C₄-alkoxy, halogen or cyano;

R⁷ represents -SR¹⁰, -SO-R¹⁰, -SO₂-R¹⁰, -SO₂NR⁸R⁹ or -SO(NR⁸)-R¹⁰;

30 R⁸ and R⁹ are the same or different and represent, independently from each other, hydrogen, C₁-C₃-alkyl or trifluoroacetyl, or together with the nitrogen atom to which they are attached form a 4- to 6-membered

nitrogen containing heterocyclic ring, said ring optionally containing one additional heteroatom selected from O, S, NH, NR^a in which R^a represents a C₁-C₄-alkyl group;

R¹⁰ represents C₁-C₄-alkyl;

5 their polymorphs, enantiomers, diastereomers, racemates, tautomers, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

2. The compound according to claim 1, wherein:

10 A represents phenyl or monocyclic heteroaryl, wherein said phenyl is substituted with R⁴ and said monocyclic heteroaryl is optionally substituted one time with R^{6a}.

X represents CR^{4a} or N;

Y represents CR³ or N;

Z represents CH or N, wherein

if X represents N, Y represents CR³ and Z represents CH, and

15 if X represents CR^{4a}, Z represents CH and Y represents CR³ or N, and

if Z represents N, Y represents N and X represents CH;

R¹ represents C₂-C₄-alkyl substituted once with R⁷ and optionally once with hydroxy, or C₃-C₆-cycloalkyl substituted once with R⁷ and optionally once with hydroxy, or 5- to 6-membered heterocycloalkyl optionally substituted once with hydroxy;

20 R² represents chloro, methyl, fluoromethyl, difluoromethyl or trifluoromethyl;

R³ represents hydrogen;

R⁴ represents methyl, halogen or cyano;

R^{4a} represents hydrogen;

R⁵ represents hydrogen;

25 R⁶ represents hydrogen;

R^{6a} represents C₁-C₄-alkyl, C₁-C₄-haloalkyl, halogen or cyano;

R⁷ represents -SR¹⁰, -SO-R¹⁰, -SO₂-R¹⁰, -SO₂NR⁸R⁹ or -SO(NR⁸)-R¹⁰;


R⁸ and R⁹ are the same or different and represent, independently from each other, hydrogen, C₁-C₃-alkyl or trifluoroacetyl;

30 R¹⁰ represents C₁-C₃-alkyl;

their polymorphs, enantiomers, diastereomers, racemates, tautomers, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

35 3. The compound according to claim 1 or 2, wherein:

A represents a group selected from:




wherein * indicates the point of attachment of said group with the rest of the molecule;

X represents CH;

5 Y represents CH or N;

Z represents CH;

R¹ represents ethyl or propyl substituted once with R⁷, or a group selected from:



10 wherein * indicates the point of attachment of said group with the rest of the molecule;

R² represents chloro or trifluoromethyl;

R⁷ represents -SR¹⁰, -SO-R¹⁰, -SO₂-R¹⁰ or -SO(NR⁸)-R¹⁰;

R⁸ represents hydrogen or trifluoroacetyl;

R¹⁰ represents methyl;

15 their polymorphs, enantiomers, diastereomers, racemates, tautomers, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.

4. The compound according to claim 1, 2 or 3 which is selected from the group consisting of:

20 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfanyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide

6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfonyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide

25 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide

6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-[(R)-methylsulfinyl]propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide

- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-[(S)-methylsulfinyl]propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-(S-methylsulfonylimidoyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 5 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-[(R)-S-methylsulfonylimidoyl]propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(2S)-1-[(S)-S-methylsulfonylimidoyl]propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfonyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 10 (trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2-(1,2-thiazol-4-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- N-[(3R,4S)-4-hydroxy-1,1-dioxo-1 λ ⁶-thiolan-3-yl]-3-oxo-2-(1,2-thiazol-4-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 15 N-[(3S,4R)-4-hydroxy-1,1-dioxo-1 λ ⁶-thiolan-3-yl]-3-oxo-2-(1,2-thiazol-4-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 2-(3-fluorophenyl)-N-[(3R,4S)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 20 (trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 2-(3-fluorophenyl)-N-[(3S,4R)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 25 N-[(3R,4S)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- N-[(3S,4R)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxamide
- 30 (trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxamide
- 2-(3-fluorophenyl)-N-[(3R,4S)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxamide

- 2-(3-fluorophenyl)-N-[(3S,4R)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-6-[6-(trifluoromethyl)pyridin-3-yl]-2,3-dihydropyridazine-4-carboxamide
- 2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfonyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 5 2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-[(R)-methylsulfinyl]propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-[(S)-methylsulfinyl]propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 10 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(3R,4S)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(3S,4R)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 15 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(1S,3S,4R)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(1R,3S,4R)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(1S,3R,4S)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 20 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(1R,3R,4S)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 25 N-[(cis)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- N-[(1S,3S,4R)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- N-[(1R,3S,4R)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 30 N-[(1S,3R,4S)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide

- N-[(1R,3R,4S)-4-hydroxy-1-oxidotetrahydrothiophen-3-yl]-3-oxo-2-(pyridin-3-yl)-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(S-methylsulfonimidoyl)propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 5 2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-[(R)-(S-methylsulfonimidoyl)]propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-[(S)-S-methylsulfonimidoyl]propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 10 2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-[S-methyl-N-(trifluoroacetyl)sulfonimidoyl]propan-2-yl]-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(3R,4S)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(3S,4R)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 15 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, diatereomer 1
- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(3S,4R)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, isomer 1
- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(3S,4R)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide, isomer 2
- 20 N-[(cis)-4-hydroxy-1-imino-1-oxidotetrahydro-1H-1lambda4-thiophen-3-yl]-2-(1-methyl-1H-pyrazol-4-yl)-3-oxo-6-[4-(trifluoromethyl)phenyl]-2,3-dihydropyridazine-4-carboxamide
- 6-(4-chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfanyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 25 6-(4-chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfinyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 6-(4-chlorophenyl)-2-(1-methyl-1H-pyrazol-4-yl)-N-[(2S)-1-(methylsulfonyl)propan-2-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide
- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[2-(methylsulfinyl)ethyl]-3-oxo-2,3-
- 30 dihydropyridazine-4-carboxamide
- 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[2-(S-methylsulfonimidoyl)ethyl]-3-oxo-2,3-dihydropyridazine-4-carboxamide

6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-{2-[(R)-S-methylsulfonimidoyl]ethyl}-3-oxo-2,3-dihydropyridazine-4-carboxamide

6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-{2-[(R)-S-methylsulfonimidoyl]ethyl}-3-oxo-2,3-dihydropyridazine-4-carboxamide

5 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(cis)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide

6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(3R,4S)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide


10 6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[(3S,4R)-4-hydroxy-1,1-dioxidotetrahydrothiophen-3-yl]-3-oxo-2,3-dihydropyridazine-4-carboxamide

6-(4-chlorophenyl)-N-(1,1-dioxidotetrahydrothiophen-3-yl)-2-(3-fluorophenyl)-3-oxo-2,3-dihydropyridazine-4-carboxamide

6-(4-chlorophenyl)-2-(3-fluorophenyl)-N-[2-(methylsulfonyl)ethyl]-3-oxo-2,3-dihydropyridazine-4-carboxamide

15 their polymorphs, enantiomers, diastereomers, racemates, tautomers, N-oxides, hydrates and solvates, as well as their physiological acceptable salts and solvates of these salts, as well as mixtures of the same.


5. A method of preparing a compound of general formula (I) according to any one of claims 1
20 to 4, said method comprising the step of allowing an intermediate compound of general formula (VII):



(VII),

25 in which R², A, X, Y and Z are as defined for the compound of general formula (I) according to any one of claims 1 to 3,


to react with a compound of general formula (VIII):



(III),

30 in which R¹ is as defined for the compound of general formula (I) according to any one of claims 1 to 3,

thereby giving a compound of general formula (I) :



(I),

in which R¹, R², A, X, Y and Z are as defined for the compound of general formula (I) according to any one of claims 1 to 3.

5

6. A compound of general formula (I) according to any one of claims 1 to 4 for use in the treatment or prophylaxis of a disease.

10 7. A pharmaceutical composition comprising a compound of general formula (I) according to any one of claims 1 to 4 and one or more pharmaceutically acceptable excipients.

8. A pharmaceutical combination comprising:

15


- one or more first active ingredients, in particular compounds of general formula (I) according to any one of claims 1 to 4, and
- one or more pharmaceutical active anti cancer compounds or
- one or more pharmaceutical active immune checkpoint inhibitors.

20 9. Use of a compound of general formula (I) according to any one of claims 1 to 4 for the treatment or prophylaxis of a disease.

10. Use of a compound of general formula (I) according to any one of claims 1 to 4 for the preparation of a medicament for the treatment or prophylaxis of a disease.

25 11. Use according to claim 9 or 10, wherein the disease is cancer or conditions with dysregulated immune responses or other disorders associated with aberrant AHR signaling, such as liquid and solid tumours, for example.

12. A compound of general formula (VII):




(VII),

in which R^2 , A, X, Y and Z are as defined for the compound of general formula (I) according to any one of claims 1 to 3.

5

13. Use of a compound of general formula (VII)



(VII),

in which R^2 , A, X, Y and Z are as defined for the compound of general formula (I) according to any one of claims 1 to 3, for the preparation of a compound of general formula (I) according to any one of claims 1 to 4.

10

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2018/081545

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07D237/16 C07D237/20 A61P35/00 A61K31/501
 ADD.
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 C07D A61P A61K
 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 1 611 121 B1 (AVENTIS PHARMA SA [FR]) 8 August 2007 (2007-08-08) claims 1-15; examples 46, 98, 139-143 -----	1-13
X,P	WO 2017/202816 A1 (BAYER PHARMA AG [DE]; DEUTSCHES KREBSFORSCHUNGSZENTRUM [DE]) 30 November 2017 (2017-11-30)	12
A,P	page 45, intermediate 17; page 47, intermediate 20; page 54 intermediate 33; page 57, intermediate 37; page 62, intermediate 45; page 67, intermeiate 53; page 71, intermediate 59; claims 1-13 -----	1-11,13

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 18 December 2018	Date of mailing of the international search report 04/01/2019
--	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Kleidernigg, Oliver
--	--

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2018/081545

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
EP 1611121	B1	08-08-2007	AR 042089 A1	08-06-2005
			AR 042202 A1	15-06-2005
			AT 368651 T	15-08-2007
			AT 369356 T	15-08-2007
			AU 2003283414 A1	15-06-2004
			AU 2003296586 A1	15-06-2004
			BR 0316720 A	18-10-2005
			CA 2506022 A1	03-06-2004
			CA 2518917 A1	03-06-2004
			CN 1741999 A	01-03-2006
			CO 5700722 A2	30-11-2006
			DE 60315354 T2	08-05-2008
			DE 60315516 T2	17-04-2008
			EP 1581505 A1	05-10-2005
			EP 1611121 A1	04-01-2006
			FR 2847253 A1	21-05-2004
			JP 2006509748 A	23-03-2006
			KR 20050083918 A	26-08-2005
			MA 27814 A1	03-04-2006
			MX PA05005270 A	25-07-2005
TW 200418478 A	01-10-2004			
TW 200503727 A	01-02-2005			
WO 2004046117 A1	03-06-2004			
WO 2004046130 A1	03-06-2004			
WO 2017202816	A1	30-11-2017	AU 2017269870 A1	06-12-2018
			CA 3025227 A1	30-11-2017
			TW 201742859 A	16-12-2017
			UY 37256 A	02-01-2018
			WO 2017202816 A1	30-11-2017