

## REFERENCIAS

- Aguirre, J., Martín, J.M., Braga, J.C., Betzler, C., Berning, B. and Buckeridge, J.S. (2008). Densely-packed concentrations of acorn barnacles (Cirripedia: Sessilia) from the early Pliocene of SE Spain. *Facies*, 54: 193-206.
- Berggren, W.A., Kent, D.V., Swisher III, C.C. and Aubry, M.P. (1995). A revised Cenozoic geochronology and chronostratigraphy. In: Geochronology, Time Scales and Global Stratigraphic Correlation (W.A. Berggren, D.V. Kent, M.P. Aubry and J. Hardenbol, Editors). *Society of Economic Paleontologists and Mineralogists Special Publication*, 54: 129–212.
- Betzler, C., Brachert, T., Braga, J.C. and Martín, J.M. (1997). Nearshore, temperate, carbonate depositional systems (Lower Tortonian, Agua Amarga Basin, Southern Spain): implications for carbonate sequence stratigraphy. *Sedimentary Geology*, 113: 27-53.
- Betzler, C., Martín, J.M. and Braga, J.C. (2000). Non-tropical carbonates related to rocky submarine cliffs (Miocene, Almería, southern Spain). *Sedimentary Geology*, 131: 51-65.
- Brachert, T., Betzler, C., Braga, J.C. and Martín, J.M. (1996). Record of climatic change in neritic carbonates: turnover in biogenic associations and depositional modes (Late Miocene, southern Spain). *Geologische Rundschau*, 85: 327-337.
- Braga, J.C. and Martín, J.M. (1988). Neogene coralline-algal growth-forms and their palaeoenvironments in the Almanzora river valley (Almería, S.E. Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 67: 285-303.
- Braga, J.C. and Martín, J.M. (1992). Messinian carbonates of the Sorbas basin: sequence stratigraphy, cyclicity and facies. In: *Late Miocene Carbonate Sequences of Southern Spain: A Guidebook for the Las Negras and Sorbas area*, in conjunction with the SEPM/IAS Research Conference on Carbonate Stratigraphic Sequences: Sequence Boundaries and Associated Facies, August 30-September 3, La Seu, Spain, p. 78-108.
- Braga, J.C. and Martín, J.M. (1996). Geometries of reef advance in response to relative sea-level changes in a Messinian (uppermost Miocene) fringing reef (Cariatz reef, Sorbas Basin, SE Spain). *Sedimentary Geology*, 107: 61-81.
- Braga, J.C. and Martín, J.M. (2000). Subaqueous siliciclastic stromatolites. A case history from Late Miocene beach deposits in the Sorbas Basin of SE Spain. In: *Microbial Sediments* (R. Riding and S.W. Awramik, Editors), Springer, p. 226-232.
- Braga, J.C. y Martín, J.M. (2003). El registro sedimentario neógeno del Bloque de Cabo de Gata. En: *Itinerarios geológicos por el Terciario de la Cordillera Bética* (J. Fernández, F. García, J.M. Soria y C. Viseras, Editores). V Congreso del Grupo Español del Terciario. Guía de Excusiones. Publicaciones del Departamento de Estratigrafía y Paleontología de la Universidad de Granada, p. 53-65.

Braga, J.C., Martín, J.M. and Riding, R. (1995) Controls on microbial dome fabric development along a carbonate-siliciclastic shelf-basin transect, Miocene, SE Spain. *Palaios*, 10: 347-361

Braga, J.C., Martín, J.M. Betzler, C. and Brachert, T. (1996a). Miocene temperate carbonates in the Agua Amarga Basin (Almería, SE España). *Revista Sociedad Geológica de España*, 9 (3-4): 285-296.

Braga, J.C., Martín, J.M. and Riding, R. (1996b). Internal structure of segment reefs: *Halimeda* algal mounds in the Mediterranean Miocene. *Geology*, 24: 35-38.

Braga, J.C., Martín, J.M. and Wood, J.L. (2001). Submarine lobes and feeder channels of redeposited, temperate carbonate and mixed siliciclastic-carbonate platform deposits (Vera Basin, Almería, southern Spain). *Sedimentology*, 48: 99-116.

Braga, J.C., Baena, J., Calaforra, J.M., Coves, J.V., Dabrio, C.J., Feixas, C., Fernández-Soler, J.M., Gómez, J.A., Goy, J.L., Harvey, A.M., Martín, J.M., Martín-Penela, A., Mather, A.E., Stokes, M., Villalobos, M. y Zazo, C. (2003a). Geología del entorno árido almeriense. Guía didáctica de campo. (English version: Geology of the arid zone of Almería (South East Spain). An educational field guide) (M. Villalobos, Editor). (Tecnología de la Naturaleza SL-TECNA). *Aguas de la Cuenca del Sur*, SA (ACUSUR) – Consejería de Medio Ambiente de la Junta de Andalucía (ISBN: 84-933537-0-1), 163 pp.

Braga, J.C., Martín, J.M. and Quesada, C. (2003b). Patterns and average rates of late Neogene-Recent uplift of the Betic Cordillera, SE Spain. *Geomorphology*, 50, 3-26.

Braga, J.C., Martín, J.M., Betzler, C. and Aguirre, J. (2006). Models of temperate carbonate deposition in Neogene basins in SE Spain: a synthesis. In: *Cool-water Carbonates: Depositional Systems and Palaeoenvironmental Control* (M. Pedley and G. Carannante, Editors). *Geological Society, London, Special Publications*, 255, 121-135.

Braga, J.C., Martín, J.M., Riding, R., Aguirre, J., Sánchez-Almazo, I.M. and Dinarès-Turell, J. (2006). Testing models for the Messinian salinity crisis: the Messinian record in Almería, SE Spain. *Sedimentary Geology*, 188-189, 131-154.

Comas, M.C., Zahn, R., Klauss, A. et al. (1996). *Proceeding of the ODP Initial Reports*, 161. Ocean Drilling Program, College Station, TX.

Dabrio, C.J., Esteban, M. and Martín, J.M. (1981). The coral reef of Níjar, Messinian (Uppermost Miocene), Almería Province, S.E. Spain. *Journal Sedimentary Petrology*, 51: 521-539.

Dabrio, C.J., Martín, J.M. and Megías, A.G. (1985). The tectosedimentary evolution of Mio-Pliocene reefs in the Province of Almería. In: *6th European Regional Meeting of Sedimentologists, Excursion Guidebook* (M.D. Milá and J. Rosell, Editors), Lleida, Spain, p. 269-305.

Dronkert, H. (1977). The evaporites of the Sorbas basin. *Revista Instituto Investigaciones Geológicas Diputación Provincial y Universidad de Barcelona*, 32: 55-76.

Esteban, M., Braga, J.C., Martín, J.M. and Santisteban, C. (1996). Western Mediterranean reef complexes. In: *Models for Carbonate Stratigraphy from Miocene Reef Complexes of the Mediterranean Regions* (E. Franseen, M. Esteban, B. Ward and J.M. Rouchy, Editors). *SEPM, Concepts in Sedimentology and Paleontology Series 5*, Tulsa, Oklahoma, p. 55-72.

Haq, B.U., Hardenbol, J. and Vail, P.R. (1987). Chronology of fluctuating sea levels since the Triassic. *Science*, 235: 1156-1167.

Hilgen, F.J. (1991). Extension of the astronomically calibrated (polarity) time scale to the Miocene/Pliocene boundary. *Earth Planetary Science Letters*, 107: 349-368.

Hsü, K.J., Montadert, L., Bernoulli, D., Cita, M.B., Erickson, A., Barrison, R.E., Kidd, R.B., Melières, F., Müller, C. and Wright, R. (1977). History of the Messinian salinity crisis. *Nature*, 267: 399-403.

Jiménez, A.P. and Braga, J.C. (1993). Occurrence and taphonomy of bivalves from the Níjar reef (Messinian, Late Miocene, SE Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 102: 239-251.

Kleverlaan, K. (1987). Gordo megabed: a possible seismite in a Tortonian submarine fan, Tabernas basin, Province Almería, southeast Spain. *Sedimentary Geology*, 51: 165-180.

Kleverlaan, K. (1989). Three distinctive feeder-lobe systems within one time slice of the Tortonian Tabernas fan, SE Spain. *Sedimentology*, 36: 25-45.

Martín, J.M. and Braga, J.C. (1994). Messinian events in the Sorbas Basin in southeastern Spain and their implications in the recent history of the Mediterranean. *Sedimentary Geology*, 90: 257-268.

Martín, J.M. and Braga, J.C. (1996). Tectonic signals in the Messinian stratigraphy of the Sorbas basin (Almería, SE Spain). In: *Tertiary Basins of Spain: the stratigraphic record of crustal kinematics* (P.F. Friend and C.J. Dabrio, Editors), Cambridge University Press, Cambridge, p. 387-391.

Martín, J.M. and Braga, J.C. (1997). Sierra Nevada: historia del levantamiento de un relieve deducida de las unidades conglomeráticas de su borde. En: *Avances en el conocimiento del Terciario Ibérico* (J.P. Calvo y J. Morales, Editores). Universidad Complutense de Madrid-Museo Nacional de Ciencias Naturales, p. 117-120.

Martín, J.M. and Braga, J.C. (2001). Shallow Marine Sedimentation. In: *A Field Guide to the Neogene Sedimentary Basins of the Almería Province, South-East Spain* (A. Mather, J.M. Martín, A.M. Harvey and J.C. Braga, Editors). Blackwell, Oxford, pp.134-185.

Martín, J.M. and Braga, J.C. (2010). Arrecifes. En. *Sedimentología* (A. Arche, Editor). *Consejo Superior de Investigaciones Científicas, Servicio de Publicaciones*.

Martín, J.M., Braga, J.C. and Rivas, P. (1989). Coral successions in Upper Tortonian reefs in SE Spain. *Lethaia*, 22: 271-286.

Martín, J.M., Braga, J.C. and Riding, R. (1993). Siliciclastic stromatolites and thrombolites, late Miocene, S.E. Spain: *Journal Sedimentary Petrology*, 63: 131-139

Martín, J.M., Braga, J.C., Betzler, C. and Brachert, T. (1996). Sedimentary model and high-frequency cyclicity in a Mediterranean, shallow-shelf, temperate-carbonate environment (uppermost Miocene, Agua Amarga Basin, Southern Spain). *Sedimentology*, 43: 263-277.

Martín, J.M., Braga, J.C. and Riding, R. (1997). Late Miocene *Halimeda* alga-microbial segment reefs in the marginal Mediterranean Sorbas Basin, Spain. *Sedimentology*, 44: 441-456.

Martín, J.M., Braga, J.C. and Sánchez-Almazo, I (1999). The Messinian record of the outcropping marginal Alboran basin deposits: significance and implications. In: *Proceeding ODP, Scientific Results* (R. Zahn, M.C. Comas and A. Klaus, Editors), 161: 543-551.

Martín, J.M., Braga, J.C. and Betzler, C. (2003). Late Neogene-Recent uplift of the Cabo de Gata volcanic province, SE Spain. *Geomorphology*, 50, 27-42.

Martín, J.M., Braga, J.C., Aguirre, J. and Betzler, C. (2004). Contrasting models of temperate-carbonate sedimentation in a small Mediterranean embayment: the Pliocene Carboneras Basin, SE Spain. *Journal of the Geological Society, London*, 161, 387-399.

Mather, A. (1991). *Late Caenozoic drainage evolution of the Sorbas basin, southeast Spain*. PhD thesis, University of Liverpool, UK.

Mather, A. and Stokes, M. (2001). Marine to continental transition. In: *A Field Guide to the Neogene Sedimentary Basins of the Almería Province, South-East Spain* (A. Mather, J.M. Martín, A.M. Harvey and J.C. Braga, Editors). Blackwell, Oxford, pp. 186-224.

Mather, A. Braga, J.C. Martín, J.M. and Harvey, A.M. (2001). Introduction to the Neogene Geology of the Sorbas Basin and Transect of the Sorbas Basin. In: *A Field Guide to the Neogene Sedimentary Basins of the Almería Province, South-East Spain* (A. Mather, J.M. Martín, A.M. Harvey and J.C. Braga, Editors). Blackwell, Oxford, pp. 9-23.

Ott d'Estevou, P. and Montenat, C. (1990). Le bassin de Sorbas-Tabernas. *Documents et Travaux IGAL*, 12-13: 101-128.

Puga-Bernabéu, A., Martín, J.M. and Braga, J.C. (2007a). Tsunami-related deposits in temperate carbonate ramps, Sorbas basin, southern Spain. *Sedimentary Geology*, 199: 107-127.

Puga-Bernabéu, A., Braga, J.C. and Martín, J.M. (2007b). High-frequency cycles in upper-Miocene, ramp temperate carbonates (Sorbas basin, SE Spain). *Facies*, 53: 329-345.

Riding, R., Martín, J.M. and Braga, J.C. (1991). Coral-stromatolite reef framework, Upper Miocene, Almería, Spain. *Sedimentology*, 38: 799-818.

Riding, R., Braga, J.C., Martín, J.M. and Sánchez-Almazo, I. M.(1998). Mediterranean Messinian Salinity Crisis: constraints from a coeval marginal basin, Sorbas, SE Spain. *Marine Geology*, 146: 1-20.

Riding, R., Braga, J.C. and Martín, J.M (1999). Late Miocene Mediterranean desiccation: topography and significance of the "Salinity Crisis" erosion surface on-land in southeast Spain. *Sedimentary Geology*, 123: 1-7.

Riding, R., Braga, J.C. and Martín, J.M (2000). Late Miocene Mediterranean desiccation: topography and significance of the "Salinity Crisis" erosion surface on-land in southeast Spain: Reply. *Sedimentary Geology*, 133: 175-184.

Roep, Th.B., Beets, D.J., Dronkert, H. and Pagnier, H. (1979). A prograding coastal sequence of wave-built structures of Messinian age, Sorbas, Almería, Spain. *Sedimentary Geology*, 22: 135-163.

Roep, Th.B., Dabrio, C.J., Fortuin, A.R. and Polo, M.D. (1998). Late highstand patterns of shifting and stepping coastal barriers and washover-fans (late Messinian, Sorbas Basin, SE Spain). *Sedimentary Geology*, 116: 27-56.

Ruegg, G. J. H. (1964.). *Geologische onderzoeken in het bekken van Sorbas, Spanje*. Amsterdam Geological Institute, University of Amsterdam, Holland, 64 pp.

Sánchez-Almazo, I.M., Spiro, B., Braga, J.C. and Martín, J.M. (2001). Constraints of stable isotope signatures on the depositional palaeoenvironments of upper Miocene reef and temperate carbonates in the Sorbas Basin, SE Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 175: 153-172.

Sánchez-Almazo, I.M., Braga, J.C., Dinarès-Turell, J., Martín, J.M. and Spiro, B. (2007). Palaeoceanographic controls on reef deposition: the Messinian Cariatz reef (Sorbas basin, Almería, SE Spain). *Sedimentology*, 54, 637-660.

Sierro, F.J., Flores, J.A., Civis, J., González-Delgado, J.A. and Frances, G. (1993). Late Miocene globorotaliid event-stratigraphy and biogeography in the NE-Atlantic and Mediterranean. *Marine Micropaleontology*, 21: 143-168.

Stokes, M. (1997). *Plio-Pleistocene drainage evolution of the Vera basin, SE Spain*. PhD thesis, University of Plymouth, UK.

Weijermars, R. (1988). Neogene tectonics in the Western Mediterranean may have caused the Messinian Salinity Crisis and an associated glacial event. *Tectonophysics*, 148, 211-219.