

Originales

- » Colorimetric method for simultaneous estimation of amlodipine besylate from plasma.
Doijad RC, Sankpal PS, More HN, Pishwikar SA, Pathan AB, Suryawanshi GB.
- » Optimization of lovastatin self-nanoemulsifying solid dosage form
Patel MJ, Patel SS.
- » El extracto acuoso de *Phyllanthus orbicularis* K protege al ADN plasmídico del daño inducido por las radiaciones ultravioletas
Vernhes M, González-Pumariega M, Passaglia A, Martins FC, Sánchez-Lamar A.
- » Las funciones desempeñadas por los farmacéuticos titulares en la provincia de Valencia en 1954
Parrilla Valero F.
- » Preparation and characterization of rufinamide HP- β -cyclodextrin complexes prepared by the kneading method for solubility enhancement.
Patel Ravish J, Dave Dhara A.

Artículo Especial

- » The manufacture of gelatine capsules in the XIX century based on Aleksander Karwacki's publication dating from 1859
Rutkowska E.

Optimization of Lovastatin Self-Nanoemulsifying Solid Dosage Form.

Maulik J. Patel, Sanjay S. Patel

Department of Pharmaceutics, Shri B. M. Shah College of Pharmaceutical Education and Research, Modasa, (Gujarat, India).

Original Paper Artículo Original

Correspondence/Correspondencia:

Maulik J. Patel
Department of Pharmaceutics,
Shri B. M. Shah College of Pharmaceutical
Education and Research,
Modasa- 383 315, Gujarat, India.
Email:- maulik2121@gmail.com

Competing interest / Conflicto de intereses: No declared

Fundings / Financiación: No declared

Received: 18.04.2012
Accepted: 06.03.2013

RESUMEN

Objetivo: El objetivo del presente estudio fue desarrollar y optimizar la auto-nanoemulsión forma de dosificación sólida (SNESDF) de la Lovastatina para aumentar su solubilidad. Lovastatina (cuya solubilidad en agua es $0,4 \times 10^{-3}$ mg / ml) se considera que es un fármaco razonable debido a su alto valor de log P (4,3) y una buena solubilidad en aceites.

Materiales y Métodos: Las formulaciones fueron optimizadas por el diseño estadístico Box-Behnken en el cual las variables independientes como relación de tensioactivo: tensioactivo co-(X₁), aceite: tensioactivo co tensioactivo (X₂), y % Aerosil (X₃). Las formulaciones se caracterizan por sus variables dependientes, tales como tamaño de la gota (Y₁), la transmitancia (Y₃), el porcentaje de fármaco liberado en 5 minutos (Y₃), y dentro de 15 minutos (Y₄).

Resultados y Conclusiones: tamaño de la gota y el potencial zeta del lote optimizado resultó ser 21,89 nm y -6,4 mV, respectivamente. 44,32% y 90,78% del fármaco se encontró que se libera dentro de 5 min y 15 min, respectivamente. Por lo tanto, mediante la formulación en SNESDF de lovastatina, se encontró que la solubilidad mejoraba significativamente.

PALABRAS CLAVE: Diseño de Box-Behnken, lovastatina, Potencial Zeta, Diagrama de superficie de respuesta

ABSTRACT

Aim: The aim of present study was to develop and optimized self-nanoemulsifying solid dosage form (SNESDF) of Lovastatin for enhancing its solubility. Lovastatin (whose water solubility is 0.4×10^{-3} mg/mL) is considered to be a reasonable drug because of its high log P value (4.3) and good solubility in oils.

Materials and Methods: The formulations were optimized by Box-Behnken statistical design in which the independent variables like Ratio of surfactant: co-surfactant (X₁), oil: surfactant co surfactant (X₂), and % Aerosil (X₃). The formulations were characterized for its dependent variables such as Droplet size (Y₁), transmittance (Y₃), percentage of drug released within 5 minutes (Y₃), and within 15 minutes (Y₄).

Results and Conclusion: Droplet size and zeta potential of the optimized batch was found to be 21.89 nm and -6.4 mV, respectively. 44.32 % and 90.78 % of the drug was found to be released within 5 min and 15 min, respectively. Hence, by formulating into SNESDF, the solubility of Lovastatin was found to be significantly improved.

KEY WORDS: Box-Behnken Design, Lovastatin, Zeta Potential, Surface Response Plot

INTRODUCTION

Successful oral delivery of drug has always remained a challenge to the drug delivery field, since approximately 40% of new drugs have poor water solubility, and thus oral delivery is frequently associated with implications of low bioavailability¹. To overcome these bioavailability problems, various formulation strategies have been reported including the use of surfactants, cyclodextrin inclusion complexes, solid dispersion, nanoparticles and absorption enhancers. However the most appropriate formulation and their metabolic products are still worth to be further investigated.

Self-nanoemulsifying drug delivery systems (SNEDDS) have attracted considerable amount of interest as potential drug delivery vehicles, largely due to simplicity of preparation, clarity and ability to be filtered and incorporate wide range of drugs of varying solubility². These SNEDDS is o/w type emulsion, most suitable formulation, which is expected to increase the solubility by dissolving compounds with low water solubility into an oil phase. They can also enhance oral bioavailability by reducing the droplet size (<100 nm), and hence increase the rate of absorption due to surfactant-induced permeability changes³. Lovastatin (LOV) lowers cholesterol levels through reversible and competitive inhibition of all HMG-CoA reductase (3-hydroxy-3methylglutaryl-coenzyme A reductase), an enzyme involved in the biosynthesis of cholesterol. Lovastatin (water solubility is 0.4×10^{-3} mg/mL) is considered to be a reasonable substrate because of its high log P value (4.3) and good solubility in oils (55.10 and 83.54 mg/gm soybean oil and sunflower oil, respectively). Hence, the objective of this study was to enhance the solubility of Lovastatin by formulating self-nanoemulsifying solid dosage form (SNESDF).

Many statistical experimental designs have been recognized as useful techniques to optimize the process variables. Different types of screening designs, such as fractional factorial and Plackett- Burman screening designs have been used for preformulation evaluations⁴. Response surface methodology (RSM) is used when only a few significant factors are involved in optimization. Different types of RSM designs include 3-level factorial design, central composite design (CCD), Box Behnken design and D-optimal design⁵. A modified central composite experimental design, Box- Behnken design, is an independent, rotatable or nearly rotatable quadratic design (contains no embedded factorial or fractional factorial design), in which the treatment combinations are at the midpoints of the edges of the process space and at the center⁶. Among all the RSM designs, Box- Behnken design requires fewer runs (15 runs) in a 3-factor

experimental design. A 3-factor, 3-level design would require a total of 27 unique runs without any repetitions and a total of 30 runs with 3 repetitions. Hence, the Box- Behnken design was applied in present investigation to optimize the Lovastatin self-nanoemulsifying solid dosage forms (SNESDFs) with constraints on the release of drug within 15 min.

MATERIALS AND METHODS

Materials

Lovastatin was received as a gift sample from the Torrent Pharmaceuticals Ltd. (Ahmedabad, India). Acrysol K 140 and Aerosil 200 were received as gift samples from Corel Chemical Ltd (Ahmedabad, India). Capmul MCM and Capmul MCM C8 were kindly gifted by Abitech Corporation (USA). Tween and Span were purchased from S.D. Fine Chemicals (Mumbai, India). All other chemicals and solvents were of analytical reagent grades.

Ternary phase diagram

Ternary phase diagram were constructed to obtain the appropriate components, and their concentration ranges that resulted in a large existence area of nanoemulsion were chosen. In order to optimize the concentration of oil phase, surfactant and co-surfactant, different batches of varied concentration were prepared and titrated with distilled water till transparency persisted. Ternary phase diagram was prepared by using a constant ratio of surfactant to co-surfactant. Three ratio of surfactant (Acrysol K 140) and co-surfactant (Capmul MCM C8) were selected. (1:1, 2:1, 3:1).

Preparation of Lovastatin Self-nanoemulsifying Solid Dosage Form (SNESDF)

Box-Behnken statistical screening design was used to optimize and evaluate main effects, interaction effects, and quadratic effects of the formulation ingredients on the *in-vitro* performance of SNESDF. A 3-factor, 3-level design is suitable for exploring quadratic response surfaces and constructing second-order polynomial models. The nonlinear computer-generated Design Expert (Trial Version 8.0.4.1 STAT-EASE, Bangalore), quadratic model is given as

$$Y = b_0 + b_1x_1 + b_2x_2 + b_3x_3 + b_{11}x_1^2 + b_{22}x_2^2 + b_{33}x_3^2 + b_{12}x_1x_2 + b_{13}x_1x_3 + b_{23}x_2x_3 \dots$$

Where Y is the measured response associated with each factor level combination; b_0 is an intercept; b_1 to b_{33} are the regression coefficients; and X_1 , X_2 , and X_3 are the independent variables⁷. The selected dependent and independent variables are shown in Table 1. The Ratio of surfactant: co-surfactant (X_1), oil: surfactant co-surfactant (X_2), and % Aerosil (X_3) used to prepare each of the 15 formulations are given in Table 2. Lovastatin SNESDFs were prepared by varying the concentrations of Sunflower oil, Acrysol K140, Capmul MCM C8 and Aerosil. Predetermined amounts of the Lovastatin (10 mg) were dissolved in the required quantity of Sunflower oil in screw-capped glass vial and were warmed in a water bath at 37°C. Acrysol K140 and Capmul MCM C8 were added to the above mixture as a fixed ratio and stirred for 1 hr. All the formulations with different concentrations of surfactant, co-surfactant, oil and solid adsorbent were filled into capsules size 3 and stored at room temperature until used in subsequent studies⁸.

Characterization of Self-nanoemulsifying Solid Dosage Form

Droplet size and Zeta potential measurements

The mean droplet size and zeta potential of the resultant nanoemulsion was determined by dynamic light scattering, using a zetasizer HSA 3000 (Malvern Instruments Ltd., UK)⁹.

Transmittance study

Stability of the optimized SNESDF formulation with respect to dilution was checked by measuring transmittance at 650 nm with a UV Spectrophotometer (UV-1800, Shimadzu)¹⁰.

In-vitro Dissolution Testing


Dissolution rates from different SNESDF were determined in 900 mL of 0.1 N HCl at $37 \pm 0.5^\circ\text{C}$ with stirrer rotation speed of 50 rpm using the USP dissolution test apparatus (TDT-08L, Electrolab, Mumbai, India) employing a basket stirrer (method 1). A 10 ml aliquot of dissolution medium was withdrawn at 5, 10 and 15 min with a pipette. The samples were suitably diluted and assayed spectrophotometrically (UV-1800, Shimadzu) at 238 nm¹¹.

RESULTS AND DISCUSSION

Preparation of Self-nanoemulsifying Solid Dosage Form

The maximum amount drug was found to dissolve in sunflower oil (83.54 ± 1.98 mg/gm). Therefore, this oil was selected for nanoemulsion formulation. The required HLB value to form o/w nanoemulsion should be between 12-18 and the selection of surfactant was mainly based on this. Co-surfactant was selected based on their capability to form a stable nanoemulsion with the relevant surfactant at a minimum concentration. Capmul MCM C8 was selected

Figure 1. Ternary phase diagram of nanoemulsion (a) Acrysol K 140/ Capmul MCM C8 (2:1) and (b) Acrysol K 140/ Capmul MCM C8 (3:1).



for Acrysol K140 containing nanoemulsion. Ternary phase diagram were constructed to obtain the appropriate components and their concentration range that can result in a large nanoemulsion existence area. From the ternary phase diagrams shown in Figure 1a and 1b, it was concluded that highest nanoemulsion zone was achieved for nanoemulsion containing Acrysol K140/Capmul MCM C8 at a ratio 3:1.

Characterization of SNESDF

Droplet size and Zeta potential measurements

Droplet size for all the formulations was found < 90 nm regardless of the content used in formulation (Table 1). Lower value of the correlation coefficient (Eq. 1) clearly indicates that the response is independent of the factors studied. Once can be concluded that all these formulations resulted in acceptable droplet size range (< 100 nm) for nanoemulsions and no particular pattern was found (Table 1). As illustrated in Table 2, a p values of < 0.05 for any factor in analysis of variance (ANOVA) indicated significant effect of the corresponding factor on the response. From

Table 1. Box-Behnken Design: Independent (X) and Dependent Variables (Y)*

Batch	X ₁	X ₂	X ₃	Y ₁	Y ₂	Y ₃	Y ₄	Y ₅
	Acrysol K-140: Capmul MCM C8 (A*)	Sunflower oil: A*	% Aerosil 200	Droplet size (nm)	Transmittance (%)	% Dissolved in 5 min	% Dissolved in 15 min	Zeta potential (mV)
F1	1	1	0	40	91±0.98	44.5±1.34	89.56±2.12	-6.7
F2	1	-1	0	32	69.8±1.02	23.3±0.87	56.23±1.67	-9.5
F3	-1	1	0	30	85±2.67	37.8±2.09	82.56±0.95	-4.3
F4	-1	-1	0	45	69.45±3.21	34.23±1.23	74.78±0.56	-5.8
F5	1	0	1	31	74.23±2.56	34.55±1.89	69.89±1.09	-3.2
F6	1	0	-1	65	89.35±1.98	22.67±2.3	43.78±2.45	-7.8
F7	-1	0	1	29	78.23±2.16	39.34±2.76	75.23±1.23	-6.7
F8	-1	0	-1	32	78.56±0.87	41.76±1.56	81.23±1.45	-5.7
F9	0	1	1	26	84.56±2.35	35.23±2.32	64.34±1.67	-4.9
F10	0	1	-1	22	94.34±2.31	36.23±1.27	70.98±2.12	-6.3
F11	0	-1	1	34	76.34±2.56	29.34±2.32	53.23±1.67	-7.1
F12	0	-1	-1	89	78.89±1.45	38.34±1.27	65.34±1.34	-6.9
F13	0	0	0	20	93.23±2.13	47.34±1.56	86.23±1.54	-5.8
F14	0	0	0	21	89.99±2.24	46.31±1.67	86.23±1.54	-6.3
F15	0	0	0	19	94.21±1.23	44.67±1.23	84.29±2.34	-5.7
Variables					Levels used			
					Low	Medium	High	
Independent Variables								
X ₁ = Surfactant: Co surfactant					1:1	2:1	3:1	
X ₂ = Oil: Surfactant Co-surfactant					7:3	8:2	9:1	
X ₃ = Percentage Aerosil					15%	20%	25%	
Dependent Variables								
Y ₁ = Droplet Size (nm)							< 100	
Y ₂ = Transmittance							near to 100	
Y ₃ = Dissolution after 5 min					22.67	47.34	Maximum	
Y ₄ = Dissolution after 15 min					53.23	89.56	> 85	

A* is Acrysol K-140: Capmul MCM C8

the results of multiple regression analysis, it was found that the dependent variables, droplet size are strongly dependent on the independent variables ($P > 0.05$).

$$\text{Droplet Size (Y}_1\text{)} = 20.00 + 4.00X_1 - 10.25X_2 - 11.00X_3 + 6.63X_1X_1 + 10.13X_2X_2 + 12.62X_3X_3 + 5.75X_1X_2 - 7.75X_1X_3 + 14.75X_2X_3 \quad (R^2 = 0.8416) \quad (1)$$

Zeta potential results of all the batches were found to be -3.2 mV to -9.5 mV. Aggregation is not expected to take place, due to the slightly negative charge of the droplets.

Transmittance Study

The results of transmittance showed wide variation (Table

1). It can be inferred that these 3 factors have a profound effect on the transmittance. Formulations numbered 1, 3, 6, 9, 10, 13, 14 and 15 showed percentage transmittance of >80 %. From the results of multiple regression analysis, it was found that transmittance are strongly dependent on the independent variables ($P < 0.05$, Table 2). The correlation coefficients indicate a good fit. Polynomial equation (Eq. 2) can be used to draw a conclusion after considering the magnitude of the coefficient and the mathematical sign it carries (positive or negative). The positive sign of variables X₁ and X₂ indicated positive effect of surfactant: co-surfactant and oil: surfactant co-surfactant ratio on transmittance.

Table 2. Results of two way ANOVA for measured response.

Droplet Size					
Source	SS	DF	MS	F Ratio	P Value
Regression	4172.33	9	463.59	2.95	0.1229
Residual	785.00	5	157.00		
Total	4957.33	14			
Transmittance					
Source	SS	DF	MS	F value	P Value
Regression	1024.30	9	113.81	17.57	0.0028
Residual	32.38	5	6.48		
Total	1056.68	14			
% Dissolution after 5 min					
Source	SS	DF	MS	F value	P Value
Regression	654.03	9	72.67	2.35	0.0793
Residual	154.30	5	30.86		
Total	808.33	14			
% Dissolution after 15 min					
Source	SS	DF	MS	F value	P Value
Regression	3498.75	9	388.75	24.14	0.0013
Residual	80.54	5	16.11		
Total	3579.28	14			

DF is degree of freedom, SS is sum of square, MS is mean sum of square and F is Fischer's ratio.

$$\text{Transmittance } (Y_2) = 92.47 + 1.64X_1 + 7.55X_2 - 3.47X_3 - 8.55X_1X_1 - 5.11X_2X_2 - 3.83X_3X_3 + 1.41X_1X_2 - 3.69X_1X_3 - 1.80X_2X_3 \quad (R^2 = 0.9694) \quad (2)$$

It is clear that the ratio of oil: surfactant co-surfactant (X_2) has a more positive effect than the surfactant: co-surfactant (X_1) ratio on transmittance, this may be due to the concentration of surfactant co-surfactant was decrease the interfacial tension between oil and water interface and give transparent nanoemulsion. Same finding was true for surfactant: co-surfactant is on the transmittance.

In-vitro Dissolution Testing

From Table 1 it can be inferred that these three factors have a profound effect on the drug release profiles. Formulations numbered 1, 8, 13, 14 and 15 was >40% and only formulation 3 showed drug release of <40% drug release after 5 min. Lower value of the correlation coefficient (Eq. 3) clearly indicates that the response is independent of the factors studied. From the results of multiple regression analysis, it was found that the dependent variables, percentage dissolution after 5 min are less dependent on

the independent variables ($P < 0.05$, Table 2). For finding best optimized batch study was further extended for drug release after 15 min.

$$D_{5 \text{ min}}(Y_3) = 46.11 - 3.5X_1 + 3.57X_2 - 0.068X_3 - 5.68X_1X_1 - 5.47X_2X_2 - 5.85X_3X_3 + 4.41X_1X_2 + 3.57X_1X_3 + 2.00X_2X_3 \quad (R^2 = 0.8091) \quad (3)$$

Formulations numbered 1, 13, 14, and 15 showed higher drug release of >85% after 15 minutes of dissolution. However, the percentage of drug released after 15 min from formulations 3 and 8 was <85%. In order to obtain a formulation having rapid drug release of >85% within 15 minutes, RSM optimization was used to determine the levels of these factors. The value of the correlation coefficient (Eq. 4) clearly indicates that the response is dependent of the factors studied. The positive value of the X_2 variable represents positive effects of oil: Surfactant co-surfactant, where as variable X_1 and X_3 have negative effects on drug released after 5 min. From the results of multiple regression analysis, it was found that the dependent variables, drug released after 15 min are dependent on the independent variables ($P < 0.05$, Table 2). The correlation coefficients indicate a good fit.

$$D_{15 \text{ min}}(Y_4) = 85.58 - 2.29X_1 + 7.23X_2 - 4.33X_3 - 7.37X_1X_1 - 2.43X_2X_2 - 19.68X_3X_3 + 6.39X_1X_2 + 17.03X_1X_3 + 1.37X_2X_3 \quad (R^2 = 0.8501) \quad (4)$$

From the 3D plots, it is clear that the ratio of oil: surfactant co-surfactant (X_2) has a major effect on determining drug release within 15 min from formulations (Figures 2a, b, and c). The Figure 2a shows that at a lower ratio of Acrysol K-140: Capmul MCM C8 (A*), the percentage drug released with an increase in the ratio of sunflower oil: A* (from 54.83 % to 82.73%). However, at a higher ratio of sunflower oil: A*, the percentage drug released from formulations decreased with an increase ratio of Acrysol K-140: Capmul MCM C8 (from 81.41% to 54.83%). This finding can be explained by the fact that co-surfactant by itself does not emulsify the oil, rather the co-surfactant acts by enhancing the emulsifying capability of surfactants. Hence, optimum ratio of surfactant and co-surfactant is a key factor in achieving an emulsion. The Figure 2b shows that at a low percentage of aerosil the percentage drug released from formulations increase with a decrease in the ratio of Acrysol K-140: Capmul MCM C8 (from 57.55% to 81.10%). Figure 2a and 2b indicated that for obtaining the higher percentage drug released after 15 min the ratio of Acrysol K-140: Capmul MCM C8 (X_1) should be kept high whereas ratio of sunflower oil: A* (X_2) should be kept medium to high. Figure 2b and 2c suggested that the % aerosil (X_3) was effective at low level so should be kept medium for further study.

Figure 2. Surface response plots for the effect of (a) independent variable X_1 and X_2 on % Dissolution after 15 min (b) independent variables X_1 and X_3 on % Dissolution after 15 min (c) independent variable X_2 and X_3 on % Dissolution after 15 min




Figure 4 is a representative contour plot, which further elucidates the effects of varying ratio of Acrysol K-140: Capmul MCM C8 (A*) and Sunflower oil: A* with a fixed amount of aerosil on all responses. Figure 3 illustrates that the emulsification of Acrysol K-140: Capmul MCM C8 (A*) increases as the ratio of sunflower oil: A* is increased. Maximum transmittance and *in-vitro* drug released in 15 min were found at Acrysol K-140: Capmul MCM C8 (A*) from 0 to 1 with lower levels of Sunflower oil: A* from -0.25 to 1 as indicated by the yellow portion of the plot. Batch F1 falls in this region and therefore was selected as the best of the batches prepared according to the Box-Behnken design. Droplet size and percentage transmittance of batch F1 were found 40 nm and 91 ± 0.98 , respectively. The *in-vitro* dissolution study indicated that more than 85% of the drug released in 15 min, indicating the non interference

Figure 3. Drug release pattern of Lovastatin SNESDF formulations





Figure 4: Overlay plot in which the yellow region is the region where the constraints set on Particle size (< 19), Transmittance (> 90), % Dissolution in 5 min (>45) and in 15 min (> 80).



of the surfactant, co-surfactant, oil and aerosil with drug release. Batch F1 was selected as best optimized batch for further study.

Validation of the evolved mathematical models

To validate the evolved mathematical models, two check points were selected. Two batches CH1 and CH2 were prepared and evaluated (Table 3). As a confirmation process, a fresh formulation of Lovastatin SNCDF was prepared with Lovastatin (10 mg), Acrysol K-140: Capmul MCM C8 (A*) (2151 mg), Sunflower oil: A* (239 mg) and Aerosil (600 mg). The optimized levels of factors yielded a formulation (batch OPT) with droplet size was <30%, transmittance was >90%, rapid drug release of >40% within 5 min and of >90% within 15 min. The observed and predicted values were in very close agreement, thus strengthening the predictability of the mathematical model. Further the optimized formulation had zeta potential of -6.4 mV.

CONCLUSION

Optimization of Lovastatin SNESDF using RSM, Box-Behnken design, was performed. The ratio of independent variables, Sunflower oil, Acrysol K 140, Capmul MCM C8 and Aerosil 200 showed a significant effect on the transmittance and drug release characteristics of the formulation. The optimum ratio of these factors at 3 levels was chosen based on the quantitative effect and the polynomial equations generated by RSM. The optimized formulation prepared by using these predicted levels of factors provided desired observed responses forming nanoemulsions with more than 94% transmittance and greater than 90% drug release within 15 min.

ACKNOWLEDGEMENTS:

I am very thankful to Dr Sanjay S. Patel for his great support for this research work. We are also thankful to Manager of Abitech Corporation, USA, Corel Chemical Ltd., Ahmedabad, and Torrent Pharmaceuticals Ltd., Ahmedabad for providing us necessary ingredients. We are thankful to Shri B. M. Shah College of Pharmaceutical Education and Research, Modasa for providing technical support in form of instruments and guidance.

REFERENCES

1. Dahan A, Hoffman A. Rationalizing the selection of oral lipid based drug delivery systems by an in vitro dynamic lipolysis model for improved oral bioavailability of poorly water soluble drugs. *J Control Release*. 2008; 129(1): 1-10.
2. Nazzal S, Khan MA. Response surface methodology for the optimization of ubiquinone self-nanoemulsified drug delivery system. *AAPS PharmSciTech*. 2002; 3(1): 23-31.
3. Pouton CW. Lipid formulations for oral administration

Table 3. Validation of the evolved mathematical models

Response	Batch CH1		Batch CH2		Batch OPT	
	A= 0.4 & B= 0.6		A= 0.6 & B= 0.4		A= 0.2 & B= 0.8	
	Observed	Predicted	Observed	Predicted	Observed	Predicted
Y ₁	20.08	21.54	24.78	23.69	21.89	20.26
Y ₂	93.89	94.79	90.89	92.92	94.67	95.45
Y ₃	43.45	45.03	46.23	43.58	44.32	45.24
Y ₄	89.75	88.48	84.67	85.59	90.78	90.07

- of drugs: non-emulsifying, self-emulsifying and 'self-microemulsifying' drug delivery systems. *Eur J Pharm Sci.* 2000; 11(Suppl 2): S93-8.
4. Hamed E, Sakr A. Application of multiple response optimization technique to extended release formulations design. *J Control Release.* 2001; 73(2-3): 329-38.
 5. Box GEP, Wilson KB. On the experimental attainment of optimum multifactorial conditions. *J Roy Statist Soc.* 1951; 13: 1-15.
 6. Khan MA, Dib J, Reddy IK. Statistical optimization of Ketoprofen Eudragit S100 coprecipitates to obtain controlled-release tablets. *Drug Dev Ind Pharm.* 1996; 22(2): 135-41.
 7. Box GEP, Behnken DW. Some new three level designs for the study of quantitative variables. *Technometrics.* 1960; 2(4): 455-75.
 8. Palamakula A, Mohammad TH, Khan MA. Response surface methodology for optimization and characterization of limonene-based coenzyme Q10 self-nanoemulsified capsule dosage form. *AAPS PharmSciTech.* 2004; 5(4): 114-21.
 9. Zhang P, Liu Y, Feng N, Xu J. Preparation and evaluation of selfmicroemulsifying drug delivery system of oridonin. *Int J Pharm.* 2007; 355(1-2): 269-76.
 10. Shen H, Zhong M. Preparation and evaluation of self-microemulsifying drug delivery systems (SMEDDS) containing atorvastatin. *J Pharm Pharmacol.* 2006; 58(9): 1183-91.
 11. Kang BK, Lee JS, Chon SK, Jeong SY, Yuk SH, Khang G, et al. Development of self-microemulsifying drug delivery systems (SMEDDS) for oral bioavailability enhancement of simvastatin in beagle dogs. *Int J Pharm.* 2004; 274(1-2): 65-73.