

RESEARCH PAPER

Dual regulation of cytosolic ascorbate peroxidase (APX) by tyrosine nitration and S-nitrosylation

Juan C. Begara-Morales¹, Beatriz Sánchez-Calvo¹, Mounira Chaki¹, Raquel Valderrama¹, Capilla Mata-Pérez¹, Javier López-Jaramillo², María N. Padilla¹, Alfonso Carreras¹, Francisco J. Corpas^{3,*} and Juan B. Barroso¹

¹ Área de Bioquímica y Biología Molecular, Departamento de Biología Experimental, Facultad de Ciencias Experimentales, Ed. B3. Campus Universitario “Las Lagunillas” s/n, Universidad de Jaén, E-23071 Jaén, Spain

² Instituto de Biotecnología, Universidad de Granada, Spain

³ Departamento de Bioquímica, Biología Celular y Molecular de Plantas, Estación Experimental del Zaidín (EEZ), Consejo Superior de Investigaciones Científicas, E-18080 Granada, Spain

* To whom correspondence should be addressed. E-mail: javier.corpas@eez.csic.es

Received 4 March 2013; Revised 29 October 2013; Accepted 30 October 2013

Abstract

Post-translational modifications (PTMs) mediated by nitric oxide (NO)-derived molecules have become a new area of research, as they can modulate the function of target proteins. Proteomic data have shown that ascorbate peroxidase (APX) is one of the potential targets of PTMs mediated by NO-derived molecules. Using recombinant pea cytosolic APX, the impact of peroxyntirite (ONOO⁻) and S-nitrosoglutathione (GSNO), which are known to mediate protein nitration and S-nitrosylation processes, respectively, was analysed. While peroxyntirite inhibits APX activity, GSNO enhances its enzymatic activity. Mass spectrometric analysis of the nitrated APX enabled the determination that Tyr5 and Tyr235 were exclusively nitrated to 3-nitrotyrosine by peroxyntirite. Residue Cys32 was identified by the biotin switch method as S-nitrosylated. The location of these residues on the structure of pea APX reveals that Tyr235 is found at the bottom of the pocket where the haem group is enclosed, whereas Cys32 is at the ascorbate binding site. Pea plants grown under saline (150 mM NaCl) stress showed an enhancement of both APX activity and S-nitrosylated APX, as well as an increase of H₂O₂, NO, and S-nitrosothiol (SNO) content that can justify the induction of the APX activity. The results provide new insight into the molecular mechanism of the regulation of APX which can be both inactivated by irreversible nitration and activated by reversible S-nitrosylation.

Key words: Ascorbate peroxidase, nitration, nitric oxide, S-nitrosoglutathione, S-nitrosylation, peroxyntirite, reactive nitrogen species, salinity stress.

Introduction

The discovery of nitric oxide (NO) generation in cells and related molecules known as reactive nitrogen species (RNS) has shown that these molecules can mediate additional post-translational modifications (PTMs) such as nitration and S-nitrosylation. Protein tyrosine nitration adds a nitro group (-NO₂) to one of the two equivalent ortho-carbons of the tyrosine residue aromatic ring. This converts tyrosine into a negatively charged hydrophilic nitrotyrosine moiety and causes a marked shift in the local p*K*_a of the hydroxyl

group from 10.07 in tyrosine to 7.50 in nitrotyrosine (Turko and Murad, 2002). This PTM is regarded as a process which depends on factors such as protein structure, the nitration mechanism, and the environmental compartments where the protein is located. These covalent changes may result in effects such as loss or gain in protein function or no change in function (Souza *et al.*, 2008; Radi, 2013). S-nitrosylation consists of binding an NO group to a protein cysteine residue and can also change the function of many proteins. Through

Abbreviations: NO, nitric oxide; NO₂-Tyr, 3-nitrotyrosine; PTM, post-translational modification; ROS, reactive oxygen species; RNS, reactive nitrogen species.

© The Author 2013. Published by Oxford University Press on behalf of the Society for Experimental Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

proteomics analysis, a significant number of plant protein target candidates for *S*-nitrosylation have been identified, including the cytoskeleton, metabolic, redox-related, stress-related and signalling/regulating proteins. However, up to now, only a limited number of proteins have been studied in order to determine how they are regulated by this PTM at the molecular level (Astier *et al.*, 2012; Begara-Morales *et al.*, 2013b).

Ascorbate peroxidase (APX) together with glutathione reductase, monodehydroascorbate reductase, and dehydroascorbate reductase plus the antioxidant metabolites ascorbate, glutathione, and NADPH constitute the ascorbate–glutathione cycle. This metabolic pathway is essential for the detoxification and regulation of the cellular level of hydrogen peroxide (H_2O_2) (Asada, 1992; Noctor and Foyer, 1998). APX catalyses the electron transfer from ascorbate to H_2O_2 , thus giving rise to dehydroascorbate and water as products. This enzyme has been identified in many higher plants and comprises a family of isoenzymes with distinct characteristics which are distributed throughout the different cell compartments including the cytosol, chloroplasts, peroxisomes, and mitochondria (for a review, see Shigeoka *et al.*, 2002). In higher plants, APX is an essential element in the fine-tuning regulation mechanism of H_2O_2 during plant development and under different environmental stresses. Consequently, APX has been analysed in terms of both physiological and biochemical aspects such as catalytic regulation, enzyme–ligand interactions, molecular properties, structure, subcellular localization, gene regulation, and responses to biotic and abiotic stress (Nakano and Asada, 1981; Bunkelmann and Trelease, 1996; Corpas and Trelease, 1998; Karpinski *et al.*, 1997; Jiménez *et al.*, 1998; Yoshimura *et al.*, 1999, 2000; Rossel *et al.*, 2002; Wada *et al.*, 2003; Sharp *et al.*, 2003; Koussevitzky *et al.*, 2008).

Recently, APX has been identified as a potential target of tyrosine nitration in *Arabidopsis* (Lozano-Juste *et al.*, 2011) and *Citrus aurantium* (Tanou *et al.*, 2012), and NO has been shown to be capable of modulating its activity in different ways through either inactivation (Clark *et al.*, 2000) or activation (Keyster *et al.*, 2011; Lin *et al.*, 2011). Proteomic analysis has also identified APX as a target of *S*-nitrosylation in *Arabidopsis* plants (Fares *et al.*, 2011). With the aim of determining which mechanism(s) is(are) involved in the modulation of APX by NO-derived species, an initial pharmacological analysis using recombinant pea cytosolic APX was carried out. Furthermore, the analysis of APX under salinity stress also supports that the *S*-nitrosylation of APX contributed in the mechanism of response against the nitro-oxidative stress provoked by salinity in pea plants. Data enabled the demonstration that pea APX is modulated by both irreversible tyrosine nitration and reversible *S*-nitrosylation which lead to antagonistic effects: nitration of Tyr235 inhibits APX activity while *S*-nitrosylation of Cys32 causes an increase in APX activity, indicating an interplay between NO metabolism and a relevant antioxidant enzyme involved in ROS metabolism.

Materials and methods

Plant material and growth conditions

Pea (*Pisum sativum* L., cv. Lincoln) seeds were obtained from Royal Sluis (Enkhuizen, The Netherlands). Seeds were surface sterilized

with 3% (v/v) commercial bleaching solution for 3 min, and then were washed with distilled water, and germinated in vermiculite for 3–4 d under the following growth chamber conditions: 24 °C/18 °C (day/night), 80% relative humidity, a 16h photoperiod, and a light intensity of 190 $\mu E m^{-2} s^{-1}$. Healthy and vigorous old seedlings were selected and grown in a nutrient solutions (Corpas *et al.*, 1993). After 14 d, plants were transplanted to similar media supplemented with 150 mM NaCl and were grown for 4 d.

Crude extract of pea leaves

Leaves from control and NaCl-treated pea plants were ground in liquid nitrogen using a mortar and pestle. The resulting powder was added to 1/3 (w/v) extraction medium of 25 mM HEPES buffer, pH 8.0, containing 1 mM diethylenetriaminepentaacetic acid (DTPA) and 0.1 mM neocuproine. The crude extracts were then filtered through one layer of Miracloth (Calbiochem, San Diego, CA, USA), centrifuged at 3000 g for 6 min (4 °C), and the supernatants were used for the *S*-nitrosylated protein analysis by the biotin switch method.

Expression and purification of cytosolic pea APX

The cDNA encoding mature pea cytosolic APX (M93051) was amplified by PCR from total pea leaf RNA using the Fast Start High Fidelity polymerase (Roche) and the specific primer sets: 5'-GGATCCTATGGGAAAATCATACCCAACCTG-3' and 5'-CTC GAGTCTTAGGCTTCAGCAAATCCAAG-3'. The PCR product (766 bp) was cloned into the pGEM-T Easy Vector (Promega). The positive clones were confirmed by sequencing and then subcloned prior to digestion with *Bam*HI and *Xho*I into the pALEXb vector. Recombinant protein carrying an N-terminal choline-binding domain was produced using *Escherichia coli* strain BIVU0811, which was routinely cultured overnight at 37 °C in LB kanamycin (25 mg l^{-1}) and ampicillin (100 mg l^{-1}). Gene expression was induced by the addition of 1 mM salicylate and 10 mM 3-methyl benzoate in 250 ml of culture grown at 20 °C overnight in order to produce a higher proportion of soluble protein. Cells were harvested by centrifugation and resuspended in 20 ml of phosphate-buffered saline (PBS) (pH 7.0) containing 25 U ml^{-1} DNase I, 10 mM $MgCl_2$, and commercial protease inhibitor (Complete, Roche). Cells were lysed with a Niro Soavi NS1001L Panda High-Pressure homogenizer at a pressure of 800–900 bar. The cell lysate was then centrifuged at 10 000 g at 4 °C for 15 min, and the supernatant was used for the purification of recombinant protein using a 1 ml LYTRAP column (Biomedal). The column was washed with 20 ml of 20 mM K phosphate buffer (pH 7.0) containing 300 mM NaCl and 5 mM choline. The protein was eluted in 1 ml fractions using a discontinuous gradient of choline prepared in the same buffer with 100 mM NaCl and 20 mM choline (fraction E1), 50 mM choline (E2), 75 mM choline (E3), 100 mM choline (E4), 150 mM choline (E5), 200 mM choline (E6), 250 mM choline (E7), and 500 mM choline (E8). The samples were analysed by 10% SDS–PAGE and stained with Coomassie (Fig. 1).

Ascorbate peroxidase activity assay. Treatment with SIN-1 (peroxynitrite donor) and GSNO (nitric oxide donor)

APX (EC 1.11.1.11) activity was determined by monitoring the initial ascorbate oxidation by H_2O_2 at 290 nm (Hossain and Asada, 1984). The molecule SIN-1 (3-morpholiniosydnominine) has been shown to generate peroxynitrite, a protein-nitrating compound (Daiber *et al.*, 2004). Cytosolic recombinant APX was therefore incubated at 37 °C for 1 h with increasing concentrations (0–5 mM) of SIN-1 (Calbiochem) freshly made up before use. For treatments with NO donors, cytosolic recombinant APX was incubated at room temperature for 30 min with 0.5 mM and 2 mM GSNO. As a control, the sample was also incubated with 0.5 mM and 2 mM glutathione (GSH). The protein concentration was determined with the aid of the Bio-Rad Protein Assay using bovine serum albumin (BSA) as standard.

Identification of nitrated tyrosine in recombinant cytosolic pea APX using mass spectrometric techniques (LC-MS/MS)

Purified recombinant pea cytosolic APX was processed according to a protocol involving reduction with dithiothreitol (DTT), derivatization with iodoacetamide (IAA), and enzymatic digestion with trypsin (37 °C, 8 h). The sample was purified using solid-phase extraction cartridges to eliminate choline interference. The resulting peptide mixture was analysed using a MALDI-TOF/TOF mass spectrometer (4800, AB Sciex) to evaluate the quality of the sample. MALDI-TOF spectra were interpreted using a peptide mass fingerprinting (PMF) database search (Protein Prospector program). The database used for identification was UniProt (release 2011_02). The sample was then analysed by liquid chromatography–tandem mass spectrometry (LC-MS/MS) using a Velos-LTQ mass spectrometer equipped with a micro-ESI (electrospray ionization) ion source (ThermoFisher, San Jose, CA, USA). The sample was evaporated to dryness and diluted up to 40 µl with water containing 5% methanol and 1% formic acid. The sample was then loaded in a chromatographic system consisting of a C18 pre-concentration cartridge (Agilent Technologies, Santa Clara, CA, USA) connected to a 10 cm long, 150 µm i.d. Vydac C18 column (Vydac, IL, USA). The separation was carried out at 1 µl min⁻¹ with a 3–40% acetonitrile gradient for 30 min (solvent A, 0.1% formic acid; solvent B, acetonitrile with 0.1% formic acid). The HPLC system contained an Agilent 1200 capillary pump, a binary pump, a thermostated microinjector, and a micro switch valve.




Fig. 1. SDS-PAGE analysis of the purification of the recombinant cytosolic ascorbate peroxidase (APX). The gel was stained with Coomassie blue. M, molecular markers; I, total protein in induced culture; SF, soluble fraction; IF, insoluble fraction; FT, flow-through; W, wash; E1–E8, elution fractions.

The Velos-LTQ instrument was operated in positive ion mode with a spray voltage of 2 kV. The scan range of each full MS scan was *m/z* 400–2000. After each MS scan, a collection of targeted MS/MS spectra was obtained in order to identify both the unmodified and nitrated form of the predicted tyrosine-containing peptides. The parent mass list of the targeted scan was selected to ensure maximum coverage of the tyrosine-containing tryptic peptides for APX. The list of targeted *m/z* values was obtained after *in silico* digestion of the proteins using nitrated tyrosine as a dynamic modification. The resulting list of predicted peptides (in both nitrated and unmodified form) was filtered to exclude all peptides not containing tyrosine residues.

MS/MS spectra were searched using Proteome Discoverer software (ThermoFisher) on the basis of the following parameters: peptide mass tolerance 2 Da, fragment tolerance 0.8 Da, enzyme set as trypsin, and no missed cleavages. The dynamic modifications were cysteine carbamidomethylation (+57 Da), methionine oxidation (+16 Da), and tyrosine nitration (+45). The searches were carried out using a database containing all the proteins listed in Table 1. Identifications were filtered with XCorr >3, P(pep) <15%. The MS/MS spectra of the nitrated tyrosines were manually validated by comparing the spectra obtained for the unmodified peptide and the nitrated peptide.

Biotin switch method

For *in vitro* S-nitrosylation, recombinant APX was incubated with S-nitrosoglutathione (GSNO) for 30 min at room temperature. S-nitrosylated APX and crude extracts obtained from control and NaCl-treated pea plants were subjected to the biotin switch method as described by Jaffrey *et al.* (2001) with some slight modifications. Blocking of the non-nitrosylated free cysteine residue was carried out by incubation with 30 mM methyl methanethioniosulphonate and 2.5% SDS at 50 °C for 20 min with frequent vortexing. Residual methyl methanethioniosulphonate was removed by precipitation with 2 vols of –20 °C acetone, and the proteins were resuspended in 0.1 ml of HENS buffer (250 mM HEPES pH 7.7 buffer containing 1 mM EDTA, 0.1 mM neocuproine, and 1% SDS) per mg protein. Biotinylation was achieved by adding 1 mM N-[6-(biotinamido)hexyl]-3'-(2'-pyridyldithio) propionamide (biotin-HPDP) and 0.1 mM ascorbate, and incubating at room temperature for 1 h. Then proteins were precipitated with 2 vols of –20 °C acetone. Biotin-labelled proteins were separated by non-reducing 10% SDS-PAGE, and transferred onto polyvinylidene difluoride (PVDF) membranes (Immobilon P, Millipore, Bedford, MA, USA) using a semi-dry transfer system (Bio-Rad Laboratories). PVDF membranes were blocked with TRIS-buffered saline (TBS)+1% BSA. The blot was incubated with anti-biotin antibody at a dilution of 1:200 000 for 1 h, and immunoreactive bands were detected using a photographic film (Hyperfilm, Amersham Pharmacia Biotech) with an enhanced chemiluminescence kit (ECL-PLUS, Amersham Pharmacia Biotech).

Table 1. List of peptides scanned and peptides identified by LC-MS/MS

Peptides identified ^a	Peptides scanned	Length (amino acids)	Mr (Da)		No. of tyrosines
			Not nitrated	Nitrated	
	EQFPVSYADFYQLAGWAVEITGGPEVPFHPGR	34	3691	–	2
DVFGKAMGLSDQDIVALSGGHTIGAANKER		30	3082	–	0
SGFEGPWTSNPLIFDINSYFTELLTGEK	SGFEGPWTSNPLIFDINSYFTELLTGEK	27	3050	–	1
SYPTVSPDYQK	SYPTVSPDYQK	11	1284	1329	2
YAADEDVFFADYAEHLK	YAADEDVFFADYAEHLK	18	2074	2119	2

Some of the detected peptides do not contain tyrosines. These peptides were not included in the targeted MS/MS detection. They were detected and identified because their molecular weight coincides with that of expected peptides.

Molecular evolution studies and analysis of the structure

Molecular evolution studies were carried out at the Evolutionary Trace server (Mihalek *et al.*, 2004) using the model of the tertiary structure of the pea APX as input, and the evolutionary conservation was ranked according to the rho parameter that deviates from 1 as the variability increases. The accessible solvent area (ASA) was analysed with DSSP (Kabsch and Sander, 1983). Molecular graphics and analyses were performed with the UCSF Chimera package (Pettersen *et al.*, 2004). Alignments and phylogenetic tree were performed using Clustal W2.1 of APX amino acid sequences found in the peroxidase database (<http://peroxibase.toulouse.inra.fr/>).

Lipid peroxidation and H₂O₂ content

Lipid peroxidation was estimated by determining the concentration of malondialdehyde (MDA) with thiobarbituric acid (Buege and Aust, 1978). Hydrogen peroxide content was determined by a spectrofluorometric assay (Creissen *et al.*, 1999)

*Detection of nitric oxide (NO) and S-nitrosothiols (SNOs)**by CLSM*

NO was detected with 10 μ M 4,5-diaminofluorescein diacetate (DAF-FM DA; Calbiochem) prepared in 10 mM TRIS-HCl (pH 7.4). Leaf cross-sections were incubated at 25 °C for 1 h, in darkness, according to Corpas *et al.* (2008). After incubation, samples were washed twice in the same buffer for 15 min each. Then leaf sections were embedded in a mixture of 15% acrylamide/bisacrylamide stock solution as described elsewhere (Corpas *et al.*, 2008), and 80–100 μ m thick sections, as indicated by the vibratome scale, were cut under 10 mM PBS. Sections were then soaked in glycerol/PBS containing azide (1:1, v/v) and mounted in the same medium for examination by confocal laser scanning microscopy (CLSM; Leica TCS SL), using standard filters and collection modalities for DAF-2 green fluorescence (excitation 495 nm; emission 515 nm).

SNOs were detected using the fluorescent reagent Alexa Fluor 488 Hg-link phenylmercury (Molecular Probes, Eugene, OR, USA) according to Chaki *et al.* (2009). Briefly, leaf cross-sections of ~25 mm² were incubated at 25 °C for 1 h, in darkness, with 10 mM *N*-ethylmaleimide (NEM) prepared in ethanol, and then were washed three times in 10 mM TRIS-HCl buffer, pH 7.4, for 15 min each. Next, the leaf samples were incubated with 10 μ M Alexa Fluor 488 Hg-link phenylmercury for 1 h at 25 °C, in darkness. After washing three times in the previous buffer, leaf sections were embedded in a mixture of 15% acrylamide–bisacrylamide stock solution and were processed as described above. The pea leaf sections were analysed with a CLSM system using standard filters for Alexa Fluor 488 green fluorescence (excitation 495 nm; emission 519 nm).

Purification of biotinylated proteins and APX immunodetection

Purification of biotinylated proteins from control and NaCl-treated pea plant leaves was carried out as described by Sell *et al.* (2008) with slight modifications. Biotinylated proteins and 30 μ l of neutravidin agarose 50% (w/v) slurry (high capacity neutravidin agarose resin, Thermo Scientific) per milligram of protein were equilibrated with a neutralization buffer [10 mM HEPES pH 7.7 containing 100 mM NaCl, 1 mM EDTA, and 0.5% (v/v) Triton X-100]. Proteins were added to the neutravidin agarose matrix and were incubated 1 h at room temperature with gentle shaking. The matrix with bound proteins was washed several times with washing buffer [20 mM HEPES pH 7.7 containing 600 mM NaCl, 1 mM EDTA, and 0.5% (v/v) Triton X-100] and was transferred to an empty column. Finally, biotinylated proteins were eluted after incubation for 30 min with elution buffer (20 mM HEPES pH 7.7 containing 0.1 M NaCl, 1 mM EDTA, and 100 mM β -mercaptoethanol) at room temperature. Purified biotinylated proteins were separated by 12% SDS-PAGE and transferred to PVDF membranes as described above.

For APX immunodetection, membrane was incubated with a rabbit polyclonal antibody against cucumber APX (Corpas and Trelease, 1998) diluted 1:3000. Immunoreactive bands were detected using a photographic film (Hyperfilm, Amersham Pharmacia Biotech) with an enhanced chemiluminescence kit (ECL-PLUS, Amersham Pharmacia Biotech).

Results*Expression and purification of cytosolic APX. Effect of peroxynitrite (ONOO⁻)*

As a means to increase our knowledge of the regulation mechanism of pea APX, the recombinant protein was obtained by sequencing the pea clone and overexpression in *E. coli* (see the Materials and methods). Figure 1 shows the electrophoretic analysis of the different fractions obtained after LYTRAP affinity column chromatography of the recombinant APX. The recombinant APX showed a molecular mass of ~49 kDa which is in range of the theoretical value predicted for the cytosolic APX protein (27.7 kDa) with the Ly-tag (21.28 kDa). The E6 fraction with an APX activity of 204 nmol ascorbate min⁻¹ mg⁻¹ protein showed an adequate purity grade for this protein and it was used for the subsequent experiments.

In order to evaluate the potential action of different NO-derived molecules, an *in vitro* assay was carried out in the presence of ONOO⁻ using SIN-1 as the peroxynitrite donor (Chaki *et al.*, 2009) and GSNO as the NO donor. Figure 2A depicts the inhibitory effect of ONOO⁻ activity that ranges from 70% with 0.1 mM SIN-1 to 100% with 5 mM SIN-1. The reliability of the nitration by SIN-1 was confirmed by immunoblot analysis of the recombinant protein using an antibody against 3-nitrotyrosine. Figure 2B shows that the degree of nitration increases as a function of the SIN-1 concentration.

Effect of S-nitrosylation of recombinant pea cytosolic APX

The effect of NO on APX is controversial. It has been reported to yield an increased APX activity in sweet potato (Lin *et al.*, 2011) and soybean (Keyster *et al.*, 2011), and an inhibition in tobacco (Clark *et al.*, 2000). Pea APX contains a single cysteine residue that is partially buried (with an ASA of 15 Å²) but that may be a good candidate for *S*-nitrosylation. In order to gain additional insight into the regulation of pea APX, the effect of increasing concentrations of GSNO, a well known NO donor, on the enzymatic activity was evaluated. As shown in Fig. 2C, 0.5 mM and 2 mM GSNO significantly increase the activity of APX. When activity was assayed in the presence of 0.5 mM and 2 mM GSH to evaluate whether this effect was due to the release and binding of NO to the protein, it was found that GSH yields a reduction in APX activity (Fig. 2D) which could be consequence of a process of *S*-glutathionylation (Dalle-Donne *et al.*, 2009). Since the enzymatic assay is based on measurement of the change of absorbance at 290 nm as a consequence of the reduction of H₂O₂ with the concomitant oxidation of ascorbate, different combinations of GSNO, ascorbate, and H₂O₂ were assayed




Fig. 2. Effect of nitration (A and B) and S-nitrosylation on recombinant ascorbate peroxidase (APX) (C–E). (A) Effect of SIN-1 (peroxynitrite donor) on recombinant APX activity. (B) Representative immunoblot showing the grade of tyrosine nitration of APX treated with different concentrations of SIN-1 and detected with an antibody against 3-nitrotyrosine (dilution 1:2500). A 5 μ g aliquot of protein was used per line. (C) Effect of S-nitrosoglutathione (GSNO). (D) Effect of glutathione (GSH). (E) S-nitrosylation of recombinant pea APX. A 5 μ g aliquot of purified recombinant APX was treated with 2 mM GSH and 2 mM GSNO and was subjected to the biotin switch method.

in the absence of the enzyme to rule out any possible interference in the assay. [Supplementary Fig. S1](#) available at *JXB* online shows that none of these combinations yields significant variations in absorbance during a standard period of time used in the APX activity assay, confirming the reliability of the observed increase in activity of recombinant APX in the presence of GSNO.

Additionally, in order to confirm that GSNO treatment of recombinant APX undergoes a process of S-nitrosylation, the biotin switch method ([Jaffrey et al., 2001](#)), which is specific for the detection of S-nitrosylated proteins, was assayed. [Figure 2E](#) shows that APX is S-nitrosylated after treatment with 2 mM GSNO (lane 3), whereas the treatment with GSH does not produce any signal in the biotin switch assay (lane 2). Given the fact that the APX sequence has only one cysteine residue, Cys32 is the target of S-nitrosylation. As expected, S-nitrosylation of APX is reversible and it can be eliminated by adding a reducing agent such as DTT to the S-nitrosylated APX (lane 4) or in the absence of ascorbate (lane 5) which is used as an SNO-specific reducing agent, further demonstrating the S-nitrosylation of Cys32.

Characterization of nitrated recombinant pea cytosolic APX

With the aim of identifying which of the seven tyrosines present in the pea plant's cytosolic APX is(are) a target(s) of this post-translational modification, peroxynitrite-treated recombinant APX was subjected to trypsin digestion followed by MALDI-TOF/TOF mass spectrometry examination. [Table 1](#) shows the list of peptides scanned and those identified by LC-MS/MS. Among the peptides identified, only two contained a nitrated tyrosine. [Figure 3](#) shows the comparison of the nitrated (top) and unmodified (bottom) MS/MS spectra of these identified peptides from the pea cytosolic APX. The nitrated peptide YAAEDVFFADYAEHLK ($Z=3$) has a total of 18 amino acids and a mass of 2119 Da (2074 Da plus 45 Da) which is compatible with the acquisition of a nitro group in Tyr235 ([Fig. 3A](#)). The nitrated peptide SYPTVSPDYQK ($Z=2$) has a total of 11 amino acids and a mass of 1329 Da (1284 Da plus 45 Da) which is also compatible with the acquisition of a nitro group in Tyr5 ([Fig. 3B](#)).

Analysis of the residues involved in the APX activity regulation by peroxynitrite and GSNO

Enzyme regulation plays an important role in homeostasis and it is reasonable to assume its preservation

Control treatments were done with water (lane 1) and 2 mM GSH (lane 2). Additionally, APX was S-nitrosylated with 2 mM GSNO and reduced again with 50 mM DTT (lane 4). Furthermore, GSNO-treated APX underwent the biotin switch method without ascorbate (lane 5). Proteins were separated under non-reducing conditions by SDS-PAGE and blotted onto a PVDF membrane. Biotinylated proteins were detected using anti-biotin antibodies. Ponceau red staining demonstrated equal loading.




Fig. 3. Comparison of the nitrated (top) and unmodified (bottom) MS/MS spectra of the identified peptides from pea APX in the corresponding panels: (A) YAAEDVFFADY*AEHLK. (B) SY*PTVSPDYQK. Peptide fragment ions are indicated by 'b' if the charge is retained on the N-terminus and by 'y' if the charge is maintained on the C-terminus. The subscript indicates the number of amino acid residues in the considered fragment from either the N-terminus or the C-terminus. The superscript indicates the charge (1+ or 2+) of the backbone fragmentation.

throughout evolution. Evolutionary analysis carried out at the Evolutionary Trace server (Mihalek *et al.*, 2004) using the tertiary structure of pea APX as input (PDB code 1apx) and the entire UniProtKB database (30,342,520 sequences) output rho values of 57.68 and 4.95 for Tyr5 and Tyr235, respectively, and 14.17 for Cys32 calculated from 416 sequences. Providing that the rho parameter deviates from 1 as the variability increases, Tyr5 is poorly preserved, Cys32 does not seem to be a critical residue, and only Tyr235 seems reasonably well preserved throughout evolution. However, when the analysis was focused on the 97 sequences with E-scores ranging from 508E-143 to 423E-117 found in the *Viridiplantae* subsection of UniProtKB (1,385,397 sequences), rho values were 1 for both Cys32 and Tyr235 and 1.79 for Tyr5. This result points to Cys32 and Tyr235 as evolutionarily relevant residues in plants, as expected from their role in the modulation of APX by S-nitrosylation and tyrosine nitration revealed by the present results. In this sense, [Supplementary Fig. S2A](#) at *JXB* online shows the multiple alignments of the amino acid sequence of pea cAPX with other 15 APX sequences of three model plants (*Arabidopsis thaliana*, *Medicago truncatula*, and *Lotus japonicus*) located in the different subcellular compartments including the cytosol, peroxisome, and chloroplast (stroma and thylakoid). Thus, it is remarkable that Tyr235 is absolutely conserved and Cys32 is present in all APXs except in APX03 and APX04 of *M. truncatula*. Additionally, [Supplementary Fig. S2B](#) shows the corresponding phylogenetic tree where the APX sequences are divided into two categories: one comprising only MtAPx04 and another for the others. Interestingly, sequences are branched according to their subcellular location. The subtree for the peroxisomal APXs is subdivided into two branches: one for MtAPx03 and the other for the rest of the peroxisomal APXs. This analysis also suggests that both Tyr235 and Cys32 are important residues and that Tyr5 seems to be a general feature of cytosolic APXs. Both MtAPx03 and MtAPx04 are exceptions, but the phylogenetic tree shows that they are somehow peculiar: MtAPx04 is different from the others, regardless of the subcellular location, and MtAPx03, although being cytosolic, is also less similar.

The structure of pea APX has been solved at 2.2 Å resolution (Patterson and Poulos, 1995). It consists of a non-covalent homodimer with one haem group per monomer located in a pocket that it is opened to the exterior by two channels (Fig. 4A, B). A closer analysis shows that Tyr235 is located at the bottom of the pocket at 3.6 Å from the haem group (Fig. 4C) and Cys32 is close to the side channel (Fig. 4D). The location of Tyr5 does not reveal any functional role and since it is an accessible residue (ASA: 83 Å²) its nitration may not have a physiological relevance.

Analysis of APX activity under salt-induced oxidative stress

In order to gain additional insight into the physiological relevance of APX activity under an oxidative stress situation, it was analysed in leaves of pea plants grown in the




Fig. 4. (A) Structure of homodimeric pea APX (PDB ID: 1apx). Residues identified as a target of tyrosine nitration and S-nitrosylation are shown as space filling. (B) The haem group is enclosed in a pocket with two channels to the exterior. (C) The view along the upper channel reveals that Y235 is at the bottom of the pocket. (D) C32 is located at the ascorbate binding site in the vicinity of the side channel. (This figure is available in colour at *JXB* online.)

presence of 150 mM NaCl. Special attention was paid to H₂O₂, NO, and SNO content since they may be interconnected with the activity of APX. As shown in [Fig. 5](#), salt-induced stress yields a 12-fold increase of the content of MDA ([Fig. 5A](#)), a 2-fold increase in the content of H₂O₂ ([Fig. 5B](#)), and a 1.4-fold increase in the APX activity ([Fig. 5C](#)). By immunoblot, the APX protein expression was also evaluated, and it was found to increase under salinity conditions ([Fig. 5C](#), lanes 1 and 2). The content and localization of NO and SNOs were analysed in leaf cross-sections by CLSM using DAF-FM DA and Alexa Fluor 488 Hg-link (Chaki *et al.*, 2009) as fluorescence probes, respectively. The results revealed a significant increase in both NO and SNO production, mainly in vascular tissue, in plants grown under saline stress ([Fig. 5E, G](#)) when compared with control plants ([Fig. 5D, F](#)).

Purification of total S-nitrosylated proteins under salinity stress and detection of S-nitrosylated APX

To evaluate if APX under salinity stress conditions undergoes a process of S-nitrosylation, total S-nitrosylated proteins were purified from leaves of pea plants grown under control and salinity stress conditions and then the presence of APX




Fig. 5. Lipid peroxidation, hydrogen peroxide, and ascorbate peroxidase (APX) activity, and representative images illustrating the CLSM detection and visualization of NO and S-nitrosothiols (SNOs) in leaves of pea plants exposed to 150 mM NaCl. (A) Malondialdehyde (MDA) content. (B) Hydrogen peroxide content. (C) APX activity and western blotting analysis using an antibody against cucumber APX (dilution 1:3000). Lanes 1 and 2 correspond to leaf extracts of pea control plants and plants exposed to 150 mM NaCl, respectively. (D) Detection of NO in a leaf cross-section of pea control plants (0 mM NaCl). (E) Detection of NO in a leaf cross-section of pea plants exposed to 150 mM NaCl. (F) Detection of SNOs in a leaf cross-section of pea control plants (0 mM NaCl). (G) Detection of SNOs in a leaf cross-section of pea plants exposed to 150 mM NaCl. NO and SNOs were detected with the fluorescent dyes DAF-FM DA and Alexa Fluor (AL) 488 Hg-link reagents, respectively, as described in the Materials and methods. The chlorophyll autofluorescence is shown. Adaxial epidermis (E1), abaxial epidermis (E2), main vein (V), palisade mesophyll (Pm), and spongy mesophyll (Sm). Bar=300 μ m. Data are means \pm SEM of at least three replicates. *Differences from control values were significant at $P < 0.05$. (This figure is available in colour at *JXB* online.)




Fig. 6. Total S-nitrosylated proteins and S-nitrosylated APX in leaves of pea plants under salinity stress conditions. (A) Detection of total S-nitrosylated proteins from leaves of pea plant controls and those exposed to 150 mM NaCl. S-nitrosylated proteins were separated under non-reducing conditions by 12% SDS-PAGE and blotted onto a PVDF membrane. Biotinylated proteins were detected using anti-biotin antibodies as described in the Materials and methods. (B) Immunoblot of total S-nitrosylated proteins of leaves of pea plant controls and those exposed to 150 mM NaCl probed with a polyclonal antibody against cucumber APX (dilution 1:3,000). A 5 μ g aliquot of protein was used per lane.

protein among these S-nitrosylated protein was evaluated by immunoblots. Figure 6A depicts the electrophoretic analysis of total S-nitrosylated proteins. Thus, under salinity stress, the pattern of S-nitrosylated proteins showed an increase in the number and in the intensity of some specific bands. Figure 6B shows the immunoblot analysis of the total S-nitrosylated proteins probed with an antibody against cucumber APX where an increase under salinity stress was observed. Taken together, the results indicate that APX is S-nitrosylated *in vivo* and this process is increased under salinity conditions, which supports the data observed in *in vitro* conditions

Discussion

APX is one of the key antioxidant enzymes involved in the regulation of H₂O₂ levels during plant development and under adverse stress conditions (Jiménez *et al.*, 1998; Gomez *et al.*, 2004; Palma *et al.*, 2006; Leterrier *et al.*, 2007). Given that independent proteomic studies have identified APX as a potential target of both nitration (Lozano-Juste *et al.*, 2011) and S-nitrosylation (Bai *et al.*, 2011; Fares *et al.*, 2011), a pharmacological study using recombinant pea cytosolic APX was undertaken to identify which amino acid residue(s) is(are) potential target(s) of these PTMs and the effect(s) on APX activity.

In higher plants, proteomic analyses of nitration have shown that a certain number of proteins are targets of this PTM mediated by NO-derived molecules. However, information on the specific impact on particular proteins involved in antioxidative systems and on the consequence of tyrosine nitration on activity and protein structure is scarce (Radi, 2013). Up to now, most analyses have shown that nitration usually causes a loss of function (Corpas *et al.*, 2013). Activity loss in the case of APX thus concurs with previous observations of tobacco APX (Clark *et al.*, 2000) and those of other protein activities such as ferredoxin-NADP reductase, carbonic anhydrase (CA), S-adenosyl homocysteine hydrolase (Chaki *et al.*, 2012, 2013b), O-acetylserine(thiol)lyase A1 (Álvarez *et al.*, 2011), and NADP-isocitrate dehydrogenase (Begara-Morales *et al.*, 2013a). In this study, mass spectrometry analysis shows that two tyrosines are targets of nitration. However, of the two nitrated tyrosines, Tyr235 is the most reliable candidate to provoke the observed inhibition of the APX activity since this residue is at the bottom of the pocket where the catalytic centre is located, only 3.6 Å away from the haem group (Patterson and Poulos, 1995; Jespersen *et al.*, 1997; Mandelman *et al.*, 1998). The addition of the nitro group may disrupt the properties of the haem group to result in a loss of the activity. However, the existence of two kinetically competent binding sites for ascorbate has been reported (Mandelman *et al.*, 1998) and the facts that Tyr5 is relatively well preserved in plants (rho value 1.79) and that Tyr5 is either present or absent, but never replaced by other residues, seem to point to a physiological role for its nitration, although its location makes it difficult to rationalize the consequences on the enzymatic activity.

In higher plants, the biotin switch method has become a reliable way of detecting protein target candidates for S-nitrosylation (Sell *et al.*, 2008). Consequently, the number of identified proteins affected by this process using the biotin switch method continues to grow. However, the specific regulatory impact is largely unknown in the majority of cases (Astier *et al.*, 2012; Lin *et al.*, 2012; Tanou *et al.*, 2012), and only in a limited number of cases have their affects been clearly identified. In *Arabidopsis thaliana*, for example, cytosolic glyceraldehyde-3-phosphate dehydrogenase undergoes a reversible inhibition by NO (Lindermayr *et al.*, 2005) and catalytic Cys155 and Cys159 appear to be targets for S-nitrosylation (Holtgreffe *et al.*, 2008). More recently, it has been reported that plant NADPH oxidase (AtRBOHD) is a target of S-nitrosylation at Cys890 *in vitro* and also *in vivo* during *Pseudomonas syringae* infection. Cys890 is situated close to Phe921 which is involved in the FAD binding site (Yun *et al.*, 2011). The impact is known in other cases such as methionine adenosyltransferase which is inhibited by S-nitrosylation (Lindermayr *et al.*, 2006) and *Arabidopsis* type-II metacaspase AtMC9 which blocks the autoprocessing and activation of AtMC9 zymogen through S-nitrosylation of its catalytic cysteine residue (Belenghi *et al.*, 2007).

Using proteomic studies, APX has been identified as a potential target of S-nitrosylation (Fares *et al.*, 2011). A previous work described the inhibition of APX activity by increasing concentrations of GSNO in tobacco leaf extracts (Clark *et al.*, 2000). However more recent results showed

the opposite behaviour, with enhanced activity of different APX isozymes in root nodules of soybean treated with an NO donor (Keyster *et al.*, 2011). Similarly, in seeds of *Anticaria toxicaria* treated with NO gas, the *S*-nitrosylation of APX was described, which also enhanced its activity which it seemed to contribute during seed desiccation (Bai *et al.*, 2011). The present results showing that GSNO enhances the activity of pea APX and *S*-nitrosylation is corroborated by the biotin switch method, with Cys32 being the target of the *S*-nitrosylation. Cys32 is near the propionate side chain of the haem group and it has been reported to form thiyl radicals through interaction of APX with H₂O₂ (Kitajima *et al.*, 2008), supporting a direct reaction with NO (Martinez-Ruiz and Lamas, 2007). However, the mechanism underlying the activation by *S*-nitrosylation is far from trivial since Cys32 does not seem to be a critical residue in the catalytic process. In fact, mutation of Cys32 only provokes a 3-fold decrease on the ascorbate peroxidase activity (Mandelman *et al.*, 1998), and the structure of the soybean APX has revealed that Cys32 has no direct interaction with ascorbate, suggesting that the 1000-fold inhibition caused by DNTB [5,5'-dithiobis-(2-nitrobenzoic acid)] is due to the blockage of the side channel (Sharp *et al.*, 2003). It has also been reported that cysteine oxidation provokes loss of APX-B enzyme activity although the reason for this is not clear. Interestingly, the oxidation of Cys32 causes enzyme inactivation, and it has been suggested that glutathionylation protects the enzyme from irreversible oxidation (Kitajima *et al.*, 2008). In this context, one might hypothesize that *S*-nitrosylation prevents APX from inactivation by H₂O₂ to yield an increase of the activity when compared with unprotected enzyme.

Very recently, proteomic analysis of *Arabidopsis* roots identified the cytosolic APX (APX1) as a target of *S*-nitrosylation, and *in vitro* *S*-nitrosylation of the recombinant APX1 showed that this process provoked an increase in its activity (Correa-Aragunde *et al.*, 2013). However, the authors, using an *in silico* analysis, proposed that among the five cysteine residues present in the *Arabidopsis* APX1, Cys168 could be the target of *S*-nitrosylation.

In order to evaluate the physiological role of APX activity under stress conditions, salinity stress was selected because it has been reported to yield both oxidative and nitrosative stress (Hernández *et al.*, 1995; Gómez *et al.*, 2004; Corpas *et al.*, 2009; Leterrier *et al.*, 2012; Tanou *et al.*, 2012). Concomitant with an enhancement of the activity of APX, key elements in the metabolism of ROS and RNS, including lipid peroxidation, H₂O₂, NO, and SNOs, are significantly increased under the saline stress induced by 150 mM NaCl. The fact that *S*-nitrosylation of Cys32 causes an increase in APX activity may suggest that this PTM might be involved in the specific case of salinity stress which is accompanied by both oxidative stress and a rise in SNOs.

In summary, the present results provide new insights into the dual mechanism of regulation of APX by post-translational modification mediated by NO-derived molecules. It is interesting to note that these NO-related PTMs produce opposite effects on the enzymatic activity, with a different consequence on the long-term functionality of the proteins,

since the modulation by *S*-nitrosylation is reversible whereas tyrosine nitration leads to an irreversible inhibition of the enzyme. To the authors' knowledge, these two PTMs have not been described in relation to plant APX which is a key element in the fine-tuning regulation of H₂O₂ and consequently in the mechanism of signalling and response during plant development and/or against adverse stress conditions such as salinity. Moreover, it is shown that *S*-nitrosylation occurs *in vivo* and that this PTM is accentuated under salinity conditions as a consequence of an increase in both NO and SNOs. In these circumstances, a rise of APX activity is observed, which, in part, is due to a process of *S*-nitrosylation. This suggests that SNOs contribute to alleviate oxidative damage induced by salinity stress. Thus, the results also support the existence of an additional interplay between the metabolism of ROS and RNS in higher plants, which is in agreement with some previous data where the metabolism of both ROS and RNS under specific environmental stress conditions can cause oxidative as well as nitrosative stress (Valderrama *et al.*, 2007; Molassiotis and Fotopoulos, 2011; Corpas and Barroso, 2013).

Supplementary data

Supplementary data are available at JXB online.

Figure S1. Control of the interaction effects on A₂₉₀ of ascorbate, GSNO, and H₂O₂.

Figure S2. Multiple sequence alignments (A) and phylogenetic tree (B) of the amino acid sequence of pea cAPX and other 15 APX sequences.

Acknowledgements

JB M acknowledges a PhD fellowship (F.P.U.) from the Ministry of Science and Innovation. This work was supported by an ERDF-co-financed grant from the Ministry of Science and Innovation (BIO2009-12003-C02-01, BIO2009-12003-C02-02, and BIO2012-33904) and Junta de Andalucía (group BIO286 and BIO192), Spain. LC/MS/MS analyses were carried out at the Laboratorio de Proteómica LP-CSIC/UAB, a member of the ProteoRed network. Technical and human support provided by CICT of Universidad de Jaén (UJA, MINECO, Junta de Andalucía, FEDER) is gratefully acknowledged. We acknowledge Mr Carmelo Ruiz-Torres for his excellent technical support.

References

- Álvarez C, Lozano-Juste J, Romero LC, García I, Gotor C, León J. 2011. Inhibition of Arabidopsis O-acetylserine(thiol)lyase A1 by tyrosine nitration. *Journal of Biological Chemistry* **286**, 578–586.
- Astier J, Kulik A, Koen E, Besson-Bard A, Bourque S, Jeandroz S, Lamotte O, Wendehenne D. 2012. Protein *S*-nitrosylation: what's going on in plants? *Free Radical Biology and Medicine* **53**, 1101–10.
- Asada K. 1992. Ascorbate peroxidase: a hydrogen peroxide-scavenging enzyme in plants. *Physiologia Plantarum* **85**, 235–241.

- Bai X, Yang L, Tian M, Chen J, Shi J, Yang Y, Hu X.** 2011. Nitric oxide enhances desiccation tolerance of recalcitrant *Antiaris toxicaria* seeds via protein S-nitrosylation and carbonylation. *PLoS One* **6**, e20714.
- Begara-Morales JC, Chaki M, Sánchez-Calvo B, Mata-Pérez C, Leterrier M, Palma JM, Barroso JB, Corpas FJ.** 2013a. Protein tyrosine nitration in pea roots during development and senescence. *Journal of Experimental Botany* **64**, 1121–1134.
- Begara-Morales JC, López-Jaramillo FJ, Sánchez-Calvo B, Carreras A, Ortega-Muñoz M, Santoyo-González F, Corpas FJ, Barroso JB.** 2013b. Vinyl sulfone silica: application of an open preactivated support to the study of transnitrosylation of plant proteins by S-nitrosoglutathione. *BMC Plant Biology* **13**, 61.
- Belenghi B, Romero-Puertas MC, Vercammen D, Brackenier A, Inzé D, Delledonne M, Van Breusegem F.** 2007. Metacaspase activity of *Arabidopsis thaliana* is regulated by S-nitrosylation of a critical cysteine residue. *Journal of Biological Chemistry* **282**, 1352–1358.
- Bunkelmann JR, Trelease RN.** 1996. Ascorbate peroxidase. A prominent membrane protein in oilseed glyoxysomes. *Plant Physiology* **110**, 589–598.
- Buege JA, Aust SD.** 1978. Microsomal lipid peroxidation. *Methods in Enzymology* **52**, 302–310.
- Chaki M, Fernández-Ocaña AM, Valderrama R, Carreras A, Esteban FJ, Luque F, Gómez-Rodríguez MV, Begara-Morales JC, Corpas FJ, Barroso JB.** 2009. Involvement of reactive nitrogen and oxygen species (RNS and ROS) in sunflower–mildew interaction. *Plant and Cell Physiology* **50**, 265–279.
- Clark D, Durner J, Navarre DA, Klessig DF.** 2000. Nitric oxide inhibition of tobacco catalase and ascorbate peroxidase. *Molecular Plant-Microbe Interactions* **13**, 1380–1384.
- Corpas FJ, Barroso JB.** 2013. Nitro-oxidative stress vs oxidative or nitrosative stress in higher plants. *New Phytologist* **199**, 633–635.
- Corpas FJ, Chaki M, Fernández-Ocaña A, Valderrama R, Palma JM, Carreras A, Begara-Morales JC, Airaki M, Río LA, Barroso JB.** 2008. Metabolism of reactive nitrogen species in pea plants under abiotic stress conditions. *Plant and Cell Physiology* **49**, 1711–1722.
- Corpas FJ, Gómez M, Hernández JA, del Río LA.** 1993. Metabolism of activated oxygen in peroxisomes from two *Pisum sativum* cultivars with different sensitivity to sodium chloride. *Journal of Plant Physiology* **141**, 160–165.
- Corpas FJ, Hayashi M, Mano S, Nishimura M, Barroso JB.** 2009. Peroxisomes are required for in vivo nitric oxide accumulation in the cytosol following salinity stress of *Arabidopsis* plants. *Plant Physiology* **151**, 2083–2094.
- Corpas FJ, Palma JM, del Río LA, Barroso JB.** 2013. Protein tyrosine nitration in higher plants grown under natural and stress conditions. *Frontiers in Plant Science* **4**, 29.
- Corpas FJ, Trelease RN.** 1998. Differential expression of ascorbate peroxidase and a putative molecular chaperone in the boundary membrane of differentiating cucumber seedling peroxisomes. *Journal of Plant Physiology* **153**, 332–338.
- Correa-Aragunde N, Foresi N, Delledonne M, Lamattina L.** 2013. Auxin induces redox regulation of ascorbate peroxidase 1 activity by S-nitrosylation/denitrosylation balance resulting in changes of root growth pattern in *Arabidopsis*. *Journal of Experimental Botany* **64**, 3339–3349.
- Creissen G, Firmin J, Fryer M, et al.** 1999. Elevated glutathione biosynthetic capacity in the chloroplasts of transgenic tobacco plants paradoxically causes increased oxidative stress. *The Plant Cell* **11**, 1277–1292.
- Daiber A, Bachschmid M, Beckman JS, Munzel T, Ullrich V.** 2004. The impact of metal catalysis on protein tyrosine nitration by peroxynitrite. *Biochemical and Biophysical Research Communications* **317**, 873–881.
- Dalle-Donne I, Rossi R, Colombo G, Giustarini D, Milzani A.** 2009. Protein S-glutathionylation: a regulatory device from bacteria to humans. *Trends in Biochemical Science* **34**, 85–96.
- Fares A, Rossignol M, Peltier JB.** 2011. Proteomics investigation of endogenous S-nitrosylation in *Arabidopsis*. *Biochemical and Biophysical Research Communications* **416**, 331–336.
- Gómez JM, Jiménez A, Olmos E, Sevilla F.** 2004. Location and effects of long-term NaCl stress on superoxide dismutase and ascorbate peroxidase isoenzymes of pea (*Pisum sativum* cv. Puget) chloroplasts. *Journal of Experimental Botany* **55**, 119–130.
- Hernández JA, Olmos E, Corpas FJ, Sevilla F, del Río LA.** 1995. Salt-induced oxidative stress in chloroplasts of pea plants. *Plant Science* **105**, 151–167.
- Holtgreve S, Gohlke J, Starmann J, Druce S, Klocke S, Altmann B, Wojtera J, Lindermayr C, Scheibe R.** 2008. Regulation of plant cytosolic glyceraldehyde 3-phosphate dehydrogenase isoforms by thiol modifications. *Physiologia Plantarum* **133**, 211–228.
- Hossain MA, Asada K.** 1984. Inactivation of ascorbate peroxidase in spinach chloroplasts on dark addition of hydrogen peroxide: its protection by ascorbate. *Plant and Cell Physiology* **25**, 1285–1295.
- Jaffrey SR, Erdjument-Bromage H, Ferris CD, Tempst P, Snyder SH.** 2001. Protein S-nitrosylation: a physiological signal for neuronal nitric oxide. *Nature Cell Biology* **3**, 193–197.
- Jespersen HM, Kjaersgård IV, Ostergaard L, Welinder KG.** 1997. From sequence analysis of three novel ascorbate peroxidases from *Arabidopsis thaliana* to structure, function and evolution of seven types of ascorbate peroxidase. *Biochemical Journal* **326**, 305–310.
- Jiménez A, Hernández JA, Pastori G, del Río LA, Sevilla F.** 1998. Role of the ascorbate–glutathione cycle of mitochondria and peroxisomes in the senescence of pea leaves. *Plant Physiology* **118**, 1327–1335.
- Kabsch W, Sande C.** 1983. Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers* , **22**, 2577–2637.
- Karpinski S, Escobar C, Karpinska B, Creissen G, Mullineaux PM.** 1997. Photosynthetic electron transport regulates the expression of cytosolic ascorbate peroxidase genes in *Arabidopsis* during excess light stress. *The Plant Cell* **9**, 627–640.
- Keyster M, Klein A, Egbichi I, Jacobs A, Ludidi N.** 2011. Nitric oxide increases the enzymatic activity of three ascorbate peroxidase isoforms in soybean root nodules. *Plant Signaling and Behaviour* **6**, 956–961.
- Kitajima S, Kurioka M, Yoshimoto T, Shindo M, Kanaori K, Tajima K, Oda K.** 2008. A cysteine residue near the propionate

side chain of heme is the radical site in ascorbate peroxidase. *FEBS Journal* **275**, 470–480.

Koussevitzky S, Suzuki N, Huntington S, Armijo L, Sha W, Cortes D, Shulaev V, Mittler R. 2008. Ascorbate peroxidase 1 plays a key role in the response of *Arabidopsis thaliana* to stress combination. *Journal of Biological Chemistry* **283**, 34197–34203.

Leterrier M, Barroso JB, Valderrama R, Palma JM, Corpas FJ. 2012. NADP-dependent isocitrate dehydrogenase from *Arabidopsis* roots contributes in the mechanism of defence against the nitro-oxidative stress induced by salinity. *ScientificWorldJournal* **2012**, 694740.

Leterrier M, del Río LA, Corpas FJ. 2007. Cytosolic NADP-isocitrate dehydrogenase of pea plants: genomic clone characterization and functional analysis under abiotic stress conditions. *Free Radical Research* **41**, 191–199.

Lin A, Wang Y, Tang J, Xue P, Li C, Liu L, Hu B, Yang F, Loake GJ, Chu C. 2012. Nitric oxide and protein S-nitrosylation are integral to hydrogen peroxide-induced leaf cell death in rice. *Plant Physiology* **158**, 451–464.

Lin CC, Jih PJ, Lin HH, Lin JS, Chang LL, Shen YH, Jeng ST. 2011. Nitric oxide activates superoxide dismutase and ascorbate peroxidase to repress the cell death induced by wounding. *Plant Molecular Biology* **77**, 235–249.

Lindermayr C, Saalbach G, Bahnweg G, Durner J. 2006. Differential inhibition of *Arabidopsis* methionine adenosyltransferases by protein S-nitrosylation. *Journal of Biological Chemistry* **281**, 4285–4291.

Lindermayr C, Saalbach G, Durner J. 2005. Proteomic identification of S-nitrosylated proteins in *Arabidopsis*. *Plant Physiology* **137**, 921–930.

Lozano-Juste J, Colom-Moreno R, León J. 2011. *In vivo* protein tyrosine nitration in *Arabidopsis thaliana*. *Journal of Experimental Botany* **62**, 3501–3517.

Mandelman D, Jamal J, Poulos TL. 1998. Identification of two electron-transfer sites in ascorbate peroxidase using chemical modification, enzyme kinetics, and crystallography. *Biochemistry* **37**, 17610–17617.

Martínez-Ruiz A, Lamas S. 2007. Signalling by NO-induced protein S-nitrosylation and S-glutathionylation: convergences and divergences. *Cardiovascular Research* **75**, 220–228.

Mihalek I, Res I, Lichtarge O. 2004. A family of evolution–entropy hybrid methods for ranking protein residues by importance. *Journal of Molecular Biology* **336**, 1265–1282.

Molassiotis A, Fotopoulos V. 2011. Oxidative and nitrosative signaling in plants: two branches in the same tree? *Plant Signaling and Behavior* **6**, 210–214.

Nakano Y, Asada K. 1981. Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant and Cell Physiology* **22**, 867–880.

Noctor G, Foyer CH. 1998. Ascorbate and glutathione: keeping active oxygen under control. *Annual Review of Plant Physiology and Plant Molecular Biology* **49**, 249–279.

Palma JM, Jiménez A, Sandalio LM, Corpas FJ, Lundqvist M, Gómez M, Sevilla F, del Río LA. 2006. Antioxidative enzymes from

chloroplasts, mitochondria, and peroxisomes during leaf senescence of nodulated pea plants. *Journal of Experimental Botany* **57**, 1747–1758.

Patterson WR, Poulos TL. 1995. Crystal structure of recombinant pea cytosolic ascorbate peroxidase. *Biochemistry* **34**, 4331–4341.

Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. 2004. UCSF Chimera—a visualization system for exploratory research and analysis. *Journal of Computational Chemistry* **25**, 1605–1612.

Radi R. 2013. Protein tyrosine nitration: biochemical mechanisms and structural basis of functional effects. *Accounts of Chemical Research* **46**, 550–559.

Rossel JB, Walter PB, Hendrickson L, Chow WS, Poole A, Mullineaux PM, Pogson BJ. 2006. A mutation affecting ASCORBATE PEROXIDASE 2 gene expression reveals a link between responses to high light and drought tolerance. *Plant, Cell and Environment* **29**, 269–281.

Sell S, Lindermayr C, Durner J. 2008. Identification of S-nitrosylated proteins in plants. *Methods in Enzymology* **440**, 283–293.

Sharp KH, Mewies M, Moody PCE, Raven EL. 2003. Crystal structure of the ascorbate peroxidase–ascorbate complex. *Nature Structural Biology* **10**, 303–307.

Shigeoka S, Ishikawa T, Tamoi M, Miyagawa Y, Takeda T, Yabuta Y, Yoshimura K. 2002. Regulation and function of ascorbate peroxidase isoenzymes. *Journal of Experimental Botany* **53**, 1305–1319.

Souza JM, Peluffo G, Radi R. 2008. Protein tyrosine nitration: functional alteration or just a biomarker? *Free Radical Biology and Medicine* **45**, 357–366.

Tanou G, Filippou P, Belghazi M, Job D, Diamantidis G, Fotopoulos V, Molassiotis A. 2012. Oxidative and nitrosative-based signaling and associated post-translational modifications orchestrate the acclimation of citrus plants to salinity stress. *The Plant Journal* **72**, 585–599.

Turko IV, Murad F. 2002. Protein nitration in cardiovascular diseases. *Pharmacological Reviews* **54**, 619–634.

Valderrama R, Corpas FJ, Carreras A, Fernández-Ocaña A, Chaki M, Luque F, Gómez-Rodríguez MV, Colmenero-Varea P, del Río LA, Barroso JB. 2007. Nitrosative stress in plants. *FEBS Letters* **581**, 453–461.

Wada K, Tada T, Nakamura Y, Ishikawa T, Yabuta Y, Yoshimura K, Shigeoka S, Nishimura K. 2003. Crystal structure of chloroplastic ascorbate peroxidase from tobacco plants and structural insights into its instability. *Journal of Biochemistry* **134**, 239–244.

Yoshimura K, Yabuta Y, Ishikawa T, Shigeoka S. 2000. Expression of spinach ascorbate peroxidase isoenzymes in response to oxidative stresses. *Plant Physiology* **123**, 223–234.

Yoshimura K, Yabuta Y, Tamoi M, Ishikawa T, Shigeoka S. 1999. Alternatively spliced mRNA variants of chloroplast ascorbate peroxidase isoenzymes in spinach leaves. *Biochemical Journal* **338**, 41–48.

Yun BW, Feechan A, Yin M, et al. 2011. S-nitrosylation of NADPH oxidase regulates cell death in plant immunity. *Nature* **478**, 264–268.