


# Journal Pre-proof

Early anthropogenic change in western Mediterranean mountains (Sierra Nevada, SE Spain)

Francisca Alba-Sánchez, Daniel Abel-Schaad, José Antonio López-Sáez, Silvia Sabariego-Ruiz, Sebastián Pérez-Díaz, Reyes Luelmo-Lautenschlaeger, José Antonio Garrido-García



PII: S2213-3054(21)00001-1

DOI: <https://doi.org/10.1016/j.ancene.2021.100278>

Reference: ANCENE 100278

To appear in: *Anthropocene*

Received Date: 2 July 2020

Revised Date: 4 January 2021

Accepted Date: 6 January 2021

Please cite this article as: Alba-Sánchez F, Abel-Schaad D, López-Sáez JA, Sabariego-Ruiz S, Pérez-Díaz S, Luelmo-Lautenschlaeger R, Garrido-García JA, Early anthropogenic change in western Mediterranean mountains (Sierra Nevada, SE Spain), *Anthropocene* (2021), doi: <https://doi.org/10.1016/j.ancene.2021.100278>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Early anthropogenic change in western Mediterranean mountains (Sierra Nevada, SE Spain)

Francisca Alba-Sánchez [falba@ugr.es](mailto:falba@ugr.es)

Departamento de Botánica, Facultad de Ciencias, Universidad de Granada, Spain

Daniel Abel-Schaad [dabels@ugr.es](mailto:dabels@ugr.es) CORRESPONDING AUTHOR

Departamento de Botánica, Facultad de Ciencias, Universidad de Granada, Avenida Fuente Nueva s/n, 18071 Granada, Spain. email: [dabels@ugr.es](mailto:dabels@ugr.es)

José Antonio López-Sáez [joseantonio.lopez@cchs.csic.es](mailto:joseantonio.lopez@cchs.csic.es)

Instituto de Historia, Centro de Ciencias Humanas y Sociales, Consejo Superior de Investigaciones Científicas, Madrid, Spain

Silvia Sabariego-Ruiz [ssabarie@farm.ucm.es](mailto:ssabarie@farm.ucm.es)

Departamento de Biología Vegetal I (Botánica y Fisiología Vegetal), Universidad Complutense de Madrid, Spain

Sebastián Pérez-Díaz [sebastian.perezdiaz@unican.es](mailto:sebastian.perezdiaz@unican.es)

Department of Geography, Urban and Regional Planning, University of Cantabria, Santander, Spain

Reyes Luelmo-Lautenschlaeger [reyes.luelmo@cchs.csic.es](mailto:reyes.luelmo@cchs.csic.es)

Instituto de Historia, Centro de Ciencias Humanas y Sociales, Consejo Superior de Investigaciones Científicas, Madrid, Spain

José Antonio Garrido-García [jsntngrrdgc@gmail.com](mailto:jsntngrrdgc@gmail.com)

Estación paleontológica Valle del río Fardes – Instituto Geológico y Minero de España, Fonelas, Spain

\*Corresponding Author: Daniel Abel-Schaad, Departamento de Botánica, Facultad de Ciencias, Universidad de Granada, Avenida Fuente Nueva s/n, 18071 Granada, Spain. Email: [dabels@ugr.es](mailto:dabels@ugr.es)

#### Abstract

Human impact on western Mediterranean mountains is gaining more attention, as they harbour a rich plant diversity threatened by global change. This paper presents an analysis of two pollen records from both sides of Sierra Nevada, the prime plant-diversity centre of the Mediterranean, spanning the last 5400 years. The analysis sought to answer the following key questions: When did natural systems transform to cultural landscapes? What was the intensity and extent of alterations driven by the diverse civilizations? A

compilation of archaeological and palaeoclimatic data supported this analysis, as well as a multi-proxy palaeoecological study using pollen, non-pollen palynomorphs and microcharcoal, searching for the imprint that diverse societies left on these mountains. Results suggest that human influences started earlier than expected, with a strong impact on forest cover, oriented more towards crops on the southern face and towards livestock on the northern one, whereas mining and metallurgy were initial key drivers of population dynamics. Irrigation has supported farming since at least the Islamic period. More recent anthropogenic control and land-use management have allowed a further spread of the tree cover. Proper conservation strategies need long-term perspectives, including palaeoecological studies. The preservation of traditional human activities like grazing, high-elevation cropping or irrigation systems become essential to maintain current biodiversity.

Keywords: human impact, long-term approach, pollen, palaeoecology, landscape changes

## 1. INTRODUCTION

Mediterranean mountains are dynamic landscape systems, highly sensitive to climate change, which have hosted biotic refugia during glacial and interglacial stages (Vogiatzakis, 2012; Hughes and Woodward, 2016). The persistence until today of “biological relics” over geological times have turned these systems into key areas of genetic diversity (Taberlet and Cheddadi, 2002; Tzedakis et al., 2002; Carrión et al., 2003; Regato and Salman, 2008). A core priority is species and habitat conservation in the face of threats related to environmental change currently affecting the Mediterranean (Medail and Diadema, 2009).

Sierra Nevada (SN), a mountain range located in the south-western Mediterranean, has a special identity, harbouring a rich biodiversity of species among a unique orography (Blanca et al., 2019). A less explored feature of this mountain, however, is the complex framework of cultural landscapes shaped by humans over the last several millennia (McNeill, 2003; Gómez-Ortiz et al., 2013).

At millennial scale, climate has been one of the main agents shaping patterns of plant species distributions on SN. Numerous palaeoecological studies have highlighted the climate-induced plant dynamics (Fig. 1) on this mountain, most of them examining high-elevation deposits (e.g. Anderson et al., 2011; Jiménez-Moreno and Anderson, 2012; Ramos-Román et al., 2016). The low-elevation deposit of El Padul deserves mention as one of the longest Quaternary records (> 100 kyr) of Western Europe (e.g. Pons and Reille, 1988; Ramos-Román et al., 2018a, 2018b; Camuera et al., 2018, 2019). In addition to climate, human activity, at a millennial scale has moulded the diversity and abundance of plant taxa in the mesophilous and orophilous ecosystems of SN. This ecological diversity has provided a wide range of resources for human exploitation, giving rise to cultural landscapes since ancient times (Woodward, 2009; Mercuri and Sadori, 2014).

Early human influence as key driver shaping Mediterranean mountains has recently attracted growing interest (e.g. Smith and Zeder, 2013; Ruddiman, 2017; Roberts et al., 2019). Prehistoric human activities in Mediterranean high-mountain areas has been detected later than at low elevations, between 5500-4000 cal yr BP (López-Sáez et al., 2014; Pini et al., 2017; Cheddadi et al., 2019). At this time, humans also settled Sierra Nevada in mid- and low-elevation lands, more oriented to agriculture and grazing (Jiménez Olivencia et al., 2006), while mining was linked more to montane areas (Bertrand and Sánchez Viciano, 2008; Catalán et al. 2019).

Nevertheless, the scarcity of specific fossil records and palaeobotanical analysis conducted so far at proper elevations hinders the reconstruction of the transition from a

natural landscape in SN, shaped by climate, towards a cultural one, despite the long human presence evidenced by the extensive archaeological surveys of the area (Fig. 2). To improve the design of more realistic conservation measures in this area of high ecological value (Gillson, 2015; Catalán et al., 2019), this study addressed the following questions: (1) When did this tipping point take place in the landscape of SN, from natural toward a cultural one? (2) What was the intensity and extent of the alteration of the landscape by the diverse civilizations, regarding the successive cultural models or socioeconomic strategies?

In short, the present work sought to estimate the degree of ecological transformation in the orophilous-mesophilous landscapes of SN during the last millennia attributable to human activity (or to climate). Results thereby allowed evaluation of the imprint left by the distinct societies on this area.

To meet this aim, we conducted a multi-proxy palaeoecological study using pollen, non-pollen palynomorphs and microcharcoal. Two pollen sequences (northern and southern faces of SN) revealed the dynamics of vegetation patterns at an elevation that enabled detection of the signal of human settlements. Likewise, we compiled both archaeological and palaeoclimatic data to shed light on the feasible links between socioeconomic activities, climatic changes, and the depth of landscape transformation.




Figure 1. Location of Dehesa del Camarate and Cañar mires (white stars), showing the location of previous palynological works cited in the text (red stars): 1. Padul (Ramos-Román et al., 2018a, 2018b; Camuera et al., 2018, 2019); 2. Laguna de la Mula (Jiménez-Moreno et al., 2013); 3. Borreguiles de la Virgen (Jiménez Moreno and Anderson, 2012); 4. Laguna de Río Seco (Anderson et al., 2011); 5. Laguna de la Mosca (Manzano et al., 2019); 6. Borreguil de la Caldera (Ramos-Román et al., 2016); 7. Laguna Hondera (Mesa-Fernández et al., 2018); 8. Sierra de Gádor (Carrión et al., 2003); 9. Sierra de Baza (Carrión et al., 2007)

## 2. MATERIAL AND METHODS

## 2.1. Study area

The deposit of Dehesa del Camarate (DC: 37°10'11"N/3°15'50"W) is located at 1965 m above sea level (asl), on the northern slopes of SN, in Lugros (Granada), while the deposit of C  nar (36°57'28"N/3°25'16"W) is situated at 1593 m asl on the southern side, in C  nar (Granada; Fig. 1). Both records were the first studied in SN outside the surroundings of the highest summits of Veleta and Mulhac  n (3398 and 3479 m asl, respectively).

They form part of the central core of this range, composed of micaschists and quartzites (Blanca and L  pez, 2002). The climate varies from one point to another, with slightly higher annual rainfall in C  nar (>700 mm) despite the elevation differences, as well as higher average temperatures (*ca.* 13  C) than in DC (*ca.* 700 mm and 5  C). The whole range lies within a Mediterranean semiarid high-mountain environment, with a marked contrast between the hot-dry season and the cold-wet one (Oliva et al., 2014). SN is considered the main plant-diversity center of Mediterranean Region, with more than 100 endemic and 2000 vascular species (Blanca et al., 2019). The main species from the wet pastureland of DC are Poaceae (*Nardus stricta*, *Festuca iberica*), Cyperaceae (*Carex nigra*, *C. echinata*, *Cyperus longus*, *Eleocharis palustris*) and Juncaceae (*Juncus articulatus*, *J. effusus*). In the case of C  nar, the pastureland is surrounded by an oak (*Quercus pyrenaica*) wood and composed mainly of Cyperaceae (*Carex camposii*, *Cyperus longus*, *Scirpus holoschoenus*), Juncaceae (*Juncus effusus*, *J. conglomeratus*), Poaceae (*Holcus lanatus*, *Anthoxanthum odoratum*, *Poa trivialis*), and Fabaceae (*Lotus uliginosus*, *Trifolium pratense*).

## 2.2. Land-use patterns of human settlement

Neolithic culture in south-eastern Iberia developed from the 8th millennium cal yr BP onwards (Fig. 2). During the 7th and 6th millennia cal yr BP, the recorded pollen indicators of human activity remained weak, linked with a low grazing pressure in the surroundings of SN (L  pez-S  ez et al., 2011). The abundant available resources for farming and mining, as well as the strategic location of this area, encouraged further population growth (Cort  s S  nchez et al., 2012).

Los Millares culture characterized the Chalcolithic Period (*ca.* 5100-4250 cal yr BP). Archaeological sites occupy south-eastern Iberia, usually in low-elevation areas. These settlements were firstly oriented towards mineral deposits, easily exploitable with primitive technologies, and then associated exclusively with farming (Lull et al., 2010; Zoido Naranjo and Jim  nez Olivencia, 2015). Charcoal analyses for Los Millares site (Rodr  guez Ariza and Esquivel, 1989) provide few signals of human impact on the mountain landscape. Conversely, widespread human impact becomes more apparent in most western Mediterranean sites both at low (Mercuri and Sadori, 2014) and high altitudes (Pini et al., 2017; Catal  n et al., 2019), resulting in a significant reduction of the forest cover (Davis et al., 2015; Birks, 2019).

The Argaric culture represents the Bronze Age (*ca.* 4250-2850 cal yr BP) in the region, with a major surge in population (Lillios et al., 2016; Fyfe et al., 2019), which increased human pressure on all natural resources (Rodr  guez Ariza, 1992), especially mining, leading to a large-scale deforestation (Murillo-Barroso et al., 2017). This qualitative leap is also recorded throughout the western Mediterranean, prompting many authors to mark this period as first in which human activity became the primary driver of vegetation change (Kaplan et al., 2009; Roberts et al., 2019). Despite the decline of this culture *ca.* 3500 cal yr BP (Schirrmacher et al., 2020), the forest cover did not appear to regenerate but rather underwent a further decline, firstly with the spread of Iron Age cultures *ca.* 2850 cal yr BP (Bux  , 2008; Gil-Romera et al., 2010). Once again, the link with mining appears strong in the northern and western parts of SN, while data remain scarce for

southern slopes (Adroher Auroux et al., 2002; Martín Civantos, 2007). The few palaeobotanical studies available indicate that the forest supplied essential wood for smelting and construction (Ruíz and Rodríguez-Ariza, 2002).

The arrival of Roman culture (*ca.* 2150 cal yr BP) triggered an overall landscape transformation throughout the Mediterranean (Carrión et al., 2010; Roberts et al., 2019). In the study area, the Roman landscape heritage is linked to rural settlements. The palaeobotanical studies confirmed the widespread human transformation of landscape with settlements and open areas for pastures, together with the likely introduction of olive cultivation (Rodríguez-Ariza and Montes Moya, 2005).

Roman land-use patterns persisted for centuries (Salvador Ventura, 2012) until the Islamic Period beginning in the early AD 8th century, when a major landscape change occurred on SN, especially concerning irrigation and terracing, as well as the diversification of human activities (Salvador Oyonate, 2013; Martín Civantos, 2014). The transition from the medieval to the Modern Period is marked in the area by the repopulation impelled by the Christian Kingdom during AD 16<sup>th</sup>-17<sup>th</sup> centuries (Sayadi and Calatrava, 2001; Trillo San José, 2004). Population grew steadily in the following centuries, causing a further intensification in the use of natural resources, but also the establishment of different forest-protection measures (Mesa Garrido, 2016), which culminated in the declaration of the National Park of Sierra Nevada in AD 1999.




Figure 2. Pre-Roman settlements in the surrounding study area. 1. Barranco León; 2. Cueva Horá; 3. Fuente Nueva; 4. Las Angosturas; 5. Cerro del Castillo; 6. Cerro de San Cristóbal; 7. Peñón de la Reina; 8. Las Majolicas; 9. Cerro de la Virgen; 10. Dólmenes de Gorafe; 11. El Malagón; 12. Cueva Romero; 13. Necrópolis de Gádor; 14. Los Millares; 15. Loma de la Galera; 16. Peñón de las Juntas; 17. Cueva de Nieves; 18. Castellon Alto; 19. Cerro de la Encina; 20. Cuesta del Negro; 21. La Terrera del Reloj; 22. El Cardal; 23. Juan Canal; 24. Peñón de Arruta; 25. Cuesta de los Chinos; 26. La Quinta; 27. Barranco Tio Gabriel; 28. Cuevas de la Zubia; 29. Loma de la Balunca; 30. Cerro Cepero; 31. Tútugi; 32. Acci; 33. El Forruchu; 34. Cerro de la Calera; 35. Iliberri; 36. Campo de Marte; 37. Cueva de Montealegre;

### 2.3. Coring and Chronology

We extracted a core of 112 cm with a Russian peat corer in DC and divided into samples 2 cm thick for subsequent analyses. Five radiocarbon dating were performed at Poznan Radiocarbon Laboratory (Table 1). We used a similar method to draw a 64-cm core from the mire of Cáñar. Four radiocarbon datings were carried out at the same laboratory (Table 2). We calibrated ages BP (Stuiver et al., 2019) with confidence intervals calculated at 95% ( $2\sigma$ ). We then constructed age-depth models (Fig. 3) applying a smooth-spline solution with Clam 2.2 software (Blaauw, 2010).




Figure 3. Age-depth models for Dehesa del Camarate and Cáñar mires

Depth (cm)	Lab Code	Age $^{14}\text{C}$	Age cal yrBP		Probability (%)
			Mín	Max	
35	Poz-64946	$780 \pm 30$ BP	670	738	95.0
63	Poz-62302	$1940 \pm 30$ BP	1822 1965	1949 1965	94.7 0.2
83	Poz-64947	$3340 \pm 30$ BP	3480 3544 3671	3538 3640 3678	23.1 70.3 1.4
100	Poz-64948	$3750 \pm 30$ BP	3988 4067 4168 4199	4047 4162 4179 4228	20.8 64.7 1.9 7.5
109	Poz-68674	$4525 \pm 35$ BP	5049 5212	5194 5308	61.5 33.4

Table 1. Results of Radiocarbon ( $^{14}\text{C}$ ) dating of core samples from Dehesa del Camarate, showing calibrated age ranges ( $2\sigma$ ).

Depth (cm)	Lab Code	Age $^{14}\text{C}$	Age cal yrBP		Probability (%)
			Mín	Max	
16	Poz-92960	$260 \pm 30$ BP	151	171	12.2

			280	331	56.7
			357	431	29.7
32	Poz-92961	745 ± 30 BP	662	726	100.0
48	Poz-92962	1220 ± 30 BP	1063	1188	78.3
			1203	1258	21.7
62	Poz-64990	2040 ± 30 BP	1903	1908	0.8
			1923	2068	91.8
			2079	2111	7.4

Table 2. Results of Radiocarbon ( $^{14}\text{C}$ ) dating of core samples from Cáñar, showing calibrated age ranges ( $2\sigma$ )

#### 2.4. Pollen, non-pollen palynomorphs (NPPs) and microcharcoal analyses

Sample treatment for pollen analysis followed the methodology of Faegri and Iversen (1989), using Thoulet solution for the densimetric separation of pollen and NPPs, and adding a *Lycopodium* tablet to each sample in order to estimate the pollen concentration. We used TILIA to process the data and graphics (Grimm, 2004).

We identified pollen and NPPs with the help of several keys and atlases (e.g. Reille, 1999; Van Geel, 2001) whereas nomenclature followed Miola (2012). We counted microcharcoals on the same pollen slides (Mooney and Tinner, 2011) and classified them into 2 categories ( $>100$  and  $<100$   $\mu\text{m}$ ) to indicate local or regional fires, respectively. A synthetic pollen diagram of each deposit was included (Figs. 4 and 5), where a number of pollen types and NPPs were merged according to their ecological affinities (Table 3), as well as charts showing the Total Pollen Concentration and the Charcoal Accumulation Rates (CHAR). This latter chart was determined by dividing the concentration of microcharcoals by the sedimentation rate of each sample obtained from the age-depth model.

Group	Pollen and NPPs
Riparian woods	<i>Alnus</i> , <i>Betula</i> , <i>Corylus</i> , <i>Fraxinus</i> , <i>Juglans</i> , <i>Prunus</i> , <i>Salix</i> , <i>Ulmus</i>
Crops	<i>Cerealia</i> , <i>Secale cereale</i> , <i>Vicia faba</i> , <i>Linum</i>
Anthropogenic Perennial Pastures (APP)	<i>Allium</i> , Apiaceae, <i>Artemisia</i> , Brassicaceae, <i>Campanula</i> , Caryophyllaceae, Fabaceae, Liliaceae, <i>Ranunculus</i> , Rosaceae, Scrophulariaceae, <i>Viola</i>
Anthropogenic Nitrophilous Communities (ANC)	<i>Anthemis</i> , <i>Aster</i> , Boraginaceae, Cardueae, <i>Centaurea cyanus</i> , <i>C. nigra</i> , Cichorioideae, <i>Erodium</i> , <i>Galium</i> , <i>Geranium</i> , <i>Malva sylvestris</i>
Anthropozoogenic Nitrophilous Communities (AZNC)	Chenopodiaceae, <i>Plantago</i> spp., <i>Rumex</i> spp., <i>Urtica dioica</i>
Coprophilous fungi	<i>Gelasinospora</i> sp. (HdV 1), <i>Gelasinospora</i> cf. <i>retispora</i> (HdV 2), <i>Chaetomium</i> sp. (HdV 7A), <i>Sordaria</i> -type (HdV 55), <i>Cercophora</i> sp. (HdV 112), <i>Sporormiella</i> sp. (HdV 113), <i>Coniochaeta</i> cf. <i>ligniaria</i> (HdV 172), <i>Podospora</i> -type (HdV 368)
Erosive phases	<i>Glomus</i> , <i>Pseudoschizaea circula</i> , <i>Entorrhiza</i> (HdV 527)
Dry conditions	<i>Pleospora</i> sp. (HdV 3B), HdV 16C, HdV 63C
Wet conditions	HdV 18

Table 3. Groups of pollen and non-pollen palynomorphs according to their ecological affinities.

### 3. RESULTS

The sequence of DC starts ca. 5410 cal yr BP, while the one of Cáñar begins ca. 2080 cal yr BP. Both of them were divided into four distinct subzones (Tables 4 and 5; Figs. 4 and 5). Likewise, main results are summarized and compared with other proxies and studies in Figure 6.






Figure 4. Summary diagram of Dehesa del Camarate (DC), including selected pollen types, NPPs, CHAR and Total Pollen Concentration. APP: Anthropogenic Perennial Pastures; ANC: Anthropogenous Nitrophilous Communities; AZNC: Anthropozoogenous Nitrophilous Communities; TR: Trees (= Arboreal Pollen –AP-); SHR: Shrubs; HER: Herbs; Stars indicate Radiocarbon datings.




Figure 5. Summary diagram of Cãñar, including selected pollen types, NPPs, CHAR and Total Pollen Concentration. APP: Anthropogenic Perennial Pastures; ANC: Anthropogenous Nitrophilous Communities; AZNC: Anthropozoogenous Nitrophilous Communities; TR: Trees (= Arboreal Pollen – AP-); SHR: Shrubs; HER: Herbs; Stars indicate Radiocarbon datings.

SUBZONE Depth (cm)/ Age (cal BP)	Trees/Shrubs	Herbs	NPPs	Pollen Concentration/ CHAR
CAMA1 112-101 cm ca. 5410-4300 cal yr BP/ 3460-2350 cal yr BC	AP (TR in Figs. 4 and 5): 7-12,8% <i>Pinus sylvestris</i> (5,9-1,7%), <i>Quercus pyrenaica</i> (1-2%), <i>Q. ilex</i> (1-2%)  Riparian woods (>1%) Sporadic occurrences of <i>Abies</i> and <i>Cedrus</i> <i>Cytisus</i> (1-2%), <i>Cistus</i> (0-2%), <i>Erica arborea</i> and <i>Helianthemum</i> (<1%)	Poaceae (51-58%) APP (13,5-21,8%) ANC (7,2-9,4%) AZNC (2,4-4%) Sporadic <i>Cerealia</i> <i>Cyperaceae</i> (4-1,3%)	Coprophilous fungi only ca. 5000 cal yr BP Low levels of Erosive and Dry phases indicators	Lowest Pollen concentration Low levels of CHAR, higher of bigger ones

CAMA2 101-67 cm ca. 4300-2250 cal yr BP/ 2350-300 cal yr BC	AP: 5-8,4% <i>Pinus sylvestris</i> (2,4-0,2%), <i>Quercus pyrenaica</i> (1-4%), <i>Q. ilex</i> (1-2%) Continuous curve of <i>Cedrus</i> from 3800 to 3300 cal yr BP <1% Riparian woods (~1%) <i>Cytisus</i> (1-2,6%), <i>Cistus</i> (0,5-3%), <i>Erica</i> (0-2,4%)	Poaceae (53-64%) APP (20,5-13,1%) ANC (12,7-7,8%) AZNC (2-6,8%) Continuous curve of <i>Cerealia</i> (<1%), <i>Vicia faba</i> appears sporadically Cyperaceae (2,4-3,6%)	Coprophilous fungi nearly absent Increase of Erosive phases indicators Alternative low levels of Dry and Wet phases indicators	Slight increase of Pollen concentration Low levels of CHAR <100 and medium ones of CHAR >100 $\mu\text{m}$
CAMB1 67-23 cm ca. 2250-410 cal yr BP/ 300 cal yr BC-1540 cal yr AD	AP: 4-9,5% <i>Pinus sylvestris</i> (0-2%), <i>Quercus pyrenaica</i> (3,9-2%), <i>Q. ilex</i> (2-3%) Continuous curve of <i>Castanea</i> ~1400 cal yr BP and <i>Olea</i> from ~1050 cal yr BP Sporadic occurrences of <i>Cedrus</i> Riparian woods (~1%) <i>Cytisus</i> (1-3,6%), <i>Cistus</i> (0,2-1,5%), <i>Erica</i> (0-2%), <i>Helianthemum</i> (1%)	Poaceae (62,7-54%) APP (16,7-12,3%) ANC (6,3-14,1%) AZNC (4,1-5,2%) Almost continuous curve of <i>Cerealia</i> (<1%), <i>Vicia faba</i> appears sporadically Maxima of Cyperaceae (8,3%)	Coprophilous fungi increase initially Significant increases of Wet and Erosive phases indicators ~1135-860 cal yr BP Dry phases indicators nearly absent	Initial increase and final decrease of Total Pollen Concentration Higher values of CHAR
CAMB2 23-0 cm ca. 410 cal yr BP/1540 cal yr AD-present	AP: 9,7-21,1% <i>Olea</i> (1-10,4%), <i>Pinus sylvestris</i> (2-4,1%), <i>Quercus pyrenaica</i> (2-3%), <i>Q. ilex</i> (2-4%) Riparian woods (1-2%) <i>Cytisus</i> (3,2-2,7%), <i>Cistus</i> (<1%), <i>Erica</i> (<1%), <i>Helianthemum</i> (0,6-1,5%)	Poaceae (43,2-42,8%) APP (15,1-13,4%) ANC (20-12%) AZNC (6,2-4,3%) <i>Cerealia</i> (<1%), sporadic <i>Vicia faba</i> Cyperaceae (0,2-3,7%)	Maxima of Coprophilous fungi and Erosive phases indicators High levels of Wet phases indicators and lower of Dry phases indicators	Final drop of Pollen Concentration Maxima of CHAR

Table 4. Description of pollen zones from Dehesa del Camarate (DC). AP: Arboreal Pollen = TR in Figs. 4 and 5; APP: Anthropogenic Perennial Pastures; ANC: Anthropogenous Nitrophilous Communities; AZNC: Anthropozoogenous Nitrophilous Communities; CHAR: Charcoal Accumulation Rates.

SUBZONE Depth (cm)/ Age (cal BP)	Trees/Shrubs	Herbs	NPPs	Pollen Concentration/ CHAR
CAÑA1 62-56 cm ca. 2000-1560 cal yr BP/ 50 cal yr BC- 390 cal yr AD	AP (TR in Figs. 4 and 5): 2-3,3% <i>Pinus pinea/halepensis</i> , <i>Quercus pyrenaica</i> and <i>Q. ilex</i> below 1% respectively; Riparian woods (~1%) <i>Cytisus</i> (3-6,6%), <i>Cistus</i> , <i>Erica</i> and <i>Helianthemum</i> (<1%) respectively	Poaceae (41,7-49,3%) APP (13,1-14,4%) ANC (14,1-22,1%) AZNC (0,4-2,3%) <i>Cerealia</i> (8-5%), <i>Secale</i> (8-4%) Cyperaceae (1,7-3,5%)	Coprophilous fungi, Dry and Wet phases indicators almost absent Medium levels of Erosive phases indicators	Medium Pollen concentration Low levels of CHAR
CAÑA2 56-44 cm ca. 1560-1000 cal yr BP/ 390-950 cal yr AD	AP: 1,8-3,8% <i>Pinus pinea/halepensis</i> , <i>Quercus pyrenaica</i> and <i>Q. ilex</i> below 1% respectively; Riparian woods (~1%) <i>Cytisus</i> (1,9-3,4%), <i>Cistus</i> , <i>Erica</i> and <i>Helianthemum</i> (<1%) respectively	Poaceae (42,2-59,3%) APP (12,5-16,7%) ANC (19,3-12,9%) AZNC (<1%) <i>Cerealia</i> (10-3%), <i>Secale</i> (9-6%) Cyperaceae (~1%)	Final slight increase of Coprophilous fungi Erosive phases indicators decrease, while Dry phases indicators prevail over Wet phases ones	Increase of Pollen concentration Maximum of CHAR <100 and low ones of CHAR >100 $\mu\text{m}$
CAÑB1 44-16 cm ca. 1000-335 cal yr BP/ 950- 1590 cal yr AD	AP: 2,6-10,2% <i>Quercus pyrenaica</i> (0,5-6,1%), <i>Q. ilex</i> (0,7-1,8%), <i>Pinus pinea/halepensis</i> (0,5-2%), <i>Olea europaea</i> (<1%) Riparian woods almost disappear <i>Cytisus</i> (4,2-2%), <i>Cistus</i> , <i>Erica</i> and <i>Helianthemum</i> (<1%) respectively	Poaceae (58,8-47%) APP (15,5-11,1%) ANC (10,4-13,3%) AZNC (>1%) <i>Cerealia</i> (2,2-0,5%), <i>Secale</i> (4,6-14%) Cyperaceae (0,7-5,6%)	Coprophilous fungi final maximum Dry, Wet and Erosive phases indicators keep their levels	Increase of Pollen Concentration Maxima of CHAR
CAÑB2 16-0 cm	AP: 10,8-28,1% <i>Quercus pyrenaica</i> (6,4-15,9%), <i>Pinus pinea/halepensis</i> (0,4-3,1%), <i>Olea</i>	Poaceae (57,4-51%) APP (9,4-9,9%) ANC (13,4-3,5%) AZNC (2,8-3,3%)	High levels of Coprophilous fungi High levels of Wet phases indicators and lower of Dry	Final maximum of Pollen Concentration

ca. 335 cal yr BP/1590 cal yr AD-present	<i>europaea</i> (0,4-2,3%), <i>Castanea</i> , <i>Q. ilex</i> and Riparian Woods (>1%) respectively <i>Cytisus</i> (1,3-0,7%), <i>Cistus</i> (<1%)	Cerealia (<2%), <i>Secale</i> (<1%) Cyperaceae (0,2-3,7%)	and Erosive phases indicators	Minor but significant levels of CHAR
------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------	-------------------------------	--------------------------------------

Table 5. Description of pollen zones from Cañar. AP: Arboreal Pollen = TR in Figs. 4 and 5; APP: Anthropogenic Perennial Pastures; ANC: Anthropogenous Nitrophilous Communities; AZNC: Anthropozoogenous Nitrophilous Communities; CHAR: Charcoal Accumulation Rates.

#### 4. DISCUSSION

4.1. The last remaining ancient forests during the Chalcolithic (*ca.* 5410-4300 cal yr BP) The deepest part of DC (CAMA1) core corresponds to the transition from Middle to Late Holocene (Walker et al., 2012), a period climatically defined by an aridification process throughout the western Mediterranean (Jalut et al., 2009; Pérez-Obiol et al., 2011). This aridity trend (Fig. 6) has also been detected in SN and nearby ranges (e.g. Anderson et al., 2011; Oliva et al., 2014), starting *ca.* 5700 cal yr BP, and could be behind the origin of this deposit (García-Alix et al., 2012).

This pollen diagram (Fig. 4) shows a remarkably low percentage (*ca.* 13%) of arboreal pollen (AP) in this subzone (Table 4) compared to previous high-mountain fossil records from SN (Fig. 1), where it is represented mainly by *Pinus* and consistently over 20%. Furthermore, all of them are located at significantly higher elevations (*ca.* 2500-3000 m asl), further from both the timberline and human signal during that period. This anomaly might be attributed to higher aridity in DC, as rainfall decreases eastwards in SN and temperatures also fall along the northern slopes (Olivares et al., 2011), both limiting forest development (Oliva et al., 2014). Nevertheless, the forest decline detected in DC could be rather ascribed to human impact, which probably started earlier on this mountain side, as shown by the presence of significant levels of Anthropogenous Nitrophilous Communities (ANC) and minor ones of Anthropozoogenous Nitrophilous Communities (AZNC) and microcharcoals, as well as the sporadic appearances of coprophilous fungi and pollen of *Cerealia*. It would have given rise to an open landscape where scrubland and mostly grasslands became a predominant part, after the depletion of the last remaining ancient pine forests. This latter demise occurred at *ca.* 5200 cal yr BP in Laguna de Río Seco (Anderson et al., 2011) and somewhat later (*ca.* 4300-4000 cal yr BP) in the other records. This delayed response could be related to the cited higher location of these deposits.

At that time, human populations expanded as a result of advances in agriculture and farming (Martín Socas et al., 1999; Carrasco Rus et al., 2016). From Neolithic to Chalcolithic human traces (Fig. 2) spread not only to coastal zones but also to montane areas of Southern Iberia (Rodríguez Ariza, 2000; Nocete, 2006).

The Chalcolithic (*ca.* 5300-4250 cal yr BP), represented mainly by the Los Millares copper culture (Lull et al., 2010; García San Juan and Murillo Barroso, 2013), registered a further rise after *ca.* 4800 cal yr BP (Schirmacher et al., 2020), which matched the retreat of pine forests as well as the parallel minor signs of human activities near the study area.

Beyond the dynamics of major forest taxa, the pollen diagram records the sporadic presence of some noteworthy types, such as *Betula* or *Cedrus*. Although these underwent different dynamics throughout the Holocene, their appearance in the pollen diagram remained under 1%. The DC pollen diagram shows that the *Betula* populations were no larger than the current ones for the last 5000 years.

*Cedrus* presence in several pollen diagrams of south-eastern Spain has long been attributed to the Saharian winds (Magri, 2012), mainly in arid periods (see details on

Alba-Sánchez et al., 2018). However, its appearance in the present study and other works for the SN (Anderson et al., 2011; Jiménez-Moreno et al., 2013; Manzano et al., 2019) does not correspond (Fig. 6) to the periods with the highest incidence of African winds (Jiménez-Espejo et al., 2014), a finding that supports the persistence of small, isolated pockets of *Cedrus* in SN during the Holocene, which would have gone extinct as a result of human impact (Postigo-Mijarra et al., 2010). Nevertheless, further studies are needed to clarify this controversy.

#### 4.2. Forest demise and increasing human pressure during Bronze and Iron Ages (*ca.* 4300-2250 cal yr BP)

Conditions in the study area remained arid and even became drier as well as cooler (Anderson et al., 2011; Zielhofer et al., 2019; Schirrmacher et al., 2020) at least until 3100 cal yr BP (Jiménez-Moreno et al., 2013; Ramos-Román et al., 2016). Later, a wet period beginning *ca.* 2600 cal yr BP occurred in SN and nearby areas (Martín-Puertas et al., 2010; Nieto-Moreno et al., 2011; Ramos-Román et al., 2018b).

For this initial phase of subzone CAMA2, the pollen diagram (Figs. 4 and 6) shows minimum percentages of AP (Table 4). *Pinus* maintains its steady decline, while, among herbs, ANC significantly advances, whereas *Vicia faba* joins Cerealia for the first time. Indicators of both Erosive and Wet phases also become notable. The former is made up mainly of *Glomus* (HdV 207), which has been also related to the decline of conifer forests (Montoya et al., 2018). The appearance of the latter, HdV 18, in a marked arid phase has been linked to the advent of rudimentary irrigation (Abel Schaad and López-Sáez, 2013). These clear traces of human presence near the study area are also revealed by a higher incidence of local fires, denoted by the increasing levels of larger particles of microcharcoal, which reach a maximum *ca.* 3800 cal yr BP.

The pine forest demise in DC coincides with this date. This depletion also appears to be more or less coetaneous in the other records of SN (Fig. 1), but not always so strong (Anderson et al., 2011; Jiménez-Moreno and Anderson, 2012; Mesa-Fernández et al., 2018). In addition to the aforementioned harsh climatic conditions, human activities would have reduced the tree cover to its minimum. A peak of fire incidence has also been reported between *ca.* 4200 and 3700 cal yr BP in most of the records studied in the area (Carrión et al., 2007; Gil-Romera et al., 2010; Manzano et al., 2019). The concurrent record of other indicators cited in DC suggests a human origin to these fires. In fact, this period coincides with the broadest expansion of the Argaric culture (*ca.* 4200-3500 cal yr BP) in south-eastern Iberia during the Early Bronze Age, entailing a surge in population and an intensification of metallurgy (García-Alix et al., 2013; Fyfe et al., 2019; Schirrmacher et al., 2020). The cultivation of legumes, such as *Vicia faba*, as well as small-scale irrigation (Lull et al., 2013) could point to the presence of this culture in this high-mountain environment.

Likewise, the collapse of this culture at *ca.* 3500 cal yr BP could be reflected in the pollen record (Figs. 4 and 6; Table 4) by the decrease of ANC, the disappearance of *Vicia faba*, Erosive and Wet Phases indicators against the sporadic occurrence of coprophilous fungi and the slight increase of AZNC. These cues appear to indicate a land-use change during the Late Bronze Age, oriented more towards livestock (Lull et al., 2013). After the disintegration of the Argaric culture, a major population drop has been recorded in south-eastern Iberia (Schirrmacher et al., 2020), which could have favoured the recovery of forests, particularly at high altitudes. The persistent aridity, high rate of landscape degradation and the impact of grazing would have prevented the growth of woody vegetation (Fyfe et al., 2019).

In any case, mining formed a major part of human activities at this time in this region (Carrión et al., 2003), as shown by the numerous sites located close to the study area (Fig. 2), most of them in use since the Bronze and even the Copper Age (Adroher Aroux et al., 2017). They became more robust with the onset of Iron Age (i.e. Iberian Period), *ca.* 2850 cal yr BP, when the population again increased, especially from 2550 cal yr BP onwards (Buxó, 2008).

The Iberian Period is considered a time of major human impact on south-eastern Spain (Gil-Romera et al., 2010; Anderson et al., 2011), where a high incidence of human-induced fires has also been documented (Carrión et al., 2007). Moreover, a wet period occurred in this region (Fig. 6), beginning *ca.* 2700 cal yr BP (Martín-Puertas et al., 2010; Jiménez-Moreno et al., 2013; Ramos-Román et al., 2018b), which appeared to favour a forest recovery. Increasing wetness was confirmed in the pollen diagram (Fig. 4; Table 4) by the almost continuous presence of HdV 18. However, no significant land-use change is detected in the record of DC, where the landscape appears to have remained almost unaltered during this period, compared to the previous one. A slight intensification of grazing activities, favoured by fires, presumably impoverished the grasslands, as indicated by the overlap of the increase of AZNC and Poaceae, the lowering of APP as well as the permanent levels of coprophilous fungi and microcharcoal particles.

#### 4.3. Crops vs. livestock on each slope of SN (*ca.* 2250-410 cal yr BP)

##### 4.3.1. Roman Period (*ca.* 2150-1530 cal yr BP)

The arrival of Romans (*ca.* 2150 cal yr BP) to the south-eastern Iberian Peninsula resulted in an overall landscape change (Carrión et al., 2010). The rise of temperatures, together with increasing rainfall (Martín-Puertas et al., 2010; Nieto-Moreno et al., 2011) helped the intensification of human pressure, particularly related to agriculture (McCormick et al., 2012). Nonetheless, mining activities were likely the first attraction for Romans in this area (González Román, 2001; García-Alix et al., 2013). Later, during the High Empire (*ca.* 1950-1750 cal yr BP), the general population underwent a further surge strongly associated with agriculture, which gave rise to a large number of *villae* as well as smaller and more scattered farms (Salvador Oyonate, 2012). Finally, the population levels fell during the Low Empire (*ca.* 1750-1530 cal yr BP), when agrarian lands became consolidated into fewer hands and the populace moved to higher elevations (González Martín, 2014).

The core of DC (Fig. 4 and 6; Table 4) initially registered a minimum of AP, perhaps related to the simultaneous rise in metallurgy (Bertrand and Sánchez Viciano, 2008). Thereafter, a slight recovery of oak forests matches the reported wettest period around *ca.* 1700 cal yr BP (Martín-Puertas et al., 2010; Nieto-Moreno et al., 2011), despite greater human pressure, also observed in nearby deposits (Carrión et al., 2003, 2007; Anderson et al., 2011). Fires prevented further forest recovery, firstly affecting heathlands and later also forests, as landowners sought to provide grazing for the increasing livestock density, this being reflected by the growing levels of coprophilous fungi, although pastoralism has not been particularly related to Romans (Gómez Pantoja, 2016). The DC record also indicates an increasing trend in AZNC, pointing to a parallel increase in grazing pressure, possibly related to the spread of different kinds of farms in the nearby area (Salvador Oyonate, 2012). Moreover, the synchronous high levels of microcharcoals coincide with maxima in most of the records of SN (Jiménez-Moreno et al., 2013), which show no imprint of grazing activities, whereas they are located at higher elevations. By contrast, livestock pressure clearly increased in Sierra de Gádor during this period (Carrión et al., 2003).




Figure 6. Main environmental changes in Sierra Nevada during the last 5300 years based on the analysis of Dehesa del Camarate (DC) and Cañar mires. (Aeolian dust input from Jiménez-Espejo et al., 2014; Aridity trend of Río Seco Lake from Anderson et al., 2011; Dry versus humid climate conditions of the west Balearic-Algerian Basin 306G core from Nieto-Moreno et al., 2011; NAO Index from Olsen et al., 2012; Palaeodemographic proxy (SPDs) for SE Spain from Fyfe et al., 2019; Pb/Al of Río Seco Lake from García-Alix et al., 2013). Humid (blue) and Dry (yellow) phases have been set based on Martín-Puertas et al., 2010; Nieto-Moreno 2011; 2013; and García-Alix et al., 2017. MOD: Modern Period; CO: Contemporary Period.

On the other hand, crops appear to have diminished, since Cerealia became scarcer, albeit *Vicia faba* appears in the record again. The almost continuous curves of *Castanea* and *Olea* should also be noted, as characteristic cultivated tree species of this period

(Rodríguez Ariza and Montes Moya, 2005), despite the fact that both of them appeared sporadically much earlier in the profile.

The pollen diagram of Cádiz (Figs. 5 and 6; Table 5) started shortly after the arrival of the Romans (*ca.* 2080 cal yr BP) and shows an open landscape, widely dominated by grasslands, with a significant area occupied by cereal crops. The remarkable low levels of AP indicate intense deforestation in former times, likely related to mining or agriculture (Martín Civantos, 2014). The curves for both *Cerealia* and *Secale cereale* (rye) indicate their local cultivation (López Sáez and López Merino, 2005). *Vicia faba* is also recorded in the first sample. The low levels of coprophilous fungi and AZNC reflect the minor amount of livestock grazing on this southern slope as well as the limited associated fires, pointed by microcharcoals. Indicators of Erosive phases could be linked to agricultural activities, focused mainly on herbaceous crops, as pointed by the absence of *Castanea* or *Olea* pollen during this period.

Therefore, over the Roman period, a distinct land use was taking shape on each side of SN. While crops grew on the southern slopes, even at rather high altitudes, livestock grazing appeared to be better adapted to northern slopes, where cooler temperatures would have hindered a major expansion of agriculture. At the end of this period, a slight decrease is revealed both by crops and by coprophilous fungi, presumably linked to the aforementioned Low Empire crisis, also recorded in DC.

#### 4.3.2. Middle Ages (*ca.* AD 420-1492)

After the fall of the Roman Empire (*ca.* AD 420), a period of political instability began in south-eastern Iberia, marked by the Visigothic Period *ca.* AD 589 (Salvador Ventura, 2012). Meanwhile, dryness progressively increased from about 350 AD until peaking around AD 650, accompanied by cooler temperatures, typifying Dark Ages (Nieto-Moreno et al., 2011). In DC, Poaceae reaches a maximum in the profile (Figs. 4 and 6), apparently in relation to the abandonment of livestock grazing and cereal cultivation, but it appears to be more closely related to human than to climate-induced changes, as cereal crops recover and even surpass their levels in the following samples. This trend is more pronounced in Cádiz, where *Cerealia* reaches its highest percentages, with notable levels of *Secale cereale* and, for the first time, the sporadic occurrence of *Vicia faba*, while coprophilous fungi, indicating livestock grazing, remain almost absent.

In the early 8<sup>th</sup> century (AD 711), the Islamic Period begins in Iberia, lasting until the end of 15<sup>th</sup> century (AD 1492). The beginning coincides roughly with the onset of a new phase of aridity, the so-called Medieval Climate Anomaly (MCA), from AD 800 to 1300 (Nieto-Moreno et al., 2011), more intense between AD 1000 and 1300 (Nieto-Moreno et al., 2013), and also warmer than the previous one (Moreno et al., 2012).

Overall, al-Andalus communities were based on a multiproductive system. Although agriculture was the main land use, both metallurgy and livestock were well represented in the area, the former mostly near north-western slopes (Martínez Ruiz, 1999; Martín Civantos, 2001, 2014; Liarte Alcaine, 2010). Farms were not large, particularly in mountain areas, but grew a large number of crops, through the installation of new irrigation systems as well as the communal use of more distant lands (Trillo San José, 2004). Both the deposit of DC and Cádiz (Fig. 1) are currently located next to an irrigation channel (MEMOLA Project, 2014). All of these infrastructures are believed to have been built towards the first half of 10<sup>th</sup> century (Trillo San José, 2004).

During the initial period of Islamic Period, until *ca.* AD 950, few changes appear in the pollen diagram of Cádiz (Figs. 5 and 6), revealing the transition towards new ways of land management. After this short event, a shift begins with the first appearance of *Linum*, the decline in *Cerealia* as well as the increase in *Secale cereale* and, to a lesser extent, of

*Vicia faba*. In addition, coprophilous fungi reached significant levels, coinciding with the return of HdV 18. This NPP, indicating wet phases in the most arid segment of the MCA (Nieto-Moreno et al., 2013), could be related to the first traces of irrigation in this area. The initial stages of the following subzone, CAÑB1, until ca. AD 1250, follow the same trend, beginning with a major increase of coprophilous fungi and the virtual demise of Cerealia, while *Secale cereale* remains the main crop. This severe human pressure is accompanied by a stronger incidence of fires, reflected in the remarkable proliferation of microcharcoals, presumably to reduce the extent of shrubland and increase the grassland areas. This earlier part of the subzone spans the establishment of the Caliphate in the 10<sup>th</sup> century, and the start of the “al-Andalus agricultural revolution” during the 11<sup>th</sup> century (Hernández Bermejo and García Sánchez, 2000). Irrigation systems were developed and new crops were grown. In this sense, dry farming was virtually abandoned, these lands being transformed to pastureland, mostly in mountain areas (Trillo San José, 2004). The profile of Cádiz clearly shows this dynamic (Fig. 5). Cereal was previously grown without irrigation, as in the Roman agricultural system (Beltrán Lloris and Willi, 2011) in wet areas, but at this time the crop disappears from the pollen diagram, parallel to the spread of coprophilous fungi, pointing to the exploitation of those lands by livestock grazing. *Secale cereale*, better adapted to poor soils and the lower temperatures of high-altitude areas (Hernández Bermejo and García Sánchez, 2008), would eventually be irrigated in spring, as described for this area even in recent times (Guzmán Álvarez, 2010; Martín Civantos 2014).

Moreover, in the following phase, coinciding with the Nasrid Kingdom (ca. AD 1250-1500), dry farming lands were virtually abandoned, allowing the marked expansion of *Quercus pyrenaica* (Figs. 5 and 6), and irrigation networks would have been expanded to water pasturelands also, as indicated by the increase in Poaceae and, especially, Cyperaceae, with the parallel occurrence of high peaks of coprophilous fungi. In addition, new fields were ploughed for the cultivation of rye, as reflected by the concurrent maxima of *Secale cereale* and Erosive phases indicators. Herds, mainly of goats, would stay only seasonally in these high-altitude areas, as indicated also by the low percentages of AZNC, within a short-distance transhumance system (Martínez Ruiz, 1999; Trillo San José, 2004). It should also be noted that a new climatic phase, the Little Ice Age (LIA) began ca. AD 1300 (Nieto-Moreno et al., 2013), shifting to a colder and slightly wetter climate, which could have further improved the cultivation of rye against other grains.

Similarly, the pollen diagram of DC (Fig. 4) reveals the likely broadening of irrigation ca. AD 850, with the occurrence of HdV 18 shortly after the onset of MCA (Nieto-Moreno et al., 2013), parallel to peaks for indicators of Erosive phases. The retreat of Poaceae and the concurrent increase in APP, mainly due to Apiaceae, could be attributed to an enrichment in palatable species among grasslands through irrigation, as reported by other records in the Iberian Central System (Abel-Schaad and López-Sáez, 2013). However, the absence of concurrent grazing indicators could point to the cultivation of any crop belonging to this family, currently fallen in disuse, such as *Smiranium olusatrum* or *Foeniculum vulgare* (Hernández Bermejo and García Sánchez, 2000). Later, during the Nasrid Kingdom (ca. AD 1250-1500), grazing pressure increased slightly, as shown by the levels of coprophilous fungi, the seasonal nature of the grazing being indicated by the limited percentages of AZNC. On the other hand, *Olea* undergoes a notable advance, especially during the last two centuries of this period. By contrast, *Castanea* is the main tree crop on the southern slopes in Cádiz, both trees attesting to agricultural intensification during Islamic Period, even in mountain areas (Trillo San José, 2004).

#### 4.4. The final landscape shaping of SN (ca. AD 1500-present)



#### 4.4.1. Modern Period (ca. AD 1500-1800)

Wet and dry phases alternated within the colder climate corresponding to LIA in this area (Nieto-Moreno et al., 2013).

The pollen diagrams (Figs. 4, 5 and 6) suggest an earlier change in northern slopes, ca. AD 1540, than in the southern ones, where prior conditions remained until ca. AD 1600. These dates could be considered the beginning of the repopulation on each side of the mountain by new settlers coming from northern areas (Trillo San José, 2004). This overall change initially involved a severe depopulation (Bañuelos Arroyo, 2008), the slow reconstruction of villages and irrigation systems (Romero Funes, 2016), as well as a greater amount of livestock raising and rainfed farming within a cereal-fallow-pasture system (Sayadi and Calatrava, 2001).

The DC core shows a slight early advance of AP (Figs. 4 and 6), led by *Pinus sylvestris* and *Quercus ilex*, perhaps favoured by colder temperatures in addition an easing of human pressure. *Castanea*, absent since Roman times, recovers its continuous curve, while *Olea* maintains previous levels and Cerealia slightly diminishes, almost disappearing. Conversely, livestock grazing appears to intensify, as indicated by the rising levels of coprophilous fungi, aided by the use of fire on shrublands, with maximum values for microcharcoals. The drop in both HdV 18 and Cyperaceae, as well as the rising levels of Dry phases indicators, could be related to the abandonment of the prior irrigation system. A similar trend was noted in Cádiz (Figs. 5 and 6), although a few decades later. The pattern of higher percentages for Cerealia and lower ones for *Secale cereale* marks the beginning of such a change. AP also increases, but on this south side led by *Quercus pyrenaica* and *Q. ilex*, as well as the tree crops *Castanea* and *Olea*. Likewise, irrigation appears initially to disappear. Rainfed cereal farming alternated with pastures for livestock, which would be more stable, as indicated by the increase in AZNC.

By AD 1650 and soon afterwards, cultivated areas doubled through the ploughing of new lands, and livestock herds substantially increased (Sayadi and Calatrava, 2001; Bañuelos Arroyo, 2008).

At this time, microcharcoals reach their maxima in DC (Figs. 4 and 6; Table 4) coinciding with high levels of coprophilous fungi and increasing percentages of ANC and AZNC. In addition, Cerealia and *Olea* maintain their levels, whereas *Quercus ilex* and *Q. pyrenaica* diminish, likely related to their use as charcoal and firewood. These trends imply a panorama of overexploitation of natural resources, well described in the chronicles of this period (Mesa Garrido, 2016).

By contrast, the diagram from Cádiz (Figs. 5 and 6) appears to point to less human pressure, as might be inferred from the increase of AP (Table 5), not only by cultivated trees such as *Castanea*, *Olea* or *Juglans*, but also, and especially, by *Quercus pyrenaica*. In the mid 18<sup>th</sup> century the first management measures were implemented for forests in Spain, in order to increase the timber production for shipbuilding (Martínez González, 2014). Most oak forests from Cádiz and other nearby areas were included in a catalogue of these guarded forests (Mesa Garrido, 2016). On the other hand, both Cerealia and mostly *Secale cereale* increased but far from previous percentages. Conversely, coprophilous fungi fall but still display significant levels, presumably revealing more stable livestock rearing, as shown by the concurrent rising levels of AZNC, within a likely integrated cycle associated with crops (Sayadi and Calatrava, 2001). Both the low levels of microcharcoals and ANC, as well as the retreat of shrubland with the advance of grasslands would support this scenario. Likewise, the significant rise of HdV 18 levels as well as the slighter one of Cyperaceae and riparian woods, could point to the restart of the prior irrigation channel.

#### 4.4.2. Contemporary Period (ca. AD 1800-present)

The overall impact on vegetation during this period on the one hand is negative, as a result of the opening of new mines, large-scale ploughing, and increasing livestock herds. On the other hand, it is positive for the establishment of a set of regulations intended to protect extant forests and restore degraded areas (Sayadi and Calatrava, 2001; Valbuena-Carabaña et al., 2010; Mesa Garrido, 2016). This new time period also overlaps with the so-called Industrial Period, during a period characterized by increasing moisture in this region, which lasted until the 2<sup>nd</sup> half of the 20<sup>th</sup> century. Then, rainfall followed a negative trend until the present (Nieto-Moreno et al., 2013).

A significant increase in AP begins the pollen diagram of DC (Figs. 4 and 6; Table 4), mainly due to *Olea*, but also to *Pinus sylvestris*, likely favoured by increasing wetness and by the implementation of Reforestation Plans in this area from AD 1940 onwards (Mesa Garrido, 2016). Other tree taxa, such as *Quercus* or riparian woodlands expanded only slightly in the last sample, as a consequence of the later establishment of protective measures during the 20<sup>th</sup> century, resulting in the establishment of the Sierra Nevada National Park. Previously, the successive Disentailment Laws in the mid-19<sup>th</sup> century triggered further pressure on vegetation (Valbuena-Carabaña et al., 2010), specially indicated by the high levels of microcharcoals and coprophilous fungi. The rising values of HdV 18, Cyperaceae, together with the enrichment of grasslands indicated by APP, may reflect a broader irrigation system for pastures, which would have allowed a longer presence of herds in the area, as suggested by the increase in AZNC.

The diagram from Cádiz (Figs. 5 and 6; Table 5) reveals a shift towards a more livestock-oriented land use at the expense of crop cultivation, reflected in the decreasing percentages of herbaceous crops, especially *Secale cereale*, the spread of grasslands and the concurrent advance of coprophilous fungi. The irrigation of these pasturelands, indicated by the maxima of HdV 18, would have likewise extended the stay of flocks, within the traditional short-distance transhumance system (Rubio de Lucas et al., 1995). Tree crops such as *Olea* and particularly *Castanea* also became more prevalent in the area during this period. Among the other tree taxa, *Quercus pyrenaica* maintained its percentages throughout most of 19<sup>th</sup> and 20<sup>th</sup> centuries, probably because of the exemption of their forests in Cádiz from the ones included in the Disentailment Laws and their subsequent accreditation as Public Mounts. They were likely used by the local settlers for firewood, leaves or acorns but maybe not for grazing (Mesa Garrido, 2016). The strikingly low levels of microcharcoals in the pollen diagram support this idea. In any case, *Q. pyrenaica* is considered resilient to human impact (Valbuena-Carabaña and Gil, 2013) and even benefits by it (Salomón et al., 2017). This was one of the first areas where reforestation works started (Mesa Garrido, 2016), in the beginning of 20<sup>th</sup> century. *Pinus pinea/halepensis* and *P. sylvestris* display a significant increase in the top sample, likely related to these tasks, but also to the lowering of human pressure and the new management measures after the Declaration of SN as National Park.

#### 4.5. Implications for Conservation

The conservation of cultural landscapes, such as SN (Jiménez-Olivencia et al., 2006; Gómez-Ortiz et al., 2013) requires integration of a long-term perspective for current and future management strategies (Gillson, 2015; Whitlock et al., 2018). Land-use history becomes an essential information in the Mediterranean, where human activities developed since at least the Middle Holocene (Davis, 2015), with a distinct dynamics in mountain areas (Zamora et al., 2015; Catalán et al., 2019). In this sense, our work provides some valuable issues which should be taken in consideration:

- Forest cover was quite thinner in the final stages of the mid-Holocene (*ca.* 5400 cal yr BP) than currently on the northern slopes of SN, with a minor extent of pinewoods, which show a high vulnerability against human activities. Environmental change appears to have

been induced by human influence rather than by climate since then. The collapse of ancient forests could be set at *ca.* 4000 cal yr BP.

- Current forest cover in SN has expanded quite recently. Oak forests of *Quercus pyrenaica* only spread since *ca.* AD 1250 on the southern slopes, first helped by the abandonment of dry-farming lands and the multiple resources they provided, and later by the increasing protection measures. Both *Castanea* and *Olea*, despite their early origin, gain relevance during the last three centuries. Current pine woodlands were mainly planted in 20<sup>th</sup> century.

- Grasslands have been the predominant part of vegetation during, at least, 5000 years, together with other herbaceous species. They have been fostered and kept by humans for agriculture, mostly for pastures. Irrigation systems enhanced their maintenance, especially since *ca.* AD 1000, during Islamic Period and later on, becoming the basis of the traditional multiproductive system.

Therefore, the high biodiversity of SN has been also preserved and promoted by humans over the last millennia, beyond the influence of progressive climate changes. Current and future warming is not the sole thread to biodiversity that environmental managers should cope, but also land abandonment related to depopulation, which only could be controlled with the preservation of traditional human activities like grazing, high-elevation cropping or irrigation systems. In this sense, the implementation of good practices for maintaining these long-established activities becomes essential.

## 5. CONCLUDING REMARKS

In conclusion, the analysis of two pollen records from both sides of SN in the western Mediterranean mountains provides answers to the two main research questions posed in this paper.

First, the main tipping point when natural systems transformed to cultural landscapes occurred during the Chalcolithic (*ca.* 5400 cal yr BP). This transformation was defined by very low percentages of AP (half of current forest cover), and concurrent traces of human activities. So far, human impact in SN had been established more than 1000 years later, during the Bronze Age.

Second, a further intensification of human activities occurred during the Argaric culture (*ca.* 3800 cal yr BP) and reached its maximum degree with the onset of Iron Age, when climate became more humid. Mineral resources, but not solely, apparently acted as main drivers of population growth, causing a severe decline of ancient forests.

Later, Romans triggered an increase of livestock pressure to northern slopes, whereas crops grew on the southern ones, but with a lesser extent as described at low elevations.

A significant land-use change occurred during Islamic Period, with the installation of an irrigation network within a multiproductive system, against the warm and dry conditions of the MCA.

Climate hampered an eventual forest recovery due to the arid mid-to-late Holocene conditions. Later, despite favourable climatic conditions during certain periods, only land-use changes have allowed the spread of the tree cover, in the form of orchards (*Castanea* and *Olea*), afforestation (*Pinus*) or the development of secondary forests (*Quercus pyrenaica*). Ultimately, the implementation of environmental protection measures has promoted the spread of the most extensive forest cover.

SN mountain vegetation has been strongly changing over a long temporal scale. Long-term studies involving climate-human-vegetation interactions are required for current ecosystem restoration and sustainable forest management, as well as for a better understanding of SN forest resilience and the tipping points which occurred in the past.

The preservation of traditional human activities becomes crucial to ensure orophilous-mesophilous biodiversity in the face of oncoming threats.

#### Declaration of Competing Interest

None

#### Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### Acknowledgements

**Funding:** This study was supported by Spanish government: State R&D Program Oriented to the Challenges of the Society, Project references RTI2018-101714-B-I00 and CS02017-83576-P; FEDER/ Andalusian Government - Ministry of Economy and Knowledge / Project reference B-RNM-404-UGR18; and Andalusian Plan for Research, Development and Innovation, PAIDI 2020, Project reference P18-RT- 4963. We are grateful to Consejería de Agricultura, Ganadería, Pesca y Desarrollo Sostenible (Andalusian Government) and Sierra Nevada National Park for providing sampling permissions and access to the study area.

Acknowledgements

## References

- Abel-Schaad, D., López-Sáez, J.A., 2013. Vegetation changes in relation to fire history and human activities at the Peña Negra mire (Bejar Range, Iberian Central Mountain System, Spain) during the past 4,000 years. *Veg. Hist. Archaeobot.* 22 (3), 199-214. doi: <https://doi.org/10.1007/s00334-012-0368-9>.
- Adroher Auroux, A., López Marcos, A., Pachón Romero, J.A. 2002. *La cultura ibérica*. Granada Arqueológica. Granada.
- Adroher Auroux, A.M., García Campoy, C.A., González Martín, J.A., Peregrín Pitto, E., Sol Plaza, J., 2017. Minería Ibérica en Sierra Nevada (Granada) y su perduración en el paisaje actual. El complejo arqueológico de El Cardal (Ferreira). En García-Pulido, L.J., Arboledas Martínez, L., Alarcón García, E. & Contreras Cortés, F. (Eds.): *Presente y futuro de los paisajes mineros del pasado: estudios sobre minería, metalurgia y poblamiento*, Editorial Universidad de Granada, Granada, pp. 341-353.
- Alba-Sánchez, F., Abel-Schaad, D., López-Sáez, J.A., Sabariego Ruiz, S., Pérez-Díaz, S., González-Hernández, A., 2018. Paleobiogeography of *Abies* spp. and *Cedrus atlantica* in the Western Mediterranean (Iberian Peninsula and Morocco). *Ecosistemas* 27(1), 26-37. doi.: 10.7818/ECOS.1441.
- Anderson, R.S., Jiménez-Moreno, G., Carrión, J.S., Pérez-Martínez, C. 2011., Postglacial history of alpine vegetation, fire, and climate from Laguna de Río Seco, Sierra Nevada, southern Spain. *Quaternary Sci. Rev.* 30, 1615-1629. doi: <https://doi.org/10.1016/j.quascirev.2011.03.005>.
- Bañuelos Arroyo, A., 2008. *Repoblación y demografía del concejo de Cástaras y Nieves*. Asociación Cultural de Cástaras y Nieves. Cástaras (Granada).
- Beltrán Lloris, F., Willi, A. 2011., *El regadío en la Hispania romana. Estado de la cuestión*. Cuadernos de Prehistoria y Arqueología de la Universidad de Granada, 21: 9-56.
- Bertrand, M., Sánchez Viciano, J.R., 2008. Production du fer et peuplement de la région de Guadix (Grenade) au cours de l'Antiquité tardive et du haut Moyen Âge. In A. Canto García A., Cressier, P (Eds.): *Minas y metalurgia en al-Andalus y Magreb occidental*, 123-157. Casa de Velázquez, Madrid.
- Birks, H.J.B. 2019., Contributions of Quaternary botany to modern ecology and biogeography. *Plant. Ecol. Divers.* 12(3-4), 189-385. doi: <https://doi.org/10.1080/17550874.2019.1646831>.
- Blanca López, G., López Onieva, M.R., 2002. *Flora amenazada y endémica de Sierra Nevada*. Consejería de Medio Ambiente. Junta de Andalucía.
- Blanca, G., Cueto, M., Romero, M.T., 2019. Rareza y endemismo en la flora vascular de Sierra Nevada. En: Peñas, J. y Lorite, J. (eds). *Biología de la Conservación de Plantas en Sierra Nevada. Principios y Retos para su Preservación*. Servicio de Publicaciones de la Universidad de Granada, Granada, 113-128.
- Blanco-González, A., Lillios, K.T., López-Sáez, J.A. Brake, B.L. 2018. Cultural, Demographic and Environmental Dynamics of the Copper and Early Bronze Age in Iberia (3300–1500 BC): Towards an Interregional Multiproxy Comparison at the Time of the 4.2

- ky BP Event. *J. World Prehist.* 31, 1–79. doi: <https://doi.org/10.1007/s10963-018-9113-3>.
- Blaauw, M., 2010. Methods and code for ‘classical’ age-modelling of radiocarbon sequences. *Quat. Geochronol.* 5(5), 512–518. doi: <https://doi.org/10.1016/j.quageo.2010.01.002>.
- Buxó, R., 2008. The agricultural consequences of colonial contacts on the Iberian Peninsula in the first millennium B. C. *Veg. Hist. Archaeobot.* 17(1), 145-154. doi: <https://doi.org/10.1007/s00334-007-0133-7>.
- Camuera, J., Jiménez-Moreno, G., Ramos-Román, M. J., García-Alix, A., Toney, J. L., Anderson, R. S., Jimenez-Espejo, F., Kaufman, D., Bright, J., Webster, C., Yanes, Y., Carrion, J.S., Ohkouchi, N., Suga, H., Yamame, M., Yokoyama, Y., Martínez-Ruiz, F., 2018. Orbital-scale environmental and climatic changes recorded in a new ~200,000-year-long multiproxy sedimentary record from Padul, southern Iberian Peninsula. *Quaternary Sci. Rev.* 198, 91-114. doi: <https://doi.org/10.1016/j.quascirev.2018.08.014>
- Camuera, J., Jiménez-Moreno, G., Ramos-Román, M. J., García-Alix, A., Toney, J. L., Anderson, R. S., Jimenez-Espejo, F., Bright, J., Webster, C., Yanes, Y., Carrión, J.S., 2019. Vegetation and climate changes during the last two glacial-interglacial cycles in the western Mediterranean: A new long pollen record from Padul (southern Iberian Peninsula). *Quaternary Sci. Rev.* 205, 86-105. doi: <https://doi.org/10.1016/j.quascirev.2018.12.013>
- Carrasco Rus, J., Morgado, A., Martínez-Sevilla, Y. F., 2016. Implantación y desarrollo de los grupos neolíticos del sur de la península ibérica. Reflexiones sobre algunos modelos interpretativos desde los inicios del siglo XXI. In *Del neolítico a l’edat del bronze en el Mediterrani occidental. Estudis en homenatge a Bernat Martí Oliver*. TV SIP 119, Valencia, pp. 159-180.
- Carrión, J.S., Sánchez-Gómez, P., Mota, J.F., Yll, R., Chaín, C., 2003. Holocene vegetation dynamics, fire and grazing in the Sierra de Gádor, southern Spain. *Holocene* 13 (6), 839-849. doi: <https://doi.org/10.1191/0959683603hl662rp>.
- Carrión, J.S., Fuentes, N., González-Sampériz, P., Sánchez Quirante, L., Finlayson, J.C., Fernández, S., Andrade, A., 2007. Holocene environmental change in a montane region of southern Europe with a long history of human settlement. *Quaternary Sci. Rev.* 26, 1455-1475. doi: <https://doi.org/10.1016/j.quascirev.2007.03.013>
- Carrión, J. S., Fernández, S., González-Sampériz, P., Gil-Romera, G., Badal, E., Carrión-Marco, Y., López-Merino, L., López-Sáez, J.A., Fierro, E. Burjachs, F., 2010. Expected trends and surprises in the Lateglacial and Holocene vegetation history of the Iberian Peninsula and Balearic Islands. *Rev. Palaeobot. Palynol.* 162(3), 458-475. doi: <https://doi.org/10.1016/j.revpalbo.2009.12.007>.
- Catalán, J., Batalla, M., Bonet, M.T., Brotons, L., Buchaca, T., Camarero, L., Clemente, I., Contreras, F., Domingo, F., Felip, M., Gassiot, E., García-Casas, D., Giralt, S., González-Sampériz, P., Jiménez-Moreno, G., Laborda, R., Lorite, J., Martín-Civantos, J.M., Mazzuco, N., Montes, L., Moreno, A., Ninterola, M., Pélachs, A., Pérez-Obiol, R., Piqué, R., Pla.Rabés, S., Saña, M., Sáez, A., Sebastián, M., Soriano, J.M., Terradas, X., Valero-Garcés, B., Villero, D., 2019. Análisis ecológico de la

culturización del paisaje de alta montaña desde el neolítico: Los parques nacionales de montaña como modelo. *Proyectos de investigación en Parques Nacionales: 2013-2017*. MITECO, Madrid.

- Cheddadi, R., Palmisano, A., López-Sáez, J. A., Nourelbait, M., Zielhofer, C., Tabel, J., Rhoujjati, A., Khater, C., Woodbridge, J., Lucarini, G., Broodbank, C., Fletcher, W.J., Roberts, C.N., 2019. Human demography changes in Morocco and environmental imprint during the Holocene. *Holocene* 29(5), 816-829. doi: <https://doi.org/10.1177/0959683619826657>.
- Cortés Sánchez, M., Jiménez Espejo, F.J., Simón Vallejo, M.D., Gibaja Bao, J.F., Carvalho, A. F., Martínez-Ruiz, F., Rodrigo-Gámiz, M., Flores, J.A., Paytan, A., López-Sáez, J.A., Peña-Chocarro, L., Carrión, J.S., Morales, A., Roselló, E., Riquelme, J.A., Dean, R., Salgueiro, E., Martínez Sánchez, R., Rubia, J.J., Lozano, F., Vera, J.L., Llorente, L., Bicho, N.F., 2012. The Mesolithic–Neolithic transition in southern Iberia. *Quaternary Res.* 77(2), 221–234. doi: [doi:10.1016/j.yqres.2011.12.003](https://doi.org/10.1016/j.yqres.2011.12.003).
- Davis, B.A., Collins, P.M., Kaplan, J.O., 2015. The age and post-glacial development of the modern European vegetation: a plant functional approach based on pollen data. *Veg. Hist. Archaeobot.* 24(2), 303-317. doi: <https://doi.org/10.1007/s00334-014-0476-9>.
- Fægri, K.; Iversen, J., 1989. *Textbook of Pollen Analysis* (4th edn). Chichester. Wiley.
- Fyfe, R.M., Woodbridge, J., Palmisano, A., Bevan, A., Shennan, S., Burjachs, F., Legarra, B., García Puchol, O., Carrión, J.S., Revelles, J., Roberts, C.N., 2019. Prehistoric palaeodemographics and regional land cover change in eastern Iberia. *Holocene* 29(5), 799-815. doi: <https://doi.org/10.1177/0959683619826643>.
- García-Ruiz, J. M., Tomás-Faci, G., Diarte-Blasco, P., Montes, L., Domingo, R., Sebastián, M., Lasanta, T., González-Sampériz, P., López-Moreno, J.I., Arnáez, J., Beguería, S., 2020. Transhumance and long-term deforestation in the subalpine belt of the central Spanish Pyrenees: An interdisciplinary approach. *Catena* 195: 104744. doi: <https://doi.org/10.1016/j.catena.2020.104744>.
- García Sanjuán, L., Murillo-Barroso, M., 2013. Social complexity in Copper Age southern Iberia (c. 3200–2200 cal BC): Reviewing the ‘state’ hypothesis at Valencina de la Concepción (Seville, Spain). In M. Cruz Berrocal, L. García Sanjuán & A. Gilman (Eds.), *The prehistory of Iberia: Debating early social stratification and the state*, 119-140. New York: Routledge.
- García-Alix, A., Jiménez-Espejo, F. J., Lozano, J.A., Jiménez-Moreno, G., Martínez-Ruiz, F., García Sanjuán, L., Aranda Jiménez, G., García Alfonso, E., Ruiz-Puertas, G., Anderson, R.S., 2013. Anthropogenic impact and lead pollution throughout the Holocene in Southern Iberia. *Sci. Total Environ.* 449, 451-460. doi: <https://doi.org/10.1016/j.scitotenv.2013.01.081>.
- García-Alix, A., Jiménez-Espejo, F.J., Toney, J.L., Jiménez-Moreno, G., Ramos-Román, M.J., Anderson, R.S., Ruano, P., Queralt, I., Delgado-Huertas, A., Kuroda, J., 2017. Alpine bogs of southern Spain show human-induced environmental change superimposed on long-term natural variations. *Sci. Rep.* 7, 7439. doi: <https://doi.org/10.1038/s41598-017-07854-w>.

- García-Alix, A., Jiménez-Moreno, G., Anderson, R.S., Jiménez-Espejo, F.J., Delgado Huertas, A., 2012. Holocene environmental change in southern Spain deduced from the isotopic record of a high-elevation wetland in Sierra Nevada. *J. Palaeolimnol.* 48, 471-484. doi: <https://doi.org/10.1007/s10933-012-9625-2>.
- Gil-Romera, G., Carrión, J. S., Pausas, J.G., Sevilla-Callejo, M., Lamb, H.F., Fernández, S., Burjachs, F., 2010. Holocene fire activity and vegetation response in South-Eastern Iberia, *Quaternary Sci. Rev.* 29, 1082–1092. doi: <https://doi.org/10.1016/j.quascirev.2010.01.006>.
- Gillson, L. 2015. Biodiversity conservation and environmental change: using palaeoecology to manage dynamic landscapes in the Anthropocene. OUP Oxford. doi: 10.1093/acprof:oso/9780198713036.001.0001
- Gómez-Ortiz, A., Oliva, M., Salvà-Catarineu, M., Salvador-Franch, F., 2013. The environmental protection of landscapes in the high semiarid Mediterranean mountain of Sierra Nevada National Park (Spain): historical evolution and future perspectives. *Appl. Geogr.* 42, 227-239. doi: <https://doi.org/10.1016/j.apgeog.2013.02.006>.
- Gómez-Pantoja, J.L., 2016. Algunos indicios de pastoralismo en Hispania romana. (Some indications of pastoralism in Roman Hispania). *Mélanges de l'École française de Rome: Antiquité*, 128(2). doi: <https://doi.org/10.4000/mefra.3521>
- González Martín, C.G., 2014. Poblamiento y territorio en el curso medio del Genil en época romana: nuevas aportaciones arqueológicas. *La villa romana de Salar. Florentia Iliberritana* 25, 157-194.
- González Román, C. 2001. Ciudad y poblamiento romano en la provincia de Granada durante el Alto Imperio. *Habis* 32, 271-296.
- Guzmán Álvarez, J.R. 2010. Usos y costumbres para el reparto del agua. En: Guzmán Álvarez, J. R. & Navarro Cerrillo, R. M. (Coords.), pp. 166-189. *El agua domesticada. Los paisajes de los regadíos de montaña en Andalucía*. Agencia Andaluza del Agua, Consejería de Medio Ambiente, Junta de Andalucía, Sevilla, 592 pp.
- Grimm, E.C., 2004. *TGView*. Springfield. Illinois State Museum.
- Hernández Bermejo, J.E., García Sánchez, E., 2000. Botánica económica y etnobotánica en al-Andalus (Península Ibérica siglos X-XV), un patrimonio desconocido de la humanidad. *Arbor* 166(654), 311-331. doi: <https://doi.org/10.3989/arbor.2000.i654.1016>.
- Hernández Bermejo, J.E., García Sánchez, E., 2008. Las gramíneas en al-Andalus. *Ciencias de la naturaleza en Al-Andalus. Textos y estudios*. VIII, CSIC.
- Hughes, P.D., Woodward, J.C., 2016. *Quaternary Glaciation in the Mediterranean Mountains*. Geological Society, London, Special Publications. doi: <https://doi.org/10.1144/SP433.14>.
- Jalut, G., Dedoubat, J.J., Fontugne, M., Otto, T., 2009. Holocene circum-Mediterranean vegetation changes: Climate forcing and human impact. *Quatern. Int.* 200 (1-2), 4-18. doi: <https://doi.org/10.1016/j.quaint.2008.03.012>.



- Jiménez-Espejo, F., García-Alix, A., Jiménez-Moreno, G., Rodrigo-Gámiz, M., Anderson, R.S., Rodríguez-Tovar, F.J., Martínez-Ruiz, F., Giral, S., Delgado Huertas, A., Pardo-Igúzquiza, E. 2014. Saharan aeolian input and effective humidity variations over western Europe during the Holocene from a high altitude record. *Chem. Geol.* 374-375, 1–12. doi: <https://doi.org/10.1016/j.chemgeo.2014.03.001>.
- Jiménez-Moreno, G., Anderson, R.S., 2012. Holocene vegetation and climate change recorded in alpine bog sediments from the Borreguiles de la Virgen, Sierra Nevada, southern Spain. *Quaternary Res.* 77, 44-53. doi: <https://doi.org/10.1016/j.yqres.2011.09.006>.
- Jiménez-Moreno, G., García-Alix, A., Hernández-Corbalán, M.D., Anderson, R.S., Delgado-Huertas, A., 2013. Vegetation, fire, climate and human disturbance history in the southwestern Mediterranean area during the Late Holocene. *Quaternary Res.* 79, 110-122. doi: <https://doi.org/10.1016/j.yqres.2012.11.008>.
- Jiménez Olivencia, Y., Porcel Rodríguez, L., Piñar Álvarez, Á. 2006. Evolución histórica de los paisajes del Parque Nacional de Sierra Nevada y su entorno. *Proyectos de investigación en Parques Nacionales*: 109-128.
- Kaplan, J.O., Krumhardt, K.M., Zimmermann, N., 2009. The prehistoric and preindustrial deforestation of Europe. *Quaternary Sci. Rev.* 28(27-28), 3016-3034. doi: <https://doi.org/10.1016/j.quascirev.2009.09.028>.
- Liarte Alcaine, M.R., 2010. La conquista musulmana de Hispania. *Revista de Clases historia*, (3), 47.
- Lillios, K.T., Blanco-González, A., Drake, B.L., López-Sáez, J.A., 2016. Mid-late Holocene climate, demography, and cultural dynamics in Iberia: A multi-proxy approach. *Quaternary Sci. Rev.* 135, 138–153. doi: <https://doi.org/10.1016/j.quascirev.2016.01.011>.
- López Sáez, J.A., López Merino, L., 2005. Precisiones metodológicas acerca de los indicios paleopalinológicos de agricultura en la prehistoria de la Península Ibérica. *Portugalia: Revista de Arqueología do Departamento de Ciências e Técnicas do Património* 26, 53-64.
- López-Sáez, J.A., López Merino, L., Pérez Díaz, S., Alba Sánchez, F. 2011., Paleopaisajes de Andalucía Oriental durante la transición Mesolítico-Neolítico antiguo. In J.F. Gibaja & A.F. Carvalho (Eds.), *Os últimos caçadores-recolectores e as primeiras comunidades produtoras do sul da Península Ibérica e do norte de Marrocos*. *Promontoria Monográfica* 15, 213–220. Faro: Universidade do Algarve.
- López-Sáez, J.A., Abel-Schaad, D., Pérez-Díaz, S., Blanco-González, A., Alba-Sánchez, F., Dorado, M., Ruiz-Zapata, B., Gil-García, M.J., Gómez-González, C., Franco-Múgica, F., 2014. Vegetation history, climate and human impact in the Spanish Central System over the last 9000 years, *Quatern. Int.* 353: 98–122. doi: <https://doi.org/10.1016/j.quaint.2013.06.034>.
- Lull, V., Micó, R., Rihuete Herrada, C., Risch, R., 2010. Metal and social relations of production in the 3rd and 2nd millennia BCE in the Southeast of the Iberian Peninsula. *Trabajos de Prehistoria* 67(2), 323-347. doi: 10.3989/tp.2010.10042.

- Magri, D., 2012. Quaternary history of *Cedrus* in Southern Europe. *Annali di Botanica* 2, 57-66. doi: <https://doi.org/10.4462/annbotrm-10022>.
- Manzano, S., Carrión, J.S., López-Merino, L., Jiménez-Moreno, G., Toney, J.L., Amstron, H., Anderson, R.S., García-Alix, A., Guerrero, J.L., Sánchez-Mata, D., 2019. A palaeoecological approach to understanding the past and present of Sierra Nevada, a Southwestern European biodiversity hotspot. *Global Planet. Change* 175, 238-250. doi: <https://doi.org/10.1016/j.gloplacha.2019.02.006>.
- Martín Civantos, J.M., 2001. Alquife, un castillo con vocación minera en el Zenete (Granada). *Arqueología y territorio medieval* 8, 325-345. doi: <https://doi.org/10.17561/aytm.v8i0.1686>.
- Martín Civantos, J.M. 2007. Poblamiento y territorio medieval en el Zenete (Granada) Editorial Universidad de Granada. Granada.
- Martín Civantos, J.M., 2014. Mountainous landscape domestication. Management of non-cultivated productive areas in Sierra Nevada (Granada-Almería, Spain). *European Journal of Post-Classical Archaeologies* 4, 99-130.
- Martín Socas, D., Buxó, R., Camalich, M.D., Goñi, A., 1999. Estrategias subsistenciales en Andalucía Oriental durante el Neolítico. *SAGVNTVM Extra* 2, 25-30.
- Martínez González, A.J., 2014. The development of the forest Ordinance for the Navy, 31st January 1748, the oceanic political basis of the Spanish monarchy during the eighteenth century. *Anuario de Estudios Americanos* 71 (2), 571-602. doi: [10.3989/aeamer.2014.2.07](https://doi.org/10.3989/aeamer.2014.2.07).
- Martínez Ruiz, E., 1999. Los montes en la cultura islámica: El bosque de Al-Andalus desde el siglo VIII al XIII. In: Calzado, A., Domingo, J.M., Marín, F.J. (Coords) *Los montes y su historia: una perspectiva política, económica y social: I Jornadas Forestales: historia, socioeconomía y política forestal*, 63-88. Universidad de Huelva.
- Martin-Puertas, C., Jiménez-Espejo, F., Martínez-Ruiz, F., Nieto-Moreno, V., Rodrigo, M., Mata, M.P., Valero-Garcés, B.L., 2010. Late Holocene climate variability in the southwestern Mediterranean region: an integrated marine and terrestrial geochemical approach. *Clim. Past.* 6(6), 807-816. doi: <https://doi.org/10.5194/cp-6-807-2010>.
- McCormick, M., Büntgen, U., Cane, M.A., Cook, E. R., Harper, K., Huybers, P., Litt, T., Manning, S.W., Mayewski, P.A., More, A.F.M., Nicolussi, K., Tegel, W., 2012. Climate change during and after the Roman Empire: reconstructing the past from scientific and historical evidence. *J. Interdiscipl. Hist.* 43(2): 169-220. doi: <https://www.jstor.org/stable/41678664>.
- McNeill, J.R. 2003. *The mountains of the Mediterranean world*. Cambridge University Press. Doi: <https://doi.org/10.1017/CBO9780511529023>
- Médail, F., Diadema, K., 2009. Glacial refugia influence plant diversity patterns in the Mediterranean Basin. *Journal of Biogeography* 36 (7), 1333-1345. doi: <https://doi.org/10.1111/j.1365-2699.2008.02051.x>.
- MEMOLA Project (Mediterranean Mountainous Landscapes). 2014. Mapa colaborativo de regadíos históricos de Granada y Almería. <https://regadiohistorico.es>

- Mercuri, A.M., Sadori, L., 2014. Mediterranean Culture and Climatic Change: Past Patterns and Future Trends. In: Goffredo S., Dubinsky Z. (Eds) *The Mediterranean Sea*. Springer, Dordrecht. doi: [https://doi.org/10.1007/978-94-007-6704-1\\_30](https://doi.org/10.1007/978-94-007-6704-1_30).
- Mesa Garrido, M. Á., 2016. Geografía y política forestal. Análisis general de la gestión de los montes en la provincia de Granada. Siglos XVIII-XX (Doctoral dissertation, Universidad de Granada).
- Mesa-Fernández, J.M., Jiménez-Moreno, G., Rodrigo-Gámiz, M., García-Alix, A., Jiménez-Espejo, F.J., Martínez-Ruiz, F., Anderson, R.S., Camuera, J., Ramos-Román, M.J., 2018. Vegetation and geochemical responses to Holocene rapid climate change in the Sierra Nevada (southeastern Iberia): the Laguna Hondera record. *Clim. Past.* 14(11), 1687-1706. doi: <https://doi.org/10.5194/cp-14-1687-2018>.
- Miola, A., 2012. Tools for Non-Pollen Palynomorphs (NPPs) analysis: A list of Quaternary NPP types and reference literature in English language (1972–2011). *Rev. Palaeobot. Palyno.* 186, 142–161. doi: <https://doi.org/10.1016/j.revpalbo.2012.06.010>.
- Montoya, E., Rull, V., Vegas-Vilarrúbia, T., Corella, J. P., Giralt, S., Valero-Garcés, B., 2018. Grazing activities in the southern central Pyrenees during the last millennium as deduced from the non-pollen palynomorphs (NPP) record of Lake Montcortès. *Rev. Palaeobot. Palynol.* 254, 8-19. doi: <https://doi.org/10.1016/j.revpalbo.2018.04.005>.
- Mooney, S.D., Tinner, W., 2011. The analysis of charcoal in peat and organic sediments. *Mires Peat* 7(9), 1–18. <http://www.mires-and-peat.net/>, ISSN 1819-754X.
- Moreno, A., Pérez, A., Frigola, J., Nieto-Moreno, V., Rodrigo-Gámiz, M., Martrat, B., González-Sampériz, P., Morellón, M., Martín-Puertas, C., Corella, J.P., Belmonte, Á., Sancho, C., Cacho, I., Herrera, G., Canals, M., Grimalt, J.O., Jiménez-Espejo, F., Martínez-Ruiz, F., Vegas-Vilarrúbia, T., Valero-Garcés, B.L., 2012. The Medieval Climate Anomaly in the Iberian Peninsula reconstructed from marine and lake records. *Quaternary Sci. Rev.* 43, 16-32. doi: <https://doi.org/10.1016/j.quascirev.2012.04.007>.
- Murillo-Barroso, M., Martín-Torres, M., Massieu, M.D.C., Socas, D.M., González, F.M., 2017. Early metallurgy in SE Iberia. The workshop of Las Pilas (Mojácar, Almería, Spain) *Archaeol. Anthropol. Sci.* 9, 1–31. doi: <https://doi.org/10.1007/s12520-016-0451-8>.
- Nieto-Moreno, V., Martínez-Ruiz, F., Giralt, S., Jiménez-Espejo, F., Gallego-Torres, D., Rodrigo-Gámiz, M., García-Orellana, J., Ortega-Huertas, M., De Lange, G.J., 2011. Tracking climate variability in the western Mediterranean during the Late Holocene: a multiproxy approach. *Clim. Past* 7(1), 635-675. doi: <https://doi.org/10.5194/cp-7-1395-2011>.
- Nieto-Moreno, V., Martínez-Ruiz, F., Giralt, S., Gallego-Torres, D., García-Orellana, J., Masqué, P., Ortega-Huertas, M., 2013. Climate imprints during the ‘Medieval Climate Anomaly’ and the ‘Little Ice Age’ in marine records from the Alboran Sea basin. *Holocene* 23(9), 1227-1237. doi: <https://doi.org/10.1177/0959683613484613>.

- Nocete, F., Alex, E., Nieto, J.M., Sáez, R., Bayona, M.R., 2005. An archaeological approach to regional environmental pollution in the south-western Iberian Peninsula related to Third millennium BC mining and metallurgy. *J. Archaeol. Sci.* 32, 1566-1576. doi: <https://doi.org/10.1016/j.jas.2005.04.012>
- Nocete, F. 2006. The first specialised copper industry in the Iberian peninsula: Cabezo Juré (2900-2200 BC). *Antiquity*, 80(309), 646-657. doi: <https://doi.org/10.1017/S0003598X00094102>.
- Oliva, M., Ortiz, A. G., Palacios, D., Salvador-Franch, F., Salvà-Catarineu, M., 2014. Environmental evolution in Sierra Nevada (South Spain) since the Last Glaciation, based on multi-proxy records. *Quatern. Int.* 353, 195-209. doi: <https://doi.org/10.1016/j.quaint.2014.02.009>.
- Olivares, J., Barea-Azcón, J.M., Pérez-López, F.J., Tinaut, A., Henares, I., 2011. Las mariposas diurnas de Sierra Nevada. Consejería de Medio Ambiente, Junta de Andalucía. 512 pp.
- Olsen, J., Anderson, N.J., Knudsen, M.F., 2012. Variability of the North Atlantic Oscillation over the past 5,200 years. *Nature Geosci.* 5(11), 808-812. doi: <https://doi.org/10.1038/ngeo1589>.
- Pérez-Obiol, R., Jalut, G., Julià, R., Pélachs, A., Iriarte, M.J., Otto, T., Hernández-Beloqui, B., 2011. Mid-Holocene vegetation and climatic history of the Iberian Peninsula. *Holocene* 21, 75-93. doi: <https://doi.org/10.1177/0959683610384161>.
- Pini, R., Ravazzi, C., Raiteri, L., Guerreschi, A., Castellano, L., Comolli, R., 2017. From pristine forests to high-altitude pastures: an ecological approach to prehistoric human impact on vegetation and landscapes in the western Italian Alps. *J. Ecol.* 105(6), 1580-1597. doi: <https://doi.org/10.1111/1365-2745.12767>.
- Pons, A., Reille, M., 1988. The Holocene and Upper Pleistocene pollen record from Padul (Granada, Spain): a new study. *Palaeogeogr. Palaeoclim. Palaeoecol.* 66(3-4), 243-263. doi: [https://doi.org/10.1016/0031-0182\(88\)90202-7](https://doi.org/10.1016/0031-0182(88)90202-7).
- Postigo-Mijarra, J.M., Morla, C., Barrón, E., Morales-Molino, C., García Álvarez, S., 2010. Patterns of extinction and persistence of Arctotertiary flora in Iberia during the Quaternary. *Rev. Palaeobot. Palynol.* 162, 416-426. doi: <https://doi.org/10.1016/j.revpalbo.2010.02.015>.
- Ramos-Román, M.J., Jiménez-Moreno, G., Anderson, R.S., García-Alix, A., Toney, J.L., Jiménez-Espejo, F.J., Carrión, J.S. 2016. Centennial-scale vegetation and North Atlantic Oscillation changes during the Late Holocene in the southern Iberia. *Quaternary Sci. Rev.* 143, 84-95. doi: <https://doi.org/10.1016/j.quascirev.2016.05.007>.
- Ramos-Román, M.J., Jiménez-Moreno, G., Camuera, J., García-Alix, A., Anderson, R.S., Jiménez-Espejo, F.J., Carrión, J.S., 2018a. Holocene climate aridification trend and human impact interrupted by millennial and centennial-scale climate fluctuations from a new sedimentary record from Padul (Sierra Nevada, southern Iberian Peninsula). *Clim. Past* 14(1), 117. doi: <https://doi.org/10.5194/cp-14-117-2018>
- Ramos-Román, M.J., Jiménez-Moreno, G., Camuera, J., García-Alix, A., Anderson, R.S., Jiménez-Espejo, F.J., Sachse, D., Toney, J.L., Carrión, J.S., Webster, C., Yanes, Y.,

- 2018b. Millennial-scale cyclical environment and climate variability during the Holocene in the western Mediterranean region deduced from a new multi-proxy analysis from the Padul record (Sierra Nevada, Spain). *Global Planet. Change* 168, 35-53. doi: <https://doi.org/10.1016/j.gloplacha.2018.06.003>.
- Regato, P., Salman, R., 2008. *Mediterranean Mountains in a Changing World: Guidelines for developing action plans*. IUCN.
- Reille, M., 1999. *Pollen et Spores d'Europe et d'Afrique du Nord*. 2nd Edition. Marseille: Laboratoire de Botanique Historique et Palynologie.
- Roberts, C. N., Woodbridge, J., Palmisano, A., Bevan, A., Fyfe, R., Shennan, S., 2019. Mediterranean landscape change during the Holocene: Synthesis, comparison and regional trends in population, land cover and climate. *Holocene* 29(5), 923-937. doi: <https://doi.org/10.1177/0959683619826697>.
- Rodríguez Ariza, M.O., Esquivel, J., 1989. Una aplicación del análisis de correspondencias en la valoración del antracoanálisis de Los Millares. *Cuadernos de Prehistoria y Arqueología* 14, 81-108 Universidad de Granada.
- Rodríguez Ariza, M.O., 1992. Human-plant relationships during the Copper and Bronze Ages in the Baza and Guadix basins (Granada, Spain), *Bulletin de la Société Botanique de France. Actualités Botaniques* 139:2-4, 451-464, doi: <https://doi.org/10.1080/01811789.1992.10827120>
- Rodríguez-Ariza, M.O., 2000. El paisaje vegetal de la Depresión de Vera durante la Prehistoria reciente. Una aproximación desde la antracología. *Trabajos de Prehistoria*, 57, 145–156. doi: <https://doi.org/10.3989/tp.2000.v57.i1.266>.
- Rodríguez-Ariza, M. O., Montes Moya, E., 2005. On the origin and domestication of *Olea europaea* L. (olive) in Andalucía, Spain, based on the biogeographical distribution of its finds. *Veg. Hist. Archaeobot.* 14(4), 551-561. doi: <https://doi.org/10.1007/s00334-005-0012-z>.
- Romero Funes, C., 2016. El Libro de Apeo y repartimiento del lugar de Carataunas: un ejemplo de la repoblación de la Alpujarra después del levantamiento de los Moriscos, a finales del siglo XVI. *Estudios sobre patrimonio, cultura y ciencias medievales*, 18(3): 1083-1164.
- Rubio de Lucas, J.L., Albert Gamboa, M.J., Muñoz Municio, M.C., San José Gómez, S., 1995. *Cuadernos de la Trashumancia*. Sierra Nevada. Madrid. ICONA, 64 pp.
- Ruddiman, W. 2017. Geographic evidence of the early anthropogenic hypothesis. *Anthropocene*, 20: 4-14. doi: <https://doi.org/10.1016/j.ancene.2017.11.003>.
- Ruiz, A., Rodríguez Ariza, M.O., 2002. Paisaje y asentamiento entre los íberos de la Cuenca del Guadalquivir. Universidad de Jaén. Jaén.
- Salomón, R., Rodríguez-Calcerrada, J., González-Doncel, I., Gil, L., Valbuena-Carabaña, M., 2017. On the general failure of coppice conversion into high forest in *Quercus pyrenaica* stands: a genetic and physiological approach. *Folia Geobot.* 52(1), 101-112. doi: <https://doi.org/10.1007/s12224-016-9257-9>.

- Salvador Oyonate, J.A., 2012. Bastetania (I): Desde la Conquista Romana hasta las Invasiones Bárbaras (208 AC–411). Una propuesta. *Péndulo. Papeles de Bastitania* 13, 9-42.
- Salvador Ventura, F., 2012. El centro de Andalucía entre los siglos V y VII: de las ciudades herederas de Roma a las sedes episcopales hispanovisigodas. *Habis* 43, 233-247.
- Sayadi S., Calatrava, J., 2001. Análisis funcional de los sistemas agrarios para el desarrollo rural sostenible: Las funciones productiva, recreativa y estética de la agricultura en la alta Alpujarra. Ministerio de Agricultura, Pesca y Alimentación, Madrid.
- Schirrmacher, J., Kneisel, J., Knitter, D., Hamer, W., Hinz, M., Schneider, R.R., Weinelt, M., 2020. Spatial patterns of temperature, precipitation, and settlement dynamics on the Iberian Peninsula during the Chalcolithic and the Bronze Age. *Quaternary Sci. Rev.* 233, 106220. doi: <https://doi.org/10.1016/j.quascirev.2020.106220>.
- Smith, B.D., Zeder, M.A., 2013. The onset of the Anthropocene. *Anthropocene* 4, 8-13. doi: <https://doi.org/10.1016/j.ancene.2013.05.001>.
- Stuiver, M., Reimer, P.J., Reimer, R.W., 2019. CALIB 7.1 [WWW program] at <http://calib.org>.
- Taberlet, P., Cheddadi, R., 2002. Quaternary refugia and persistence of biodiversity. *Science* 297(5589), 2009-2010. doi: [10.1126/science.297.5589.2009](https://doi.org/10.1126/science.297.5589.2009).
- Trillo San José, C., 2004. Agua, tierra y hombres en Al-Andalus: la dimensión agrícola del mundo nazarí (Vol. 1). *Ajbar Colección de Historia. Motril* (Granada).
- Tzedakis, P.C., Lawson, I.T., Frogley, M.R., Hewitt, G.M., Preece, R.C., 2002. Buffered Tree Population Changes in a Quaternary Refugium: Evolutionary Implications. *Science*, 297: 2044-2047. doi: [10.1126/science.1073083](https://doi.org/10.1126/science.1073083).
- Valbuena-Carabaña, M., de Heredia, U.L., Fuentes-Utrilla, P., González-Doncel, I., Gil, L., 2010. Historical and recent changes in the Spanish forests: a socio-economic process. *Rev. Palaeobot. Palynol.* 162(3), 492-506. doi: <https://doi.org/10.1016/j.revpalbo.2009.11.003>.
- Valbuena-Carabaña, M., Gil, L., 2013. Genetic resilience in a historically profited root sprouting oak (*Quercus pyrenaica* Willd.) at its southern boundary. *Tree Genet Genomes* 9(5), 1129-1142. doi: <https://doi.org/10.1007/s11295-013-0614-z>.
- Van Geel, B., 2001. Non-pollen palynomorphs. In: Smol, J.P., Last, W.M. and Birks, J.B. (eds) *Tracking Environmental Change Using Lake Sediments*, vol. 3. Dordrecht: Springer Netherlands, pp. 99–119.
- Vogiatzakis, I., 2012. *Mediterranean mountain environments*. John Wiley & Sons. doi: [10.1002/9781119941156](https://doi.org/10.1002/9781119941156).
- Walker, M.J.C., Berkelhammer, M., Björck, S., Cwynar, L.C., Fisher, D.A., Long, A.J., Lowe, J.J., Newnham, R.M., Rasmussen, S.O., Weiss, H., 2012. Formal subdivision of the Holocene Series/Epoch: a Discussion Paper by a Working Group of INTIMATE (Integration of ice-core, marine and terrestrial records) and the Subcommittee on Quaternary Stratigraphy (International Commission on Stratigraphy). *J. Quaternary Sci.* 27(7), 649-659. doi: <https://doi.org/10.1002/jqs.2565>.

- Whitlock, C., Colombaroli, D., Conedera, M., Tinner, W. 2018. Land-use history as a guide for forest conservation and management. *Conserv. Biol.* 32(1), 84-97. doi: <https://doi.org/10.1111/cobi.12960>
- Woodward, J., 2009. *The physical geography of the Mediterranean*. Oxford University Press.
- Zamora Rodríguez, R.J., Pérez-Luque, A.J., Bonet, F.J., Barea-Azcón, J. M., Aspizua, R. 2015. La huella del cambio global en Sierra Nevada: Retos para la conservación. *Consejería de Medio Ambiente y Ordenación del Territorio. Junta de Andalucía*. 208 pp
- Zielhofer, C., Köhler, A., Mischke, S., Benkaddour, A., Mikdad, A., Fletcher, W.J., 2019. Western Mediterranean hydro-climatic consequences of Holocene ice-rafted debris (Bond) events. *Clim. Past* 15 (2), 463-475. <https://doi.org/10.5194/cp-15-463-2019>.
- Zoido Naranjo, F., Jiménez Olivencia, Y., 2015. *Catálogo de paisajes de la provincia de Granada*. *Consejería de Medio Ambiente y Ordenación del Territorio, Sevilla*.