

# Benefits of Physical Activity for Low Back Pain Across Different Age Groups

Carmen Elena Jiménez-Gutiérrez<sup>1</sup>, Pablo Redruello-Guerrero<sup>2</sup>

<sup>1</sup> Faculty of Nursing, University of Granada (UGR)

<sup>2</sup> Faculty of Medicine, University of Granada (UGR)

## Abstract

Low back pain (LBP) is one of the most common and incapacitating pain conditions globally. Its current treatment is essentially pharmacological, leading to a large consumption of painkillers and opioids. The present non-systematic review collects data from various clinical trials evaluating the effects of physical activity in low back pain management and organizes them into different age groups. Early treatment in adolescents based on physical activity combined with spinal manipulative therapy shows a decrease in chronic low back pain. In adults, various combination therapies have been reviewed with mixed results. Lumbar stimulation combined with activities such as walking has been found to be the most significant one. Physical activity shows great benefits in the elderly population, because it reduces the intake of painkillers. These findings stress the value of exercise as an alternative to pharmacological treatment in low back pain management.

**Keywords:** low back pain, physical activity, adolescents, adults, elderly.

## 1. Introduction

Low back pain (LBP) affects the lumbar region of the spine. Though it can range in intensity, LBP is considered to be one of the most common pain conditions, with an estimated 40 to 80% of individuals worldwide experiencing it at some point in their lives (1, 2). In the past two decades, there has been a 42% increase in limitations due to LBP, thereby establishing itself as the first cause of disability worldwide (3).

The high prevalence of LBP and its association to disability and deterioration in quality of life make it a public health issue (4). It imposes a high cost on society, with sick leaves due to LBP showing a similar incidence to leaves due to pathologies such as diabetes mellitus or coronary heart disease (5).

There are various forms of LBP, such as mechanical low back pain, which affects spine, joints, and muscles (6), and non-specific low back pain, which is not attributable to a known cause (7). Figure 1 illustrates both types of pathologies.

The role of cognitive and behavioral factors in the development and management of chronic pain has recently been emphasised. Chronic pain causes a cortical reorganization which sensitises the neural network that subserves pain and disinhibits the surrounding neural networks. Thus, the modifica-

tion of pain-eliciting and maintaining behaviors, cognitions and emotions has the potential to increase pain management and tolerance (8). Furthermore, physical activity and exercise can have an effect on these neural networks, which leads to a decrease in chronic pain (9).

In collecting the findings of various clinical trials, the present narrative review aims to examine the benefits of physical activity to reduce LBP, stratified by age.

## 2. Adolescents

Low back pain most often begins in adolescence (10-12), which is considered a strong predictor for LBP in adulthood. Adult pain levels appear to be reached by around the age of 18 (13-15). Rehabilitation exercises should be fostered in order to help patients manage LBP and prevent future relapses (16). Similarly, spinal manipulative therapy (SMT) combined with exercise therapy (ET) is advised before pharmacological treatment, as it has proven to be effective for low back pain (17). Encouraging patients to remain active is also key in LBP treatment in order to prevent long-term sitting, which is known to exacerbate symptoms (18). Aerobic exercise is the most recommended activity to this end (19). Few studies documenting the effectiveness of physical exercise for LBP have been conducted on children and adolescents (20-22).

### TRANSLATED BY:

Carmen Bohórquez-Gavira<sup>3</sup>, Sandra Gutiérrez-Bullón<sup>3</sup>, Javier León-Saniger<sup>3</sup>, María Pineda-Cantos<sup>3</sup>, Luís Manuel Porrero-Triguero<sup>3</sup>, Javier Tarín-Marín<sup>3</sup>

<sup>3</sup> Faculty of Translation and Interpreting, University of Granada (UGR)

In a randomized trial with a sample of 185 adolescents aged 12-18 years, the effectiveness of spinal manipulative therapy combined with exercise therapy was compared to exercise therapy alone. The first option involved spinal manipulation and mobilization, while the second required the teaching and supervision of exercises by trained personnel, as well as the unsupervised performance of the same exercises at home. Participants were provided with instructions on how to perform the physical exercises and which posture to adopt while exercising, sitting down, studying, etc. Spinal manipulation combined with exercise for over 6 months reported an increase in the functional capacity of the individuals, as shown in Graph 1. Both approaches led participants to experience a long-term decrease in LBP intensity. It is also noteworthy that adolescents reported an 80% reduction in medication use. These are important findings in light of the concern that exists around the safety and effectiveness of medication use for managing LBP (23).

One of the most common forms of LBP in adolescents is directly linked to the exercises in elite gymnastics which overstrain the lumbar spine (24). A clinical trial was conducted on a 15-year-old female gymnast with extension-based LBP. The movements overloading her lumbar spine were reeducated thanks to specific activities using a spine stabilizer and manual therapy technique. It was found that the addressing of cognitive-affective factors together with the correction of maladaptive exercises reduced the nociceptive input and desensitized the nervous system, leading to a better control of LBP (25). Another clinical trial developed a programme of exercises designed to help prevent low back injuries and reduce pain. Thirty female athletes, 10-14 years old, participated in this study. Fifteen gymnasts implemented back stretches and various exercises to improve posture and coordination in their ordinary training, while the other fifteen followed the standard training, acting as the control group. After the intervention, assessment showed a decrease of pain identified as mild (12%) or moderate (11%) and a disappearance of severe pain (26).

### 3. Adults

Clinical prediction rules analyse the feasible evolution of patients with LBP without surgical intervention (27). PERRON et al. carried out a study during a month and a half with 85 soldiers suffering from subacute or chronic low back pain and without surgical intervention. This study showed that there are a series of variables which can predict the future evolution of patients before the physical activity is performed. These variables are included in Table 1.

Of the subjects who presented between four and five variables, 77.5% obtained a favorable outcome (28). The purpose of the authors was to establish a new clinical prediction rule that would help identify the prognosis of these patients depending on the variables considered in the study.

#### 3.1. Conventional exercises

On the one hand, the aim of lumbar stabilization exercises (SE) is to strengthen the muscles in charge of stabilizing the spine (29). These exercises should be adapted to the clinical characteristics of each patient (30), which can be done by using individualised graded lumbar SE (IGLSE), since this technique permits to adapt the intensity of the exercise. HYUN SUH et al. carried out a prospective, randomized, controlled study in order to evaluate walking efficiency (WE) and put into practice IGLSE with a sample of 48 patients with LBP. The study showed that lumbar SE and WE significantly improved LBP (31).

On the other hand, promotion of physical activity is essential for a good development and for its practice (32). The Movement Coaching is an intervention which comprises three different components, (33) such as physical exercise in the same place and time with a therapist, which is very effective (34), and also includes telephone and internet-based aftercare. SCHALLER et al. performed a randomized controlled trial in Germany with 144 patients with LBP. The sample was divided into two groups of 71 and 73 patients respectively and they compared a multi-component intervention with Movement Coaching and a control intervention with online presentations without coaching. Their results were not significantly relevant and they did not prove that the suggested therapy was more effective. This could be due to a decrease in the total physical activity during the 12-months follow-up (33).

In addition, home exercise programmes have also been developed. These consist of 10 minutes of aerobic activity followed by eight types of ground exercises, which were aimed at strengthening the lumbar muscles, and five types of muscle stretches (35). These interventions, which appear in Figure 2, were carried out by two groups of 13 and 17 patients with LBP respectively. Only the first group was supervised by a physical therapist each week. Both groups experienced a symptomatic improvement in LBP, but the differences between the supervised and the unsupervised group were not significant (35).

#### 3.2. Balneotherapy

The combined therapy of physical exercise and balneotherapy was approached in order to allow workers with this chronic LBP to return to work. For

this purpose, a prospective randomized controlled trial was performed (36) using a modified Zelen design (37). This was carried out with a sample of 88 patients in France. The patients were divided into two parallel groups of 45 and 43 subjects respectively. The first group received a 5-day intensive intervention, which consisted of balneotherapy (2 hr/day), physical exercise (45 min/day) and an individualized educational program (45 min/day). On the contrary, the second group was taken as a control group and only received regular medical supervision. The findings were promising, but there was no significant evidence because the size of the sample was insufficient (36). These results leave the door open for future clinical trials in this area.

On the other hand, HUBER et al. carried out a randomized controlled trial with 80 Austrian patients. They were divided into three groups: the first group (27 patients) performed green exercise, the second one (26 patients) combined green exercise with balneotherapy, and the third one (27 patients) was the control group and did not spend 8 days in the Tyrol. With the second group, mountain hiking and a spa treatment with Mg-Ca-SO<sub>4</sub> thermal water were used in order to evaluate the symptomatic improvement and the spinal mobility. The results of the study showed that this combined therapy has benefits for the participants concerning pain, functional mobility and subjective quality of life (38).

### 3.3. Alternative Therapies

Tai Chi (TC) is a health-promoting exercise (39) that can be performed by people from different age groups (40). In addition, it shows benefits in multiple pathologies (41). It aims to integrate mind-body development to improve or maintain the health state (42). A randomized controlled trial was conducted on 43 patients with LBP to learn about the effectiveness of TC. They were divided into three groups, the first one of 15 people who were treated with TC exercises; the second one of 15 patients who performed core stabilization exercises; and the last group of 13 patients who were taken as control group (43). The results showed that TC had positive effects on pain, but did not improve the proprioception on lower limbs, which were also assessed.

Likewise, yoga is another alternative therapy that seeks to alleviate the symptoms and pain of multiple conditions (44). NEYAZ et al. conducted a prospective randomized comparative study to determine whether Hatha yoga (HY) therapy was more effective than conventional therapeutic exercises (CTEs) in patients with chronic low back pain. The Hatha yoga intervention includes a series of physical, breathing and meditation exercises. The sample size was of 70 subjects, distributed in two equal

groups that were treated with both techniques. The Hatha yoga therapy consisted of six sessions (35 min/week). Pain intensity decreased significantly in both groups with no evidence of improvement in one group over the other (45).

## 4. Elderly

The world's population is aging rapidly and it is estimated that between 76 and 82% of elderly patients suffer from some form of non-cancer pain (46, 47). LBP is common in older adults, resulting in physical limitations, disability, and decreased quality of life. The prevalence of this form of pain is estimated to affect between 32 and 58% of the elderly population (48, 49). Many of them are undergoing opioid and analgesic treatment in order to manage their chronic pain. This highlights the need to identify other safe non-pharmacological strategies for pain management in the elderly. Some studies have shown the advantages of physical exercise or spinal manipulative therapy to manage LBP (50, 51). Promoting pain management programs to learn how to manage pain in the long term has also been found to be effective (52).

A randomised clinical trial conducted on 241 people over 65 years old with LBP assessed the benefits of adding spinal manipulative therapy or supervised rehabilitation exercises to the exercise performed at home for 12 weeks. Three groups were created for this purpose. The first group exercised only at home, the second combined it with a supervised exercise program and the third with spinal manipulative therapy. The reduction of pain intensity was not very significant in the combined therapy of spinal manipulation and home exercise compared to physical exercise alone, increasing the improvement by only 10% as shown in Graph 2. It was considered more cost-effective to develop a multi-session home exercise program and in patients requiring more support to perform spinal manipulative therapy or supervised exercise (53).

Another randomized clinical trial was conducted with 60 seniors in their 70s. Home strengthening, stretching and aerobic activity interventions were performed and followed up by telephone. It was concluded that this method was effective and valid to increase adherence to exercise programs in older patients with LBP (54).

New approaches to LBP treatments include promoting healthy habits and techniques for pain management, and reducing interventions to treat pain symptoms by reducing the high consumption of medications and medical interventions (55). Physical exercise is a simple activity that can be performed at home (56). Using telecommunication for pa-



tient follow-up is a novel approach in medicine that has advantages for older patients as they can access health care comfortably from home (57).

## 5. Conclusion

Several clinical trials have shown the benefits of physical activity in reducing LBP. Spinal manipulation, along with the development of physical exercise has been found to relieve low back pain in adolescents. In addition, in this age group, preventive compensatory training has also been identified as relieving pain in elite gymnasts in the short term. Various combination therapies have been reviewed for the adult population. Balneotherapy, alternative therapies such as Tai Chi or Hatha Yoga, or spinal stimulation have been highlighted; all of them accompanied by physical exercise. Spinal manipulation has shown a more relevant symptomatic improvement. Regarding the elderly population, the improvement is not as significant in terms of intensity. However, it favors the decrease of painkiller intake and pain management by the patient. Furthermore, in this age group, telephone follow-up increased adherence to treatment, unlike other age groups.

This review suggests that physical exercise helps to lower LBP. In both adolescents and adults, it would be advisable to practice physical exercise along with other therapies to achieve a more significant pain relief. Moreover, telephone follow-up is recommended in the elderly to influence the monitoring of suggested physical activities.

However, there is still not enough evidence available, so more clinical trials on LBP and exercise can be suggested in all age groups. This limitation is present mainly in adolescents, so studies regarding this age range could increase the scientific evidence in this field.

## Statements

### Acknowledgements

This paper is part of the Teaching Innovation Project coordinated between the Faculty of Medicine and the Faculty of Translation and Interpreting of the University of Granada (UGR), within the framework of the FIDO Plan 2018-2020 of the UGR (code 563).

### Ethical concerns

This paper did not require the approval of any ethics committee.

### Conflicts of interest

The authors of this paper declare no conflicts of interest.

## Funding

No funding was received for the production of this paper.

## References

1. Frymoyer JW. Back Pain and Sciatica. *New England Journal of Medicine*. 1988 4;318(5):291-300.
2. Hoy D, Bain C, Williams G, March L, Brooks P, Blyth F, et al. A systematic review of the global prevalence of low back pain. *Arthritis Rheum*. 2012; 64(6):2028-37.
3. Hoy D, March L, Brooks P, Blyth F, Woolf A, Bain C, et al. The global burden of low back pain: estimates from the Global Burden of Disease 2010 study. *Ann Rheum Dis*. 2014; 73(6):968-74.
4. Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, et al. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012 15;380(9859):2163-96.
5. Druss BG, Rosenheck RA, Sledge WH. Health and disability costs of depressive illness in a major U.S. corporation. *Am J Psychiatry*. 2000; 157(8):1274-8.
6. Will JS, Bury DC, Miller JA. Mechanical Low Back Pain. *Am Fam Physician*. 2018; 98(7):421-8.
7. Russo M, Deckers K, Eldabe S, Kiesel K, Gilligan C, Vicecili J, et al. Muscle Control and Non-specific Chronic Low Back Pain. *Neuro-modulation*. 2018; 21(1):1-9.
8. Moseley GL, Flor H. Targeting Cortical Representations in the Treatment of Chronic Pain: A Review. *Neurorehabil Neural Repair*. 2012; 26(6):646-52.
9. Morley, S, Eccleston, C, Williams, A. Revisión sistemática y meta-análisis de ensayos controlados aleatorios de terapia cognitiva conductual y terapia conductual para el dolor crónico en adultos, excluyendo dolor de cabeza. *Dolor*. 1999 ; 80: 1 - 13.
10. Calvo-Muñoz I, Gómez-Conesa A, Sánchez-Meca J. Physical therapy treatments for low back pain in children and adolescents: a meta-analysis. *BMC Musculoskelet Disord*. 2013; 14:55.
11. Jeffries LJ, Milanese SF, Grimmer-Somers KA. Epidemiology of adolescent spinal pain: a systematic overview of the research literature. *Spine*. 2007; 32(23):2630-7.
12. Leboeuf-Yde C, Kyvik KO. At what age does low back pain become a common problem? A study of 29,424 individuals aged 12-41 years. *Spine*. 1998; 23(2):228-34.
13. Dunn KM, Hestbaek L, Cassidy JD. Low back pain across the life course. *Best Practice & Research Clinical Rheumatology*. 2013; 27(5):591-600.
14. Hestbaek L, Leboeuf-Yde C, Kyvik KO, Manniche C. The course of low back pain from adolescence to adulthood: eight-year follow-up of 9600 twins. *Spine*. 2006; 31(4):468-72.
15. Jeffries LJ, Milanese SF, Grimmer-Somers KA. Epidemiology of adolescent spinal pain: a systematic overview of the research literature. *Spine*. 2007; 32(23):2630-7.
16. Delitto A, George SZ, Van Dillen L, Whitman JM, Sowa G, Shekelle P, et al. Low back pain. *J Orthop Sports Phys Ther*. 2012;42(4):A1-57.
17. Qaseem A, Wilt TJ, McLean RM, Forcica MA, Clinical Guidelines Committee of the American College of Physicians. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med*. 2017; 166(7):514-30.
18. Chou R, Qaseem A, Snow V, Casey D, Cross JT, Shekelle P, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med*. 2007; 147(7):478-91.
19. Maher C, Underwood M, Buchbinder R. Non-specific low back pain. *Lancet*. 2017 18;389(10070):736-47.
20. Ahlqwist A, Hagman M, Kjellby-Wendt G, Beckung E. Physical therapy treatment of back complaints on children and adolescents. *Spine*. 2008; 33(20):E721-727.
21. Fanucchi GL, Stewart A, Jordaan R, Becker P. Exercise reduces the intensity and prevalence of low back pain in 12-13 year old children: a randomised trial. *Aust J Physiother*. 2009;55(2):97-104.
22. Jones M, Stratton G, Reilly T, Unnithan V. The efficacy of exercise as an intervention to treat recurrent nonspecific low back pain in adolescents. *Pediatr Exerc Sci*. 2007; 19(3):349-59.
23. Evans R, Haas M, Schulz C, Leininger B, Hanson L, Bronfort G. Spinal manipulation and exercise for low back pain in adoles-

- cents: a randomized trial. *Pain*. 2018; 159(7):1297–307.
24. Hutchinson MR. Low back pain in elite rhythmic gymnasts. *Med Sci Sports Exerc*. 1999; 31(11):1686–8.
  25. Winslow JJ, Jackson M, Getzin A, Costello M. Rehabilitation of a Young Athlete With Extension-Based Low Back Pain Addressing Motor-Control Impairments and Central Sensitization. *J Athl Train*. 2018; 53(2):168–73.
  26. Marini M, Sgambati E, Barni E, Piazza M, Monaci M. Pain syndromes in competitive elite level female artistic gymnasts. Role of specific preventive-compensative activity. *Ital J Anat Embryol* 2008;113(1):47–54.
  27. Haskins R, Osmotherly PG, Rivett DA. Validation and impact analysis of prognostic clinical prediction rules for low back pain is needed: a systematic review. *J Clin Epidemiol*. 2015; 68(7):821–32.
  28. Perron M, Gendron C, Langevin P, Leblond J, Roos M, Roy J. Prognostic factors of a favorable outcome following a supervised exercise program for soldiers with sub-acute and chronic low back pain. *BMC musculoskeletal disorders* 2018 Apr 2;19(1):95.
  29. Lee HS, Kim DJ, Oh Y, Min K, Ryu JS. The effect of individualized gradable stabilization exercises in patients with chronic low back pain: Case-control study. *J Back Musculoskelet Rehabil*. 2016; 29(3):603–10.
  30. Kim CR, Park DK, Lee ST, Ryu JS. Electromyographic Changes in Trunk Muscles During Graded Lumbar Stabilization Exercises. *PM R*. 2016;8(10):979–89.
  31. Suh JH, Kim H, Jung GP, Ko JY, Ryu JS. The effect of lumbar stabilization and walking exercises on chronic low back pain. *Medicine (Baltimore)* [Internet]. 2019 [cited 2020 Mar 8];98(26). Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6616307/>
  32. Müller-Riemenschneider F, Reinhold T, Nocon M, Willich SN. Long-term effectiveness of interventions promoting physical activity: a systematic review. *Prev Med*. 2008; 47(4):354–68.
  33. Schaller A, Petrowski K, Pfoertner T, Froboese I. Effectiveness of a theory-based multicomponent intervention (Movement Coaching) on the promotion of total and domain-specific physical activity: a randomised controlled trial in low back pain patients. *BMC musculoskeletal disorders* 2017 Nov 6;18(1):431.
  34. Vuori IM, Lavie CJ, Blair SN. Physical activity promotion in the health care system. *Mayo Clin Proc*. 2013; 88(12):1446–61.
  35. Kanas M, Faria RS, Salles LG, Sorpreso ICE, Martins DE, Cunha RA da, et al. Home-based exercise therapy for treating non-specific chronic low back pain. *Revista da Associação Médica Brasileira*. 2018; 64(9):824–31.
  36. Nguyen C, Boutron I, Rein C, Baron G, Sanchez K, Palazzo C, et al. Intensive spa and exercise therapy program for returning to work for low back pain patients: a randomized controlled trial. *Scientific reports* 2017 Dec 20;7(1):17956–10.
  37. Zelen M. A new design for randomized clinical trials. *N Engl J Med*. 1979; 300(22):1242–5.
  38. Huber D, Grafetstätter C, Proßegger J, Pichler C, Wöll E, Fischer M, et al. Green exercise and mg-ca-SO<sub>4</sub> thermal balneotherapy for the treatment of non-specific chronic low back pain: a randomized controlled clinical trial. *BMC Musculoskelet Disord* [Internet]. 2019 [Last access: 8 March 2020];20. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6524239/>
  39. Liu J, Xie H, Liu M, Wang Z, Zou L, Yeung AS, et al. The Effects of Tai Chi on Heart Rate Variability in Older Chinese Individuals with Depression. *Int J Environ Res Public Health*. 2018 07;15(12).
  40. Zou L, Wang H, Xiao Z, Fang Q, Zhang M, Li T, et al. Tai chi for health benefits in patients with multiple sclerosis: A systematic review. *PLoS ONE*. 2017;12(2):e0170212.
  41. Huston P, McFarlane B. Health benefits of tai chi: What is the evidence? *Canadian Family Physician*. 2016; 62(11):881–90.
  42. Webster CS, Luo AY, Krägeloh C, Moir F, Henning M. A systematic review of the health benefits of Tai Chi for students in higher education. *Prev Med Rep*. 2016; 3:103–12.
  43. Liu J, Yeung A, Xiao T, Tian X, Kong Z, Zou L, et al. Chen-Style Tai Chi for Individuals (Aged 50 Years Old or Above) with Chronic Non-Specific Low Back Pain: A Randomized Controlled Trial. *International journal of environmental research and public health* 2019 Feb 12;16(3):517.
  44. Wieland LS, Skoetz N, Pilkington K, Vempati R, D'Adamo CR, Berman BM. Yoga treatment for chronic non-specific low back pain. *The Cochrane database of systematic reviews* 2017 Jan 12;1(1):CD010671.
  45. Neyaz O, Sumila L, Nanda S, Wadhwa S. Effectiveness of Hatha Yoga Versus Conventional Therapeutic Exercises for Chronic Nonspecific Low-Back Pain. *The Journal of Alternative and Complementary Medicine*. 2019; 25(9):938–45.
  46. Guidance on the management of pain in older people. *Age and Ageing*. 2013; 42(suppl\_1):i1–57.
  47. Horgas AL, Snigurska U, Farland MZ, Marsiske M. Analyzing Analgesic Medications in Community-Dwelling Older Adults. *Pain Med*. 2019; 20(1):58–67.
  48. Cecchi F, Debolini P, Lova RM, Macchi C, Bandinelli S, Bartali B, et al. Epidemiology of Back Pain in a Representative Cohort of Italian Persons 65 Years of Age and Older: The InCHIANTI Study. *Spine*. 2006; 31(10):1149–1155.
  49. Jacobs JM, Hammerman-Rozenberg R, Cohen A, Stessman J. Chronic Back Pain Among the Elderly: Prevalence, Associations, and Predictors. *Spine*. 2006; 31(7):E203.
  50. Chou R, Deyo R, Friedly J, Skelly A, Hashimoto R, Weimer M, et al. Noninvasive Treatments for Low Back Pain [Internet]. Agency for Healthcare Research and Quality (US); 2016 [Last access: 8 March 2020]. Available at: <https://www.ncbi.nlm.nih.gov/sites/books/NBK350276/>
  51. Skelly AC, Chou R, Dettori JR, Turner JA, Friedly JL, Rundell SD, et al. Noninvasive Nonpharmacological Treatment for Chronic Pain: A Systematic Review [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2018 [cited 2020 Mar 8]. (AHRQ Comparative Effectiveness Reviews). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK519953/>
  52. Mansell G, Hall A, Toomey E. Behaviour change and self-management interventions in persistent low back pain. *Best Practice & Research Clinical Rheumatology*. 2016; 30(6):994–1002.
  53. Schulz C, Evans R, Maiers M, Schulz K, Leininger B, Bronfort G. Spinal manipulative therapy and exercise for older adults with chronic low back pain: a randomized clinical trial. *Chiropr Man Therap*. 2019; 27:21.
  54. Goode AP, Taylor SS, Hastings SN, Stanwyck C, Coffman CJ, Allen KD. Effects of a Home-Based Telephone-Supported Physical Activity Program for Older Adult Veterans With Chronic Low Back Pain. *Phys Ther*. 2018; 98(5):369–80.
  55. Foster NE, Anema JR, Cherkin D, Chou R, Cohen SP, Gross DP, et al. Prevention and treatment of low back pain: evidence, challenges, and promising directions. *The Lancet*. 2018; 391(10137):2368–83.
  56. Kelley GA, Kelley KS, Hootman JM, Jones DL. Exercise and Health-Related Quality of Life in Older Community-Dwelling Adults: A Meta-Analysis of Randomized Controlled Trials. *J Appl Gerontol*. 2009; 28(3):369–94.
  57. Narasimha S, Agnisarman S, Chalil Madathil K, Gramopadhye A, McElligott JT. Designing Home-Based Telemedicine Systems for the Geriatric Population: An Empirical Study. *Telemed J E Health*. 2018; 24(2):94–110.

## Annex I: Figures

Figure 1. Cases of low back pain (LBP).




Figure 2. Table showing the exercise program to be developed at home, for strength and stretching. Image adapted from Kanas M et al. (35)

AEROBICS: WALKING OR STATIONARY BIKE FOR 10 MINUTES

### STRETCHES

**1**

3X30 seconds

While lying supporting your back, cross one leg over the other. Using both hands, hold your thigh from the back and pull your legs towards your chest until you feel the crossed leg is being stretched. Hold the position for 30 seconds. You should feel like your stretching, with no pain. While stretching, breathe naturally. Repeat each movement 03 times for each leg. You should feel the stretching on the posterior muscles of the leg that is crossed.

**2**

3X30 seconds

While lying with your back on the ground, hold one of your legs, pull it towards your chest, and hold it there for 30 seconds. The other leg should remain stretched. You should feel like your stretching, with no pain. While stretching, breathe naturally. Repeat this movement 03 times for each leg. You should feel the stretching in the posterior muscles of the bent leg.

**3**

3X30 seconds

While sitting on a firm surface, flex one leg and stretch the other while bringing your hands towards the foot of the stretched leg. If you can't touch your foot, just keep your hands towards it. Try to relax your back and neck and hold for 30 seconds. You should feel like your stretching, with no pain. While stretching, breathe naturally. Repeat this movement 03 times for each leg. You should feel the stretching in the posterior muscles of the stretched leg.

**4**

3X30 seconds

Sit on your heels, stretch your arms as forward as possible, relax your back and neck and hold for 30 seconds. You should feel like your stretching, with no pain. While stretching, breathe naturally. Repeat the stretching 03 times. You should feel the stretching on the posterior muscles of your back.

**5**

3X30 seconds

Standing up, use your hand to pull one of your feet towards your glutes and hold for 30 seconds. Remember to use your free hand to support yourself and keep your back as straight as possible. You should feel like your stretching, with no pain. While stretching, breathe naturally. Repeat each movement 03 times for each leg. You should feel the stretching in the anterior muscles of the bent leg.

### MUSCLE-RECRUITMENT

**1**

10X10 seconds

Lying down with your back well-supported, bend your knees and keep your shoulders relaxed. Using your abdomen muscles (without putting too much strength in them), pull in your navel as if trying to make it touch the ground. Once it is pulled in, hold the position for 10 seconds. Repeat the movement 10 times a day.

**2**

3x30 seconds (15 seconds min.)

Lie on your stomach with your forearms resting on the ground and your elbows well below your shoulders. Carry out exercise 01 (pull navel in) and lift your knees off the floor until they are straight. Keep your back straight and hold that position for 15-30 seconds (as long as you can). If it is too difficult, keep your knees on the ground and just raise your hips, keeping your back straight. It is important that you maintain your abdomen contracted and your navel pulled in for the entire duration of the exercise. Repeat the movement 03 times.

**3**

Inicio

Final

3x30 seconds (15 seconds min.)

On a surface that is comfortable for your knees, position yourself on your hands and knees, keeping your back straight. Carry out exercise 01 (pull navel in) and then straighten the left knee up to your hips and the right arm up to your shoulders. Hold the position for 15-30 seconds (as long as you can). Return to the initial position. Straighten the right knee and the left arm and hold for another 15-30 seconds. It is important that you maintain your abdomen contracted and your navel pulled in for the entire duration of the exercise. Repeat the exercise 03 times.

**4**

3x15 repetitions

While lying with your back well-supported, bend your knees. Place a soft ball or pillow between your legs and press it while, at the same time, tightening the pelvic muscles, as if trying to "hold in pee." It is important to "hold in pee" every time you press the ball between your knees. Carry out 03 series of 15 repetitions.

**5**

3x30 seconds (15 seconds min.)

Lie on your back with your knees bent. Carry out exercise 01 (pull navel in) and raise your hips until your back is straight, forming a line between your knees and shoulders. Hold the position for 15-30 seconds (as long as you can). It is important that you maintain your abdomen contracted and your navel pulled in for the entire duration of the exercise. Repeat the movement 03 times.

**6**

3x30 seconds (15 seconds min.)

Lying on your side with your elbow aligned with your shoulder and your forearm resting on the ground, carry out exercise 01 (pull navel in) and raise your hips until your body is well stretched, forming a line from your feet to your shoulder. Hold that position for 15-30 seconds (as long as you can). It is important that you maintain your abdomen contracted and your navel pulled in for the entire duration of the exercise. Repeat this movement 03 times for each side.

**7**

3x10-15 repetitions

Lying on your back, bend your knees and put your hands behind your head, raising your trunk until your shoulders are off the ground. Try to keep your back on the ground while doing this exercise. Carry out 03 series of 10 or 15 repetitions.

**8**


3x10-15 repetitions

Lying on your side with your knees bent, move your knees away from each other by rotating or opening and closing or legs. It is important that you keep your feet together during the exercise. Repeat the exercise on both sides, with 03 series of 5 repetitions.




## Annex II: Graphs

Graph 1. This graph shows the change in mean pain severity over time in the group that performed only exercise therapy (ET) and in the group that combined it with spinal manipulative therapy (SMT+ET). Adapted from Evans R et al.



Graph 2. This graph shows the change in mean pain severity over time for the group with a home exercise program (HEP, in red), the group which combined a home exercise program with supervised exercise program (SEP+HEP, in grey) and another group which combined a home exercise program with spinal manipulative therapy (SMT+HEP, in blue). Adapted from Schulz C et al. (53)



## Annex III: Tables

Table 1. The FABQ Work consists of a Fear-Avoidance Beliefs Questionnaire. Number in square brackets represent the 95% confidence interval. Adapted from Perron M et al. (28)

Variables	Sig.	Sensitivity	Specificity	LR+	LR-	Odd Ratio
No pain in lying down	0.017	0.75 [0.58-0.87]	0.49 [0.34-0.64]	1.47 [1.05-2.06]	0.51 [0.29-0.92]	3.65 [1.3-10.6]
No use of antidepressants	0.061	0.95 [0.82-0.99]	0.24 [0.13-0.40]	1.26 [1.05-1.51]	0.20 [0.05-0.90]	5.2 [0.9-29.4]
FABQ Work <22.5	0.061	0.73 [0.56-0.85]	0.67 [0.51-0.80]	2.18 [1.38-3.43]	0.41 [0.24-0.70]	2.9 [0.9-8.6]
Number of previous treatments <5	0.144	0.68 [0.51-0.81]	0.58 [0.42-0.72]	1.60 [1.07-2.39]	0.56 [0.35-0.91]	2.2 [0.8-6.3]
Work restriction <6 months	0.161	0.85 [0.69-0.94]	0.44 [0.30-0.60]	1.53 [1.14-2.05]	0.34 [0.15-0.75]	2.48 [0.7-8.8]