



Preservers of Triple Transition Pseudo-Probabilities in Connection with Orthogonality Preservers and Surjective Isometries

Antonio M. Peralta 

Abstract. We prove that every bijection preserving triple transition pseudo-probabilities between the sets of minimal tripotents of two atomic JBW*-triples automatically preserves orthogonality in both directions. Consequently, each bijection preserving triple transition pseudo-probabilities between the sets of minimal tripotents of two atomic JBW*-triples is precisely the restriction of a (complex-)linear triple isomorphism between the corresponding JBW*-triples. This result can be regarded as triple version of the celebrated Wigner theorem for Wigner symmetries on the posets of minimal projections in $B(H)$. We also present a Tingley type theorem by proving that every surjective isometry between the sets of minimal tripotents in two atomic JBW*-triples admits an extension to a real linear surjective isometry between these two JBW*-triples. We also show that the class of surjective isometries between the sets of minimal tripotents in two atomic JBW*-triples is, in general, strictly wider than the set of bijections preserving triple transition pseudo-probabilities.

Mathematics Subject Classification. Primary 81R15, 47B49; Secondary 17C65, 46L60, 47N50.

Keywords. Wigner theorem, minimal partial isometries, minimal tripotents, triple transition pseudo-probability, preservers, Cartan factors, surjective isometry, Tingley's type theorem.

1. Introduction

As it is masterfully narrated by G. Chevalier in [10], the Geneva-Brussels School proposed the orthomodular lattice $\mathcal{P}(H)$, of all projections on a complex Hilbert space H , with the partial order defined by $p \leq q$ in $\mathcal{P}(H)$ if $pq = p$ and orthogonality determined by zero product –equivalently, the orthomodular lattice \mathbf{L} of all closed subspaces of H with the partial ordering given by inclusion and orthogonality in the Euclidean sense– as mathematical model in quantum mechanics. By a result due to C. Piron, from 1976, every isomorphism of the propositional system of all closed subspaces of a complex Hilbert space of dimension at least 3 is induced by a unitary or by an antiunitary operator (see [10, Corollary 13]). This is actually an equivalent reformulation of the celebrated Wigner’s unitary-antiunitary theorem (cf. [9]).

The elements in $\mathcal{P}(H)$ are precisely the positive partial isometries in $B(H)$. We recall that an operator e in $B(H)$ is called a *partial isometry* or a *tripotent* if $ee^*e = e$ (equivalently, ee^* or e^*e lies in $\mathcal{P}(H)$). Along this note we shall write $PI(H) = \mathcal{U}(B(H))$ for the collection of all partial isometries in $B(H)$. In [37] L. Molnár paved a new ground to establish an analogue to Piron’s version of Wigner’s unitary-antiunitary theorem for preservers of order and orthogonality between the corresponding structures of partial isometries. We should recall first that for $e, v \in \mathcal{U}(B(H))$ we write $e \leq v$ (respectively, e is orthogonal to v , $e \perp v$ in short) if $ee^* \leq vv^*$ and $e^*e \leq v^*v$ (respectively, $ev^* = v^*e = 0$). Molnár’s version of the Piron-Wigner theorem asserts that for each complex Hilbert space H with $\dim(H) \geq 3$, every bijective transformation $\Phi : \mathcal{U}(B(H)) \rightarrow \mathcal{U}(B(H))$ which preserves the partial ordering and orthogonality between partial isometries in both directions and is continuous (in the operator norm) at a single element of $\mathcal{U}(B(H))$ different from 0, extends to a real-linear triple isomorphism (cf. [37, Theorem 1]).

The Banach space $B(H)$, of all bounded linear operators on a complex Hilbert space H , is more than a prototype of C^* - and von Neumann algebra. By the Gelfand-Naimark theorem every C^* -algebra embeds as a self-adjoint subalgebra of some $B(H)$. It is known that $B(H)$ can be also regarded as particular case of type 1 Cartan factors. There are six different types of Cartan factors (see Sect. 2 for definitions), which are employed in a Gelfand-Naimark type theorem to represent every JB^* -triple isometrically as a JB^* -subtriple of an ℓ_∞ -sum of Cartan factors (cf. [22]).

As we shall see below, those complex Banach spaces whose open unit ball is a bounded symmetric domain were characterized by W. Kaup in [32] as the complex Banach spaces E admitting a continuous triple product $\{., ., .\} : E \times E \times E \rightarrow E$ (bilinear and symmetric in the outer variables and conjugate linear in the middle one) satisfying a collection of algebraic and analytic axioms (see Sect. 2). For the moment we shall simply note that every C^* -algebra is a JB^* -triple for the triple product

$$\{x, y, z\} = \frac{1}{2}(xy^*z + zy^*x), \tag{1}$$

while the class of JB^* -triples is strictly wider than the collection of all C^* -algebras, since it also contains, among other more exotic examples, all infinite dimensional complex Hilbert spaces. The fixed-points of the just commented triple product on a C^* -algebra A are the partial isometries in A . The elements e in a JB^* -triple E satisfying $\{e, e, e\} = e$ are called tripotents. The set of all tripotents in E will be denoted by $\mathcal{U}(E)$. As it will be detailed in Sect. 2, there is a natural partial order and a notion of orthogonality for elements in $\mathcal{U}(E)$, which restricted to $\mathcal{U}(B(H))$ are precisely those employed by Molnár in the theorem commented above.

A triple homomorphism between JB^* -triples E and F is a linear map $T : E \rightarrow F$ preserving triple products. Every triple homomorphism between JB^* -triples is automatically continuous [1, Lemma 1]. If T is a triple isomorphism (i.e. a bijective triple homomorphism), its restriction to $\mathcal{U}(E)$ is a surjective isometry $T|_{\mathcal{U}(E)} : \mathcal{U}(E) \rightarrow \mathcal{U}(F)$ which preserves orthogonality and partial order in both directions (we recall that very injective triple homomorphism is an isometry [1, Lemma 1]). As we shall justify later, the restriction of T to the corresponding subsets of all minimal tripotents is also a surjective isometry.

As in the case of C^* -algebras, there exist JB^* -triples E for which $\mathcal{U}(E) = \{0\}$. However, in a JB^* -triple E the extreme points of its closed unit ball are precisely the complete tripotents in E (cf. [5, Lemma 4.1], [34, Proposition 3.5] or [14, Corollary 4.8]). Thus, every JB^* -triple which is also a dual Banach space contains an abundant set of tripotents. JB^* -triples which are additionally dual Banach spaces are called *JBW*-triples*. Each JBW^* -triple admits a unique (isometric) predual and its triple product is separately weak* continuous (cf. [2]). Since each Cartan factor is a dual Banach space, each ℓ_∞ -sum of Cartan factors is a JBW^* -triple. The JBW^* -triples which are of this form are called atomic JBW^* -triples.

In a recent collaboration with Y. Friedman, we studied bijective transformations preserving orthogonality and order between the sets of tripotents of two atomic JBW^* -triples. More concretely, let $M = \bigoplus_{i \in I}^{\ell_\infty} C_i$ and $N = \bigoplus_{j \in J}^{\ell_\infty} \tilde{C}_j$ be atomic JBW^* -triples, where C_i and \tilde{C}_j are Cartan factors with rank ≥ 2 . Suppose that $\Phi : \mathcal{U}(M) \rightarrow \mathcal{U}(N)$ is a bijective transformation which preserves the partial ordering in both directions and orthogonality between tripotents. If we additionally assume that Φ is continuous at a tripotent $u = (u_i)_i$ in M with $u_i \neq 0$ for all i (or we simply assume that $\Phi|_{\mathbb{T}u}$ is continuous at a tripotent $(u_i)_i$ in M with $u_i \neq 0$ for all i), then there exists a real linear triple isomorphism $T : M \rightarrow N$ such that $T(w) = \Phi(w)$ for all $w \in \mathcal{U}(M)$ (cf. [20, Theorem 6.1]).

Back to the original statement of Wigner’s theorem, we recall the notion of transition probability between minimal (i.e. rank-one) projections in $B(H)$.

Suppose $p = \xi \otimes \xi$ and $q = \eta \otimes \eta$ are two minimal projections in $B(H)$ with ξ and η in the unit sphere of H (where $\xi \otimes \eta(\zeta) := \langle \zeta, \eta \rangle \xi$). The *transition probability* from p to q is defined as

$$TP(p, q) = \text{tr}(pq) = \text{tr}(pq^*) = \text{tr}(qp^*) = |\langle \xi, \eta \rangle|^2.$$

Let $\mathcal{P}_1(H)$ stand for the set of all minimal projections in $B(H)$. A bijective map $\Phi : \mathcal{P}_1(H) \rightarrow \mathcal{P}_1(H)$ is called a *symmetry transformation* or a *Wigner symmetry* if it preserves the transition probability between minimal projections, that is,

$$TP(\Phi(p), \Phi(q)) = \text{tr}(\Phi(p)\Phi(q)) = \text{tr}(pq) = TP(p, q), \text{ for all } (p, q \in \mathcal{P}_1(H)).$$

Wigner’s theorem proves that symmetry transformations on $\mathcal{P}_1(H)$ are characterized as those bijective maps $\Phi : \mathcal{P}_1(H) \rightarrow \mathcal{P}_1(H)$ for which there is an either unitary (i.e. a linear mapping $u : H \rightarrow H$ such that $uu^* = u^*u = 1$) or an antiunitary (i.e. a conjugate-linear mapping $u : H \rightarrow H$ such that $uu^* = u^*u = 1$) operator u on H , unique up to multiplication by a unitary scalar, such that $\Phi(p) = upu^*$ for all $p \in \mathcal{P}_1(H)$ (cf. [41], [36, page 12]).

The following version of Wigner’s theorem for minimal partial isometries is also due to L. Molnár. In the statement we see that the set of minimal projections in $B(H)$ has been enlarged to the set, $\mathcal{U}_{min}(B(H))$, of all minimal partial isometries in $B(H)$. We recall that a partial isometry e in $B(H)$ is called minimal if its left projection ee^* (equivalently, its right projection e^*e) is minimal.

Theorem 1.1 [37, Theorem 2]. *Let $\Phi : \mathcal{U}_{min}(B(H)) \rightarrow \mathcal{U}_{min}(B(H))$ be a bijective mapping satisfying*

$$\text{tr}(\Phi(e)^* \Phi(v)) = \text{tr}(e^*v), \text{ for all } e, v \in \mathcal{U}_{min}(B(H)). \tag{2}$$

Then Φ extends to a surjective complex-linear isometry. Moreover, one of the following statements holds:

- (a) *There exist unitaries u, w on H such that $\Phi(e) = uew$ ($e \in \mathcal{U}_{min}(B(H))$);*
- (b) *There exist antiunitaries u, w on H such that $\Phi(e) = ue^*w$ ($e \in \mathcal{U}_{min}(B(H))$).*

The transition probability between two minimal projections p, q in $B(H)$ coincides with $\text{tr}(pq^*) \in [0, 1]$, so the hypothesis assumed by Molnár in (2) (i.e. preservation of $\text{tr}(e^*v) \in \mathbb{C}$) is an analogue of transition probability preservation for non-necessarily positive minimal partial isometries. Let us analyse this new “generalized transition probability”. If we fix a minimal partial isometry e in $B(H)$, the functional $\varphi_e(x) = \text{tr}(e^*x)$ is the unique extreme point of the closed unit ball of $B(H)_*$, the predual of $B(H)$, at which e attains its norm, so $\text{tr}(e^*v) = \varphi_e(v)$. This is the crucial point to consider the notion of triple transition pseudo-probability from a minimal tripotent to another minimal tripotent in an arbitrary JBW*-triple as introduced in the recent reference [39]. More concretely, for each minimal tripotent e in a JBW*-triple, M , there

exists a unique pure atom (i.e. an extreme point of the closed unit ball of M_*) φ_e at which e attains its norm and the corresponding Peirce-2 projection writes in the form $P_2(e)(x) = \varphi_e(x)e$ for all $x \in M$ (cf. [21, Proposition 4]). The mapping

$$\mathcal{U}_{min}(M) \rightarrow \partial_e(\mathcal{B}_{M_*}), \quad e \mapsto \varphi_e$$

is a bijection from the set of minimal tripotents in M onto the set of pure atoms of M . Given two minimal tripotents e and v in a JBW^* -triple M , we define the *triple transition pseudo-probability* from e to v as the complex number given by

$$TTP(e, v) = \varphi_v(e). \tag{3}$$

We can no longer use the term “probability” because $TTP(e, v)$ is an element in the closed unit ball of the complex plane. In the case of $B(H)$, the triple transition pseudo-probability between two minimal projections is precisely the usual transition probability in Wigner’s theorem, while the hypothesis (2) in Theorem 1.1 simply says that Φ preserves triple transition pseudo-probabilities.

We recall for later purposes that the triple transition pseudo-probability is symmetric in the sense that $TTP(e, v) = \overline{TTP(v, e)}$, for every couple of minimal tripotents $e, v \in M$ (see [39, (2.3)]).

In view of Theorem 1.1, it is an attractive challenge to ask whether a bijection Φ between the sets of minimal tripotents of two Cartan factors (or more generally of two atomic JBW^* -triples) M and N preserving triple transition pseudo-probabilities is precisely the restriction of a (complex-)linear triple isomorphism between the corresponding JBW^* -triples. This problem has been positively solved when M and N are both Cartan factors of type 1 (i.e. Banach spaces $B(H, K)$ of bounded linear operators between complex Hilbert spaces) or when M and N are both type 4 or spin Cartan factors (see [39, Theorems 4.4 and 3.2]). It is worth to note that the proof of these results is built upon classic theorems on preservers and concrete tools for operator spaces and Hilbert spaces. The general problem remains open.

This paper presents a complete solution to the problem just presented (see Corollary 3.3). Here, instead of combining classical tools on preservers for concrete Cartan factors, we shall turn our point of view to a completely newfangled strategy with arguments and tools taken from abstract theory of JB^* -triples. As we shall see in Sect. 2, the achievements in [39, Theorem 2.3] prove that each bijective transformation Φ preserving triple transition pseudo-probabilities between the sets of minimal tripotents of two atomic JBW^* -triples M and N , admits an extension to a bijective (complex) linear mapping T_0 from the socle of M onto the socle of N whose restriction to $\mathcal{U}_{min}(M)$ is Φ , where the socle of a JB^* -triple is the subspace linearly generated by its minimal tripotents. If we additionally assume that Φ preserves orthogonality, then Φ admits an extension to a surjective (complex-)linear (isometric) triple isomorphism from M onto N (cf. [39, Corollary 2.5]). In Theorem 3.2 we prove that every bijection

preserving triple transition pseudo-probabilities between the sets of minimal tripotents of two atomic JBW*-triples automatically preserves orthogonality in both directions.

The main conclusion in this paper shows that the set of minimal tripotents in an atomic JBW*-triple together with the triple transition pseudo-probabilities among its elements is a complete invariant valid to determine the whole structure of the JB*-triple (cf. Corollary 3.3). The result should be complemented with the main conclusion of [20], which asserts that in an atomic JBW*-triple M containing no rank-one Cartan factors the poset of all tripotents in M with the partial order and the relation of orthogonality is a complete invariant for its structure of real JBW*-triple. Both results together validate the full analogy in the setting of JB*-triples with the different statements of Wigner's theorem for projections.

Another result derived from our main conclusion (see Corollary 3.4) proves that every bijection preserving triple transition pseudo-probabilities between the sets of minimal tripotents in two atomic JBW*-triples is an isometry with respect to the gap metric (i.e. the metric given by the JB*-triple norm).

It is known that the gap metric and the usual transition probability between minimal projections in $B(H)$ are mutually determined (see [36, (2.6.13) in page 127] or [23]). However, as we shall see in Remark 3.5, the triple transition pseudo-probability and the gap metric are not, in general, related each other. It naturally arises the problem of studying those bijections preserving distances between the sets of minimal tripotents in two atomic JBW*-triples. This task is culminated in Theorem 3.8. In the just quoted result we establish a variant of Tingley's theorem by proving that every surjective isometry between the sets of minimal tripotents in two atomic JBW*-triples admits an extension to a real linear surjective isometry between these two JBW*-triples. The proof is obtained by an application of the result describing the bijections preserving triple transition pseudo-probabilities between sets of minimal tripotents in atomic JBW*-triples. However, the class of surjective isometries between the sets of minimal tripotents in two atomic JBW*-triples is, in general, strictly wider than the set of bijections preserving triple transition pseudo-probabilities, since we can also find examples of extensions which are conjugate linear or which are neither complex linear nor conjugate linear.

2. Background and State-of-the-Art

This section is aimed to provide the reader the basic terminology and notions to understand the results and to fill the gaps left in the introduction. We shall also approach to a brief state-of-the-art of the main problem tackled in this paper.

Our arguments will employ tools developed in theory of JB*-triples. So, it seems pertinent to recall the definition of JB*-triple (cf. [32]), a mathematical

model originally arisen in holomorphic theory deeply studied in functional analysis.

A *JB*-triple* is a complex Banach space E together with a continuous triple product $\{\cdot, \cdot, \cdot\} : E \times E \times E \rightarrow E$, which is symmetric and bilinear in the first and third variables, conjugate-linear in the middle one, and satisfies the following axioms:

(a) (Jordan identity)

$$L(a, b)L(x, y) = L(x, y)L(a, b) + L(L(a, b)x, y) - L(x, L(b, a)y)$$

for a, b, x, y in E , where $L(a, b)$ is the operator on E given by $x \mapsto \{a, b, x\}$;

(b) $L(a, a)$ is a hermitian operator with non-negative spectrum for all $a \in E$;

(c) $\|\{a, a, a\}\| = \|a\|^3$ for each $a \in E$.

For $a \in E$, we write $Q(a)$ for the conjugate linear operator defined by $Q(a)(x) := \{a, x, a\}$. The examples of mathematical models included in the class of JB^* -triples is perhaps one of the biggest attractiveness of this notion. We have already commented that every C^* -algebra is a JB^* -triple. The same triple product employed for C^* -algebras given in (1) serves to equip the space $B(H, K)$, of all bounded linear operators between two complex Hilbert spaces H and K , with a structure of JB^* -triple. The JB^* -triples of the form $B(H, K)$ are known as *Cartan factors of type 1*. There are another 5 types of Cartan factors. Cartan factors of types 2 and 3 are subtriples of $B(H)$ defined in the following way. Fix a conjugation j (i.e. a conjugate-linear isometry or period 2) on a complex Hilbert space H , and define a linear involution on $B(H)$ by $x \mapsto x^t := jx^*j$ —this is an infinite dimensional version of the transposition in $M_n(\mathbb{C})$. *Cartan factors of type 2 and 3* are the JB^* -subtriples of $B(H)$ of all t -skew-symmetric and t -symmetric operators, respectively.

A *Cartan factor of type 4*, also called a *spin factor*, is a complex Hilbert space M provided with a conjugation $x \mapsto \bar{x}$, where the triple product and the norm are defined by

$$\{x, y, z\} = \langle x, y \rangle z + \langle z, y \rangle x - \langle x, \bar{z} \rangle \bar{y}, \tag{4}$$

and

$$\|x\|^2 = \langle x, x \rangle + \sqrt{\langle x, x \rangle^2 - |\langle x, \bar{x} \rangle|^2}, \tag{5}$$

respectively (cf. [19, Chapter 3]). The *Cartan factors of types 5 and 6* (also called *exceptional* Cartan factors) are spaces of matrices over the eight dimensional complex algebra of Cayley numbers; the type 6 consists of all 3×3 self-adjoint matrices and has a natural Jordan algebra structure, and the type 5 is the subtriple consisting of all 1×2 matrices (see [30, 31, 33] and the recent references [25, §6.3 and 6.4], [26, §3] for more details).

As we have already commented during the introduction, partial isometries in a C^* -algebras A are precisely the elements which are fixed points for its natural triple product (1). The fixed points of the triple product of a JB^* -triple E are called *tripotents*. We write $\mathcal{U}(E)$ for the set of all tripotents in E .

Each e in $\mathcal{U}(E)$ produces the following *Peirce decomposition* of the space E in terms of the eigenspaces of the operator $L(e, e)$:

$$E = E_0(e) \oplus E_1(e) \oplus E_2(e), \tag{6}$$

where $E_k(e) := \{x \in E : L(e, e)x = \frac{k}{2}x\}$ is a subtriple of E called the *Peirce- k subspace* ($k = 0, 1, 2$). *Peirce- k projection* is the name given to the natural projection of E onto $E_k(e)$ and it is usually denoted by $P_k(e)$. Triple products among elements in different Peirce subspaces obey certain laws known as *Peirce arithmetic*. Concretely, the inclusion $\{E_k(e), E_l(e), E_m(e)\} \subseteq E_{k-l+m}(e)$, and the identity $\{E_0(e), E_2(e), E\} = \{E_2(e), E_0(e), E\} = \{0\}$, hold for all $k, l, m \in \{0, 1, 2\}$, where $E_{k-l+m}(e) = \{0\}$ whenever $k - l + m$ is not in $\{0, 1, 2\}$.

The Peirce-2 subspace $E_2(e)$ is a unital JB*-algebra with respect to the product and involution given by $x \circ_e y = \{x, e, y\}$ and $x^{*e} = \{e, x, e\}$, respectively. The self-adjoint or hermitian part of $E_2(e)$ will be denoted by $E^1(e)$, that is,

$$E^1(e) = \{x \in E_2(e) : x^{*e} = \{e, x, e\} = x\} = \{x \in E : \{e, x, e\} = x\}.$$

Let us recall next the analogue to minimal partial isometry in the wider setting of JB*-triples. A non-zero tripotent e in a JB*-triple E is called (*algebraically*) *minimal* if $E_2(e) = \mathbb{C}e \neq \{0\}$. We shall denote by $\mathcal{U}_{min}(E)$ the set of all minimal tripotents in E . The tripotents $e \in E$ satisfying $E_0(e) = \{0\}$ are called *complete*.

As we have commented in the introduction, von Neumann algebras identify with those C*-algebras which are dual Banach spaces. JBW*-triples, defined as those JB*-triples which are dual Banach spaces, play the role of von Neumann algebras in the JB*-triple setting. A concrete subclass is determined by those JBW*-triples which coincide with the w*-closure of the linear span of their minimal tripotents – $B(H)$ is an example–. The triples in this subclass are known as *atomic* JBW*-triples. Deep structure results, established by Y. Friedman and B. Russo, prove that every atomic JBW*-triple is an ℓ_∞ -sum of Cartan factors (cf. [22, Proposition 2 and Theorem E]), and that every JB*-triple embeds isometrically as a JB*-subtriple of an atomic JBW*-triple.

It is now moment to concrete the definition of the partial order and the notion of orthogonality among tripotents in a JB*-triple. Let us take $e, v \in \mathcal{U}(E)$, where E is a generic JB*-triple. Following a concept that generalises the notion of orthogonality for partial isometries in $B(H)$, we shall say that e is *orthogonal* to u ($e \perp u$ in short) if $\{e, e, u\} = 0$ (equivalently, $L(e, u) = 0 \Leftrightarrow L(u, e) = 0 \Leftrightarrow e \in E_0(u) \Leftrightarrow u \in E_0(e)$ cf. [3, 7, 35]). It is known that any two orthogonal tripotents e and v in JB*-triple E are M -orthogonal, that is, $\|e \pm v\| = \max\{\|e\|, \|v\|\} = 1$ (cf. [21, Lemma 1.3(a)]).

The rank of a JB*-triple E is the minimal cardinal number r satisfying $\text{card}(S) \leq r$ for every *orthogonal* subset $S \subseteq E$, where by an orthogonal subset we mean a subset not containing zero and satisfying that $x \perp y$ for every $x \neq y$

in S (cf. [4, 6, 33] for basic background on the rank of a Cartan factor and a JBW*-triple, and its relation with reflexivity).

The natural partial order among partial isometries in $B(H)$ and, more generally, among tripotents in a JB*-triple E is defined by $e \leq u$ in $\mathcal{U}(E)$ if $u - e$ is a tripotent and $u - e \perp e$. This partial order is precisely the order considered by L. Molnár in [37], and it is a central notion in the theory of JB*-triples (cf., for example, the recent papers [24–29]). Thanks to the partial ordering we can consider tripotents which are minimal with respect to this ordering. It is easy to check that every algebraically minimal tripotent is (order) minimal, though the reciprocal implication does not necessarily hold for general JB*-triples, in a JBW*-triple order minimal and algebraic minimal tripotents coincide (cf. [14, Corollary 4.8] and [3, Lemma 4.7]).

A very useful tool in the representation theory of JB*-triples employed in our arguments and obtained in [17, Lemma 3.10], allows to describe the theoretical position of two arbitrary minimal tripotents in a Cartan factor of rank greater than or equal to 2. In order to understand the statement employed later, we recall several basic relations between tripotents. Let u, v be two tripotents in a JB*-triple E . We shall say that u and v are *collinear* ($u \top v$ in short) if $u \in E_1(v)$ and $v \in E_1(u)$. The tripotent u *governs* the tripotent v ($u \vdash v$ in short) whenever $v \in E_2(u)$ and $u \in E_1(v)$. An ordered quadruple (u_1, u_2, u_3, u_4) of minimal tripotents in a JB*-triple E is called a *quadrangle* if $u_1 \perp u_3$, $u_2 \perp u_4$, $u_1 \top u_2 \top u_3 \top u_4 \top u_1$ and $u_4 = 2\{u_1, u_2, u_3\}$ (the latter equality also holds if the indices are cyclically permuted, e.g. $u_2 = 2\{u_3, u_4, u_1\}$). An ordered triplet (v, u, \tilde{v}) of minimal tripotents in E , is called a *trangle* if $v \perp \tilde{v}$, $u \vdash v$, $u \vdash \tilde{v}$ and $v = Q(u)\tilde{v}$ (see [12, §1]).

Along this note, the unit sphere and the closed unit ball of a normed space X will be denoted by S_X and \mathcal{B}_X , respectively, and we shall write \mathbb{T} for $S_{\mathbb{C}}$.

3. Main Result

In our first result, we shall see that bijections preserving triple transition pseudo-probabilities between sets of minimal tripotents in two atomic JBW*-triples preserves the relation “being collinear” among them. It should be noted that Cartan factors of rank-one constitute a serious obstacle for the theorem describing the bijections preserving the partial order in both directions and orthogonality in one direction between the posets of all tripotents of two atomic JBW*-triples (cf. [20, Theorem 6.1 and Remark 3.6]), but in all the results in this manuscript we do not need to impose any restriction on the rank.

In general, the linear span of all minimal tripotents in a JB*-triple E , called the *socle* of E ($\text{soc}(E)$ in short), need not be a closed subspace. That is the case of the socle of $B(H)$, which coincides with the subspace, $\mathcal{F}(H)$, of all finite rank operators, and it is not closed when H is infinite dimensional.

However, if a JB*-triple E has finite rank (equivalently, E is a reflexive JB*-triple), we have $\text{soc}(E) = E$ (cf. [6, Proposition 4.5 and Remark 4.6] or [4]).

Proposition 3.1. *Let $\Phi : \mathcal{U}_{\min}(M) \rightarrow \mathcal{U}_{\min}(N)$ be a bijection preserving triple transition pseudo-probabilities, where M and N are two atomic JBW*-triples. Suppose e and v are two minimal collinear ($e \top v$) tripotents in $\mathcal{U}_{\min}(M)$. Then $\Phi(e)$ and $\Phi(v)$ are collinear ($\Phi(e) \top \Phi(v)$) in $\mathcal{U}_{\min}(N)$.*

Proof. By hypotheses, M and N can be written as ℓ_∞ -sums of two families of Cartan factors $\{C_i : i \in \Lambda_1\}$ and $\{D_j : j \in \Lambda_2\}$, respectively. Each minimal tripotent in M (respectively, in N) lies in a single summand. Let us observe that, by hypotheses, e and v must lie in the same Cartan factor C_{i_0} among those summands in M , otherwise they would be orthogonal.

By [39, Theorem 2.3] there exists a complex linear bijection $T_0 : \text{soc}(M) \rightarrow \text{soc}(N)$ whose restriction to $\mathcal{U}_{\min}(M)$ is Φ .

If $\Phi(e)$ and $\Phi(v)$ belong to different Cartan factors D_{j_1} and D_{j_2} with $j_1 \neq j_2$, then they are orthogonal. However, in such a case $\frac{1}{\sqrt{2}}e + \frac{1}{\sqrt{2}}v$ is a minimal tripotent (cf. [12, Lemma in page 306]), and thus $\Phi\left(\frac{1}{\sqrt{2}}e + \frac{1}{\sqrt{2}}v\right) = T_0\left(\frac{1}{\sqrt{2}}e + \frac{1}{\sqrt{2}}v\right) = \frac{1}{\sqrt{2}}\Phi(e) + \frac{1}{\sqrt{2}}\Phi(v)$ must be a minimal tripotent too, which is incompatible with $\Phi(e) \perp \Phi(v)$. We can therefore assume that $\Phi(e)$ and $\Phi(v)$ both belong to the same Cartan factor D_{j_0} and $\Phi(e) \not\perp \Phi(v)$.

We shall distinguish two cases. If D_{j_0} is a rank-one Cartan factor it must be a complex Hilbert space with inner product $\langle \cdot, \cdot \rangle$, regarded as a type 1 Cartan factor, and both $\Phi(e)$ and $\Phi(v)$ are norm-one elements. As we commented before, they are collinear as minimal tripotents if and only if they are orthogonal in the Euclidean sense of this complex Hilbert space. Since, for each $(\lambda_1, \lambda_2) \in S_{\ell_2^2}$, the element $\lambda_1 e + \lambda_2 v$ is a minimal tripotent (cf. [12, Lemma in page 306]), its image under Φ or T_0 , that is, $\Phi(\lambda_1 e + \lambda_2 v) = T_0(\lambda_1 e + \lambda_2 v) = \lambda_1 \Phi(e) + \lambda_2 \Phi(v)$, is a minimal tripotent in D_{j_0} , equivalently, a norm-one element of this Hilbert space. Therefore

$$\begin{aligned} 1 &= \|\lambda_1 \Phi(e) + \lambda_2 \Phi(v)\|^2 = \langle \lambda_1 \Phi(e) + \lambda_2 \Phi(v), \lambda_1 \Phi(e) + \lambda_2 \Phi(v) \rangle \\ &= |\lambda_1|^2 \|\Phi(e)\|^2 + |\lambda_2|^2 \|\Phi(v)\|^2 \\ &\quad + 2\Re \lambda_1 \bar{\lambda}_2 \langle \Phi(e), \Phi(v) \rangle, \\ &= 1 + 2\Re \lambda_1 \bar{\lambda}_2 \langle \Phi(e), \Phi(v) \rangle \end{aligned}$$

for all $(\lambda_1, \lambda_2) \in S_{\ell_2^2}$, witnessing that $\langle \Phi(e), \Phi(v) \rangle = 0$, which proves that $\Phi(e)$ and $\Phi(v)$ are orthogonal in the Euclidean sense in the Hilbert space D_{j_0} , equivalently, collinear in the Cartan factor D_{j_0} .

We assume next that D_{j_0} is Cartan factor with rank ≥ 2 . By the representation result in [17, Lemma 3.10], applied to $\Phi(e)$ and $\Phi(v)$ in D_{j_0} , one of the following statements holds:

- (1) There exist minimal tripotents $\tilde{v}_2, \tilde{v}_3, \tilde{v}_4$ in D_{j_0} and $\alpha, \beta, \gamma, \delta \in \mathbb{C}$ such that $(\Phi(e), \tilde{v}_2, \tilde{v}_3, \tilde{v}_4)$ is a quadrangle and $\Phi(v) = \alpha\Phi(e) + \beta\tilde{v}_2 + \gamma\tilde{v}_4 + \delta\tilde{v}_3$, $\alpha\delta = \beta\gamma$ and $|\alpha|^2 + |\beta|^2 + |\gamma|^2 + |\delta|^2 = 1$;
- (2) There exists a rank two tripotent $w \in D_{j_0}$, a minimal tripotent $\tilde{v} \in D_{j_0}$ and $\alpha, \beta, \gamma \in \mathbb{C}$ such that $(\Phi(e), w, \tilde{v})$ is a triangle, $\Phi(v) = \alpha\Phi(e) + \beta w + \delta\tilde{v}$, $\alpha\delta = \beta^2$ and $|\alpha|^2 + 2|\beta|^2 + |\delta|^2 = 1$.

Let us observe that $|\delta| < 1$, otherwise we would contradict $\Phi(e) \perp \Phi(v)$. We shall treat each case independently.

(1) Since e and v are collinear, the element $\frac{1}{\sqrt{2}}e + \frac{1}{\sqrt{2}}v$ is a minimal tripotent (cf. [12, Lemma in page 306]) and the same must occur to

$$\begin{aligned} \Phi\left(\frac{1}{\sqrt{2}}e + \frac{1}{\sqrt{2}}v\right) &= T_0\left(\frac{1}{\sqrt{2}}e + \frac{1}{\sqrt{2}}v\right) = \frac{1}{\sqrt{2}}\Phi(e) + \frac{1}{\sqrt{2}}\Phi(v) \\ &= \frac{1}{\sqrt{2}}(1 + \alpha)\Phi(e) + \frac{1}{\sqrt{2}}\beta\tilde{v}_2 + \frac{1}{\sqrt{2}}\gamma\tilde{v}_4 + \frac{1}{\sqrt{2}}\delta\tilde{v}_3, \end{aligned}$$

but the latter being a minimal tripotent implies that $(1 + \alpha)\delta - \beta\gamma = 0$. However, since $\alpha\delta = \beta\gamma$, it follows that $\delta = 0 = \beta\gamma$. Let us assume that $\beta = 0$ (the other case, i.e. $\gamma = 0$, is similar). Therefore, the element $\Phi\left(\frac{1}{\sqrt{2}}e + \frac{1}{\sqrt{2}}v\right) = \frac{1}{\sqrt{2}}(1 + \alpha)\Phi(e) + \frac{1}{\sqrt{2}}\gamma\tilde{v}_4$ must be a minimal tripotent, what occurs if and only if $\frac{|1+\alpha|^2}{2} + \frac{|\gamma|^2}{2} = 1$. We deduce that $\Re(\alpha) = 0$. Replacing e with ie , and having in mind that ie and v are collinear too and T_0 is complex linear, we get $\Im(\alpha) = 0$. Therefore, $\alpha = 0$, $\gamma \in \mathbb{T}$ and $\Phi(v) = \gamma\tilde{v}_4$ is collinear to $\Phi(e)$.

(2) As in the previous case, the element

$$\Phi\left(\frac{1}{\sqrt{2}}e + \frac{1}{\sqrt{2}}v\right) = T_0\left(\frac{1}{\sqrt{2}}e + \frac{1}{\sqrt{2}}v\right) = \frac{1}{\sqrt{2}}(1 + \alpha)\Phi(e) + \frac{1}{\sqrt{2}}\beta w + \frac{1}{\sqrt{2}}\delta\tilde{v},$$

must be a minimal tripotent, and thus $(1 + \alpha)\delta - \beta^2 = 0$, and consequently $\delta = \beta = 0$ and $\alpha \in \mathbb{T}$ which is impossible because Φ preserves triple transition pseudo-probabilities. So, the second case is discarded and the proof is concluded. \square

We can now establish our main result showing that every bijection preserving triple transition pseudo-probabilities between the posets of minimal tripotents of two atomic JBW*-triples automatically preserves orthogonality.

Theorem 3.2. *Let $\Phi : \mathcal{U}_{min}(M) \rightarrow \mathcal{U}_{min}(N)$ be a bijection preserving triple transition pseudo-probabilities, where M and N are two atomic JBW*-triples. Then Φ preserves orthogonality in both directions.*

Proof. We begin by observing that Φ^{-1} also preserves triple transition pseudo-probabilities. Let $T_0 : soc(M) \rightarrow soc(N)$ be the bijection extending Φ whose existence is given by [39, Theorem 2.3].

Let us take $e, v \in \mathcal{U}_{min}(M)$ with $e \perp v$. We shall prove that $\Phi(e) \perp \Phi(v)$.

By hypotheses, $M = \bigoplus_{i \in \Lambda_1}^{\ell_\infty} C_i$ and $N = \bigoplus_{j \in \Lambda_2}^{\ell_\infty} D_j$, where C_i and D_j are Cartan factors.

If $\Phi(e)$ and $\Phi(v)$ belong to different Cartan factors D_{j_1} and D_{j_2} with $j_1 \neq j_2$, the desired conclusion is clear. We shall therefore assume that $\Phi(e), \Phi(v) \in D_{j_0}$.

If D_{j_0} has rank-one, it must be a complex Hilbert space regarded as a type 1 Cartan factor, and both elements $\Phi(e)$ and $\Phi(v)$ are in its unit sphere. If $\dim(D_{j_0}) = 1$ (i.e. $D_{j_0} = \mathbb{C}$) $\Phi(v) = \mu\Phi(e)$ for some unitary $\mu \in \mathbb{C}$. However, $0 = TTP(e, v) = TTP(\Phi(e), \Phi(v)) = \mu$, which is impossible. We can therefore assume that $\dim(D_{j_0}) \geq 2$, and find a third tripotent $\hat{w} \in D_{j_0}$ (i.e. an element in the unit sphere of D_{j_0}) and $(\lambda_1, \lambda_2) \in S_{\ell_2^3}$ such that $\Phi(e) \perp_2 \hat{w}$ in the Euclidean sense and $\Phi(v) = \lambda_1\Phi(e) + \lambda_2\hat{w}$. By applying Φ^{-1} and T_0^{-1} we derive that $v = \lambda_1e + \lambda_2\Phi^{-1}(\hat{w})$. Since $\Phi(e) \top \hat{w}$ in D_{j_0} (and hence in N), Proposition 3.1, applied to Φ^{-1} , implies that e and $\Phi^{-1}(\hat{w})$ are collinear, which contradicts the fact that $v \perp e$, because $0 = \{e, e, v\} = \{e, e, \lambda_1e + \lambda_2\Phi^{-1}(\hat{w})\} = \lambda_1e + \frac{\lambda_2}{2}\Phi^{-1}(\hat{w})$, and thus $\lambda_1 = \lambda_2 = 0$. Therefore D_{j_0} must have rank ≥ 2 .

Since D_{j_0} is Cartan factor with rank ≥ 2 , Lemma 3.10 in [17], applied to $\Phi(e)$ and $\Phi(v)$ in D_{j_0} , assures that one of the following statements holds:

- (1) There exist minimal tripotents $\tilde{v}_2, \tilde{v}_3, \tilde{v}_4$ in D_{j_0} and $\alpha, \beta, \gamma, \delta \in \mathbb{C}$ such that $(\Phi(e), \tilde{v}_2, \tilde{v}_3, \tilde{v}_4)$ is a quadrangle and $\Phi(v) = \alpha\Phi(e) + \beta\tilde{v}_2 + \gamma\tilde{v}_4 + \delta\tilde{v}_3$, $\alpha\delta = \beta\gamma$ and $|\alpha|^2 + |\beta|^2 + |\gamma|^2 + |\delta|^2 = 1$;
- (2) There exists a rank two tripotent $w \in D_{j_0}$, a minimal tripotent $\tilde{v} \in D_{j_0}$ and $\alpha, \beta, \gamma \in \mathbb{C}$ such that $(\Phi(e), w, \tilde{v})$ is a triangle, $\Phi(v) = \alpha\Phi(e) + \beta w + \delta\tilde{v}$, $\alpha\delta = \beta^2$ and $|\alpha|^2 + 2|\beta|^2 + |\delta|^2 = 1$.

We treat both cases in parallel. Since Φ preserves triple transition pseudo-probabilities, $0 = TTP(e, v) = TTP(\Phi(e), \Phi(v)) = \alpha$. This implies in case (2) that $\beta = 0$ and $\Phi(v) = \delta\tilde{v} \perp \Phi(e)$, which gives the desired conclusion.

We finally handle case (1). Since $\alpha = 0 = \beta\gamma$ one of these two scalars is zero. We can assume that $\beta = 0$ (the other case is similar). Then $\Phi(v) = \gamma\tilde{v}_4 + \delta\tilde{v}_3$ with $|\gamma|^2 + |\delta|^2 = 1$. Since $\tilde{v}_4 \top \Phi(e)$ and $\tilde{v}_4 \top \tilde{v}_3$, Proposition 3.1 assures that $e \top \Phi^{-1}(\tilde{v}_4)$ and $\Phi^{-1}(\tilde{v}_4) \top \Phi^{-1}(\tilde{v}_3)$. Taking images under Φ^{-1} and T_0^{-1} we get

$$0 = \{e, e, v\} = \{e, e, \gamma\Phi^{-1}(\tilde{v}_4) + \delta\Phi^{-1}(\tilde{v}_3)\} = \frac{\gamma}{2}\Phi^{-1}(\tilde{v}_4) + \delta \{e, e, \Phi^{-1}(\tilde{v}_3)\}. \tag{7}$$

Let us make a couple of observations. First, by the preservation of triple transition pseudo-probabilities we have

$$0 = TTP(\Phi(e), \tilde{v}_3) = TTP(e, \Phi^{-1}(\tilde{v}_3)),$$

which implies that $P_2(\Phi^{-1}(\tilde{v}_3))(e) = 0$, and consequently $e = P_1(\Phi^{-1}(\tilde{v}_3))(e) + P_0(\Phi^{-1}(\tilde{v}_3))(e)$. Therefore, by Peirce arithmetic,

$$\begin{aligned} \{e, e, \Phi^{-1}(\tilde{v}_3)\} &= \{P_0(\Phi^{-1}(\tilde{v}_3))(e), P_1(\Phi^{-1}(\tilde{v}_3))(e), \Phi^{-1}(\tilde{v}_3)\} \\ &\quad + \{P_1(\Phi^{-1}(\tilde{v}_3))(e), P_1(\Phi^{-1}(\tilde{v}_3))(e), \Phi^{-1}(\tilde{v}_3)\}, \end{aligned}$$

where $\{P_0(\Phi^{-1}(\tilde{v}_3))(e), P_1(\Phi^{-1}(\tilde{v}_3))(e), \Phi^{-1}(\tilde{v}_3)\} \in M_1(\Phi^{-1}(\tilde{v}_3))$, while the second summand $\{P_1(\Phi^{-1}(\tilde{v}_3))(e), P_1(\Phi^{-1}(\tilde{v}_3))(e), \Phi^{-1}(\tilde{v}_3)\} \in M_2(\Phi^{-1}(\tilde{v}_3))$. Having in mind that $\Phi^{-1}(\tilde{v}_4) \top \Phi^{-1}(\tilde{v}_3)$, and hence $\Phi^{-1}(\tilde{v}_4) \in M_1(\Phi^{-1}(\tilde{v}_3))$, we deduce from (7) and the Peirce decomposition with respect to $\Phi^{-1}(\tilde{v}_3)$ that

$$\delta \{P_1(\Phi^{-1}(\tilde{v}_3))(e), P_1(\Phi^{-1}(\tilde{v}_3))(e), \Phi^{-1}(\tilde{v}_3)\} = 0. \tag{8}$$

If $\delta = 0$, it follows from (7) that $\gamma = 0$, and hence $\Phi(v) = 0$, which is impossible.

The other alternative from (8) is $\{P_1(\Phi^{-1}(\tilde{v}_3))(e), P_1(\Phi^{-1}(\tilde{v}_3))(e), \Phi^{-1}(\tilde{v}_3)\} = 0$. Now, an application of [21, Lemma 1.5] or [38, Theorem 2.3] gives $P_1(\Phi^{-1}(\tilde{v}_3))(e) = 0$, and thus $e = P_0(\Phi^{-1}(\tilde{v}_3))(e) \perp \Phi^{-1}(\tilde{v}_3)$. Finally, by the above arguments, we have

$$v = \Phi^{-1}(\gamma\tilde{v}_4 + \delta\tilde{v}_3) = T_0^{-1}(\gamma\tilde{v}_4 + \delta\tilde{v}_3) = \gamma\Phi^{-1}(\tilde{v}_4) + \delta\Phi^{-1}(\tilde{v}_3),$$

with $v \perp e$, $e \perp \Phi^{-1}(\tilde{v}_3)$ and $\Phi^{-1}(\tilde{v}_4) \top e$ (compare Proposition 3.1), it necessarily holds that $\gamma = 0$, and thus $\Phi(v) = \delta\tilde{v}_3 \perp \Phi(e)$. \square

By combining Theorem 3.2 with [39, Corollary 2.5] we appreciate the real impact of our conclusions in the next corollary.

Corollary 3.3. *Let $\Phi : \mathcal{U}_{min}(M) \rightarrow \mathcal{U}_{min}(N)$ be a bijective transformation preserving triple transition pseudo-probabilities (i.e. $TTP(\Phi(v), \Phi(e)) = \varphi_{\Phi(e)}(\Phi(v)) = \varphi_e(v) = TTP(v, e)$, for all e, v in $\mathcal{U}_{min}(M)$), where M and N are atomic JBW*-triples. Then Φ extends (uniquely) to a surjective complex-linear (isometric) triple isomorphism from M onto N .*

The next corollary is perhaps an interesting surprise by itself.

Corollary 3.4. *Let $\Phi : \mathcal{U}_{min}(M) \rightarrow \mathcal{U}_{min}(N)$ be a bijective transformation preserving triple transition pseudo-probabilities, where M and N are atomic JBW*-triples. Then Φ is an isometry with respect to the distances given by the triple norms.*

All previous results also hold for atomic von Neumann algebras (i.e. ℓ_∞ -sums of $B(H)$ spaces).

Let M and N be atomic JBW*-triples. Under the light of Corollary 3.4 above, it is natural to ask whether a bijection $\Phi : \mathcal{U}_{min}(M) \rightarrow \mathcal{U}_{min}(N)$ preserving distances with respect to the triple norms also preserves triple transition pseudo-probabilities. The answer is, in general, negative. The counterexamples presented below points out the different information encoded by the set of minimal tripotents equipped with the triple transition pseudo-probability

and the set of minimal projections with the usual transition probability. For example, the natural conjugation on $B(H)$, $a \mapsto a^*$, defines a conjugate-linear (isometric) triple automorphism whose restriction to $\mathcal{U}_{min}(B(H))$, defines a bijection $\Psi : \mathcal{U}_{min}(B(H)) \rightarrow \mathcal{U}_{min}(B(H))$ which preserves distances, but does not preserve triple transition pseudo-probabilities since $TTP(\Psi(\lambda e), \Psi(e)) = TTP(\overline{\lambda e^*}, e^*) = \overline{\lambda}$ is not, in general, equal to $TTP(\lambda e, e) = \lambda$ for all $e \in \mathcal{U}_{min}(B(H))$ and $\lambda \in \mathbb{T}$.

Remark 3.5. The usual operator or C^* - norm on $B(H)$ induces a metric on the set $\mathcal{P}_1(H)$, of all minimal projections in $B(H)$, which is known as the *gap metric*. The gap metric and the transition probability between elements in $\mathcal{P}_1(H)$ are mutually determined by the formula

$$\|p - q\| = \sqrt{1 - \text{tr}(pq)} = \sqrt{1 - TTP(p, q)}, \tag{9}$$

(cf. [36, (2.6.13) in page 127]). The distance, or gap metric, between two minimal tripotents e and v in a JB^* -triple E was determined in [17, Proposition 3.3] and can be computed with the following formula:

$$\|e - v\|^2 = (1 - \Re TTP(v, e)) + \sqrt{(1 - \Re TTP(v, e))^2 - \|P_0(e)(v)\|^2}. \tag{10}$$

It does not take too much time to check that (10) coincides with the formula in (9) when e and v are minimal projections (i.e. positive minimal partial isometries) in $B(H)$. To illustrate the statement that the gap metric and the triple transition pseudo-probability are not mutually determined with a concrete example, consider the tripotents $e = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $v = \begin{pmatrix} 1/3 & 1/3 \\ \sqrt{7/18} & \sqrt{7/18} \end{pmatrix}$

and $\tilde{v} = \begin{pmatrix} 1/3 & 1/4 \\ \sqrt{119/15} & \sqrt{119/20} \end{pmatrix}$ in $M_2(\mathbb{C})$.

It is easy to check that $TTP(v, e) = TTP(\tilde{v}, e) = 1/3$ while $\|e - v\| \neq \|e - \tilde{v}\|$. On the other hand, taking $\gamma, \beta \in \mathbb{R}$ such that $\gamma\beta = 1/2 \left(\sqrt{3 - \sqrt{2}} \right) / (3\sqrt{2})$, $1/4 + \beta^2 + \gamma^2 + (3 - \sqrt{2})/18 = 1$, the element $u = \begin{pmatrix} 1/2 & \beta \\ \gamma & \left(\sqrt{3 - \sqrt{2}} \right) / (3\sqrt{2}) \end{pmatrix}$ is a minimal tripotent satisfying $\|e - v\|^2 = \frac{1+2\sqrt{2}}{3\sqrt{2}} = \|e - u\|^2$ while $TTP(v, e) = 1/3 \neq 1/2 = TTP(u, e)$.

The previous discussion naturally leads to the study of surjective isometries between sets of minimal tripotents in two atomic JBW^* -triples. We are therefore connected with the celebrated Tingley’s problem in the case of atomic JBW^* -triples [16, 18]. The main result in [16] shows that every surjective isometry Δ between the unit spheres of two atomic JBW^* -triples M and N admits an extension to a real linear triple isomorphism between the JB^* -triples. Clearly, the set of minimal tripotents in a JB^* -triple E is contained in the unit sphere of E . In the case of a complex Hilbert space, regarded as a type 1 Cartan factor, the set of all minimal tripotents is precisely the whole sphere. One of

the key facts in the just commented result from [16] consists in proving that such an isometry Δ maps $\mathcal{U}_{min}(M)$ to $\mathcal{U}_{min}(N)$. In our next result this will be part of the hypothesis but at the cost of reducing the domain of our bijection.

We recall some terminology first. Following [15], the set of all *contractive perturbations* of a subset S of the closed unit ball of a Banach space X is defined as the norm-closed convex subset of \mathcal{B}_X given by

$$cp(S) = \{x \in X : \|x \pm s\| \leq 1\}.$$

For each natural $n \geq 2$, the n -th contractive perturbations of S are inductively defined by the equality $cp^{(n)}(S) = cp(cp^{(n-1)}(S))$. It is known that $S \subseteq cp^{(2)}(S)$, which gives $cp(S) = cp^{(3)}(S)$.

One of the basic tools in our previous arguments is provided by [17, Lemma 3.10], a result which describes the relative position of two minimal tripotents in a Cartan factor of rank ≥ 2 . We shall next state an analogous result for rank-one Cartan factors, which has been employed before. The statement is probably part of the folklore in JB*-triple theory and the proof is clear.

Lemma 3.6. *Let e, v be two minimal tripotents in a rank-one Cartan factor C with dimension ≥ 2 . Then there exists another minimal tripotent v_1 in C and $\alpha, \beta \in \mathbb{C}$ satisfying*

$$e \top v_1, |\alpha|^2 + |\beta|^2 = 1, \text{ and } v = \alpha e + \beta v_1.$$

Remark 3.7. Each surjective real linear isometry between two Cartan factors of rank ≥ 2 must be either complex linear or conjugate linear and a triple isomorphism (cf. [11]). A similar conclusion is, in general, false for rank-one Cartan factors. Namely, the mapping $T_0 : \ell_2^2 \rightarrow \ell_2^2, T_0(\lambda_1, \lambda_2) = (\lambda_1, \overline{\lambda_2})$ is a surjective real linear isometry which is not complex linear nor conjugate linear and does not preserve triple products. Let us see that this covers all possible possibilities. Suppose $T : H \rightarrow H$ is a surjective real linear isometry between two rank-one Cartan factors (i.e. two complex Hilbert spaces which are clearly identified). Let $\{e_j : \Lambda\}$ be an orthonormal basis of the Hilbert space H . It is easy to check that each e_j defines a minimal and maximal tripotent in H and they are all mutually collinear. It is also clear that T preserves cubes and collinearity (i.e. Euclidean orthogonality). Therefore $\{T(e_j)\}$ is an orthonormal basis of H too, and the equation $\|T(e_j) - T(ie_j)\|^2 = 2$ then implies that $T(ie_j) \in \{\pm iT(e_j)\}$ for all j . Setting $\Lambda_1 := \{j : T(ie_j) = iT(e_j)\}$, $\Lambda_2 := \{j : T(ie_j) = -iT(e_j)\}$, and $H_k = \overline{\text{span}}\{e_j : j \in \Lambda_k\}$, we have $H = H_1 \oplus_2^{\perp} H_2$, $T|_{H_1} : H_1 \rightarrow T(H_1)$ is a complex linear surjective isometry and $T|_{H_2} : H_2 \rightarrow T(H_2)$ is a conjugate linear surjective isometry. Furthermore, the natural conjugation j on H defined by $j(k_1, k_2) := (k_1, \overline{k_2})$ ($(k_1, k_2) \in T(H_1) \oplus_2^{\perp} T(H_2)$) satisfies that $j \circ T : H \rightarrow H$ is a complex linear isometry and a triple isomorphism.

Our next result determines the form of all surjective isometries between the sets of minimal tripotents in two atomic JBW*-triples.

Theorem 3.8. *Let $\Delta : \mathcal{U}_{min}(M) \rightarrow \mathcal{U}_{min}(N)$ be a surjective isometry, where M and N are atomic JBW*-triples. Then there exists a (unique) real linear isometry $T : M \rightarrow N$ such that $\Delta = T|_{\mathcal{U}_{min}(M)}$.*

Proof. The proof will be obtained by adequate adaptations of tools developed in the study of Tingley’s problem for JB*-triples (cf. [16–18, 40]) and the new conclusion in Corollary 3.3.

Let us begin by proving that Δ maps antipodal points to antipodal points, that is,

$$\Delta(-e) = -\Delta(e) \text{ for all } e \in \mathcal{U}_{min}(M). \tag{11}$$

Namely, since by hypothesis we have $\|\Delta(e) - \Delta(-e)\| = \|e - (-e)\| = 2$, Proposition 2.2 in [17] assures that

$$\Delta(-e) = -\Delta(e) + P_0(\Delta(e))(\Delta(-e)).$$

However, the fact that $\Delta(-e)$ is a minimal tripotent implies that $\Delta(-e) = -\Delta(e)$ as desired.

We shall next show that Δ preserves orthogonality among minimal tripotents in both directions, concretely,

$$e \perp v \text{ in } \mathcal{U}_{min}(M) \Leftrightarrow \Delta(e) \perp \Delta(v) \text{ in } \mathcal{U}_{min}(N). \tag{12}$$

Let us take $e, v \in \mathcal{U}_{min}(M)$ with $e \perp v$. In this case, $\|e \pm v\| = 1$, by orthogonality, and thus $\|\Delta(e) \pm \Delta(v)\| = 1$ (cf. (11)), assuring that $\Delta(v) \in cp(\Delta(e))$. As shown in [15, (6) in page 360], $cp(\{\Delta(e)\}) = \{y \in \mathcal{B}_N : y \perp \Delta(e)\} = \mathcal{B}_{N_0(\Delta(e))}$, which proves that $\Delta(v) \in \mathcal{B}_{N_0(\Delta(e))}$, and hence $\Delta(e) \perp \Delta(v)$.

In the sequel, we shall apply that M and N are atomic JBW*-triples, and hence we can write $M = \bigoplus_{i \in \Lambda_1}^{\ell_\infty} C_i$ and $N = \bigoplus_{j \in \Lambda_2}^{\ell_\infty} D_j$, where C_i and D_j are

Cartan factors. Let us comment a basic fact. Each minimal tripotent in M and in N lies in a single summand of the corresponding decomposition. Therefore, $e \not\perp v$ in $\mathcal{U}(M)_{min}$ implies that e and v belong to the same Cartan factor C_{i_0} , and by (12) $\Delta(e), \Delta(v)$ are contained in the same Cartan factor D_{j_0} .

Our next goal consists in proving that for each $i \in \Lambda_1$ there exists a unique $\sigma(i) \in \Lambda_2$ such that $\Delta(\mathcal{U}_{min}(C_i)) = \mathcal{U}_{min}(D_{\sigma(i)})$, and both summands C_i and $D_{\sigma(i)}$ have the same rank. Namely, fix any $e \in \mathcal{U}_{min}(C_i)$ and pick $\sigma(i) \in \Lambda_2$ such that $\Delta(e) \in \mathcal{U}_{min}(D_{\sigma(i)})$. Given any other $v \in \mathcal{U}_{min}(C_i)$, by [17, Lemma 3.10] and Lemma 3.6 there exists $w \in \mathcal{U}_{min}(C_i)$ such that $w \not\perp e, v$. It follows from the above comments that $\Delta(e), \Delta(v)$ and $\Delta(w)$ all lie in the same factor of the decomposition of N , therefore $\Delta(v) \in \mathcal{U}_{min}(D_{\sigma(i)})$. This proves that $\Delta(\mathcal{U}_{min}(C_i)) \subseteq \mathcal{U}_{min}(D_{\sigma(i)})$, and the equality follows from the same argument applied to Δ^{-1} . The rest is clear from the bijectivity of Δ and (12).

Pick $i \in \Lambda_1$ such that C_i and $D_{\sigma(i)}$ have rank-one. In this case $\Delta|_{\mathcal{U}_{min}(C_i)} : \mathcal{U}_{min}(C_i) \rightarrow \mathcal{U}_{min}(D_{\sigma(i)})$ is a surjective isometry, and the assumption concerning the rank implies that $\mathcal{U}_{min}(C_i) = S_{C_i}$ and $\mathcal{U}_{min}(D_{\sigma(i)}) = S_{D_{\sigma(i)}}$. We

can therefore apply Ding’s solution to Tingley’s problem for Hilbert spaces [13, Theorem 2.2] to deduce the existence of a surjective real linear isometry $T_i : C_i \rightarrow D_{\sigma(i)}$ satisfying $T_i(e) = \Delta(e)$ for all $e \in \mathcal{U}_{\min}(C_i) = S_{C_i}$. Although, T_i need not be complex linear nor conjugate linear (cf. Remark 3.7), we can find a conjugation j_i on $D_{\sigma(i)}$ such that $j_i \circ T_i : C_i \rightarrow D_{\sigma(i)}$ is an isometric (linear) triple isomorphism. This concludes the discussion for rank-one Cartan factors in the decomposition of M .

In the following we shall focus in the case in which C_i and $D_{\sigma(i)}$ are Cartan factors with rank ≥ 2 and $\Delta|_{\mathcal{U}_{\min}(C_i)} : \mathcal{U}_{\min}(C_i) \rightarrow \mathcal{U}_{\min}(D_{\sigma(i)})$ is a surjective isometry.

Let us next prove that

$$\Delta(ie) \in \{\pm i\Delta(e)\}, \text{ for all } e \in \mathcal{U}_{\min}(C_i). \tag{13}$$

Namely, pick a minimal tripotent $\tilde{v} = \Delta(v) \perp \Delta(e)$ in $\mathcal{U}_{\min}(D_{\sigma(i)})$. By (12) $v \perp e$, equivalently, $v \perp ie$, and thus $\tilde{v} = \Delta(v) \perp \Delta(ie)$ (cf. (12)). It follows that

$$\begin{aligned} \{\Delta(e)\}^\perp \cap \mathcal{U}_{\min}(D_{\sigma(i)}) &= \{\tilde{v} \in \mathcal{U}_{\min}(D_{\sigma(i)}) : \tilde{v} \perp \Delta(e)\} \\ &= \{\Delta(ie)\}^\perp \cap \mathcal{U}_{\min}(D_{\sigma(i)}) = \{\tilde{v} \in \mathcal{U}_{\min}(D_{\sigma(i)}) : \tilde{v} \perp \Delta(ie)\}. \end{aligned}$$

Since the linear combinations of minimal tripotents in the orthogonal complement of $\Delta(e)$ in $D_{\sigma(i)}$ are weak* dense in this orthogonal complement, we deduce from the above that

$$\begin{aligned} \{\Delta(e)\}^\perp \cap D_{\sigma(i)} &= \{\tilde{z} \in D_{\sigma(i)} : \tilde{z} \perp \Delta(e)\} \\ &= \{\Delta(ie)\}^\perp \cap D_{\sigma(i)} = \{\tilde{z} \in D_{\sigma(i)} : \tilde{z} \perp \Delta(ie)\}. \end{aligned}$$

Consequently,

$$\Delta(ie) \in \{\Delta(e)\}^{\perp\perp} \cap D_{\sigma(i)} = \{\tilde{z} \in D_{\sigma(i)} : \tilde{z} \perp \{\Delta(e)\}^\perp\}.$$

Since $\Delta(e)$ is a minimal tripotent in a Cartan factor with rank ≥ 2 , it cannot be complete, and hence $\{\Delta(e)\}^{\perp\perp} \cap D_{\sigma(i)} = (D_{\sigma(i)})_2(\Delta(e)) = \mathbb{C}\Delta(e)$ (cf. [8, Remark 3.4]). We can therefore find a unitary $\mu \in \mathbb{T}$ such that $\Delta(ie) = \mu\Delta(e)$. By applying that Δ is an isometry we get

$$|1 - \mu| = \|\Delta(e) - \Delta(ie)\| = \|e - ie\| = \sqrt{2},$$

which gives $\mu = \pm i$, and concludes the proof of (13).

Building upon (13) we can now prove the key step in the proof. Let $e \in \mathcal{U}_{\min}(C_i)$, then one, and precisely one, of the following statements holds:

- (14.a) $\Delta(ie) = i\Delta(e)$, $TTP(v, e) = TTP(\Delta(v), \Delta(e))$ and $\Delta(iv) = i\Delta(v)$ for all $v \in \mathcal{U}_{\min}(C_i)$.
- (14.b) $\Delta(ie) = -i\Delta(e)$, $TTP(v, e) = \overline{TTP(\Delta(v), \Delta(e))}$ and $\Delta(iv) = -i\Delta(v)$ for all $v \in \mathcal{U}_{\min}(C_i)$.

Fix arbitrary elements $e, v \in \mathcal{U}_{\min}(C_i)$. The proof relies on the relative position of the minimal tripotents e, v and their images. By [17, Lemma 3.10]

applied to e, v in C_i and their images in $D_{\sigma(i)}$ one of the statements in the following two couples holds:

- (1) There exist minimal tripotents v_2, v_3, v_4 in C_i and complex numbers $\alpha, \beta, \gamma, \delta$ such that (e, v_2, v_3, v_4) is a quadrangle, $|\alpha|^2 + |\beta|^2 + |\gamma|^2 + |\delta|^2 = 1$, $\alpha\delta = \beta\gamma$, and $v = \alpha e + \beta v_2 + \gamma v_4 + \delta v_3$;
 - (2) There exist a minimal tripotent $\tilde{v} \in C_i$, a rank two tripotent $u \in C_i$, and complex numbers α, β, δ such that (e, u, \tilde{v}) is a triangle, $|\alpha|^2 + 2|\beta|^2 + |\delta|^2 = 1$, $\alpha\delta = (\beta)^2$, and $v = \alpha e + \beta u + \delta \tilde{v}$.
- (1') There exist minimal tripotents w_2, w_3, w_4 in $D_{\sigma(i)}$, and complex numbers $\alpha', \beta', \gamma', \delta'$ such that $(\Delta(e), w_2, w_3, w_4)$ is a quadrangle, $|\alpha'|^2 + |\beta'|^2 + |\gamma'|^2 + |\delta'|^2 = 1$, $\alpha'\delta' = \beta'\gamma'$, and $\Delta(v) = \alpha'\Delta(e) + \beta'w_2 + \gamma'w_4 + \delta'w_3$;
 - (2') There exist a minimal tripotent $w \in D_{\sigma(i)}$, a rank two tripotent $\tilde{u} \in D_{\sigma(i)}$, and complex numbers α', β', δ' such that $(\Delta(e), \tilde{u}, w)$ is a triangle, $|\alpha'|^2 + 2|\beta'|^2 + |\delta'|^2 = 1$, $\alpha'\delta' = (\beta')^2$, and $\Delta(v) = \alpha'\Delta(e) + \beta'\tilde{u} + \delta'w$.

Case 1 $\Delta(ie) = i\Delta(e)$.

Assume first that (1) and (1') hold. By the formula measuring the distance between minimal tripotents (10) (cf. [17, Proposition 3.3]), the hypothesis on Δ and (11) we have

$$(1 \pm \Re(\alpha)) + \sqrt{(1 \pm \Re(\alpha))^2 - |\delta|^2} = \|e \pm v\|^2$$

$$= \|\Delta(e) \pm \Delta(v)\|^2 = (1 \pm \Re(\alpha')) + \sqrt{(1 \pm \Re(\alpha'))^2 - |\delta'|^2},$$

equivalently,

$$(\mp \Re(\alpha') \pm \Re(\alpha)) + \sqrt{(1 \pm \Re(\alpha))^2 - |\delta|^2} = \sqrt{(1 \pm \Re(\alpha'))^2 - |\delta'|^2}. \tag{14}$$

By squaring both terms in the equations and subtracting the resulting identities we get

$$2(\Re(\alpha') - \Re(\alpha)) \left(2 + \sqrt{(1 - \Re(\alpha))^2 - |\delta|^2} + \sqrt{(1 + \Re(\alpha))^2 - |\delta|^2} \right) = 0,$$

which implies that $\Re(\alpha) = \Re(\alpha')$.

Now, by applying that $\Delta(ie) = i\Delta(e)$ and repeating the above arguments we have

$$\begin{aligned} (1 \pm \Im(\alpha)) + \sqrt{(1 \pm \Im(\alpha))^2 - |\delta|^2} &= \|ie \pm v\|^2 = \|\Delta(ie) \pm \Delta(v)\|^2 \\ &= \|i\Delta(e) \pm \Delta(v)\|^2 = (1 \pm \Im(\alpha')) + \sqrt{(1 \pm \Im(\alpha'))^2 - |\delta'|^2}, \end{aligned}$$

leading to $\Im(\alpha) = \Im(\alpha')$, and hence $\alpha = \alpha'$. We have therefore proved that

$$TTP(v, e) = \alpha = \alpha' = TTP(\Delta(v), \Delta(e))$$

as desired.

The arguments in cases **(1)** and **(2')**, **(2)** and **(1')**, and **(2)** and **(2')** are exactly the same, or even particular cases, and all lead to $TTP(v, e) = \alpha = \alpha' = TTP(\Delta(v), \Delta(e))$. This concludes the proof of the first statement in (14.a).

Case 2 $\Delta(ie) = -i\Delta(e)$.

Assuming that **(1)** and **(1')** hold we have

$$(1 \pm \Re(\alpha)) + \sqrt{(1 \pm \Re(\alpha))^2 - |\delta|^2} = \|e \pm v\|^2$$

$$= \|\Delta(e) \pm \Delta(v)\|^2 = (1 \pm \Re(\alpha')) + \sqrt{(1 \pm \Re(\alpha'))^2 - |\delta'|^2},$$

and

$$\begin{aligned} (1 \pm \Im(\alpha)) + \sqrt{(1 \pm \Im(\alpha))^2 - |\delta|^2} &= \|ie \pm v\|^2 = \|\Delta(ie) \pm \Delta(v)\|^2 \\ &= \|-i\Delta(e) \pm \Delta(v)\|^2 = (1 \mp \Im(\alpha')) + \sqrt{(1 \mp \Im(\alpha'))^2 - |\delta'|^2}, \end{aligned}$$

equations which combined give $TTP(v, e) = \alpha = \overline{\alpha'} = \overline{TTP(\Delta(v), \Delta(e))}$. The other possible cases can be similarly treated, and all together prove the first statement in (14.b).

Let us now prove the final claims in (14.a) and (14.b). Suppose on the contrary that we can find $e, v \in \mathcal{U}_{min}(C_i)$ such that $\Delta(ie) = i\Delta(e)$ and $\Delta(iv) = -i\Delta(v)$.

We shall first show that $\Delta(iw) = i\Delta(w)$ (respectively, $\Delta(iw) = -i\Delta(w)$) for all $w \in \mathcal{U}_{min}(C_i)$ with $TTP(w, e) \neq 0$ (respectively, $TTP(w, v) \neq 0$). Namely, for each $w \in \mathcal{U}_{min}(C_i)$ we know that $\Delta(iw) \in \{\pm i\Delta(w)\}$ (cf. 13). If $\Delta(iw) = -i\Delta(w)$ (respectively, $\Delta(iw) = i\Delta(w)$) it follows from the first part of (14.a) (respectively, from the first part of (14.b)) that

$$\begin{aligned} -iTTP(w, e) &= TTP(-i\Delta(w), \Delta(e)) = TTP(\Delta(iw), \Delta(e)) \\ &= TTP(iw, e) = iTTP(w, e), \end{aligned}$$

(respectively,

$$\begin{aligned} i\overline{TTP(w, v)} &= \overline{TTP(i\Delta(w), \Delta(v))} = \overline{TTP(\Delta(iw), \Delta(v))} \\ &= \overline{TTP(iw, e)} = -i\overline{TTP(w, e)} \end{aligned}$$

which forces to the condition $TTP(w, e) = 0$ (respectively, $TTP(w, v) = 0$).

We deduce from the above paragraphs and the assumptions on e and v that $TTP(v, e) = 0 = \overline{TTP(e, v)}$. Combining this information with the result describing the relative position of two minimal tripotents in [17, Lemma 3.10] (as employed in many cases before), we can assume the existence of minimal tripotents v_2, v_3, v_4 in C_i and complex numbers β and δ such that (e, v_2, v_3, v_4) is a quadrangle, $|\beta|^2 + |\delta|^2 = 1$, and $v = \beta v_2 + \delta v_3$. The element $u = \frac{1}{\sqrt{2}}e + \frac{1}{\sqrt{2}}v_2$ is a minimal tripotent in C_i with $TTP(u, e) = \frac{1}{\sqrt{2}} \neq 0$ and $TTP(u, v) = \frac{\beta}{\sqrt{2}}$.

The previous conclusion shows that $\beta = 0$, and hence $v = \delta v_3$, for a unitary $\delta \in \mathbb{T}$. In such a case $\tilde{u} = \frac{1}{2}(e + v_2 + v_3 + v_4)$ is a minimal tripotent such that $TTP(\tilde{u}, e) = \frac{1}{2} \neq 0$ and $TTP(\tilde{u}, v) = \frac{\bar{\delta}}{2} \neq 0$, which is impossible. This concludes the proof of (14.a) and (14.b).

Let us define $\Lambda_{1,0} := \left\{ i \in \Lambda_1 : C_i \text{ has rank-one} \right\}$,

$$\Lambda_{1,l} := \left\{ i \in \Lambda_1 : C_i \text{ has rank } \geq 2 \text{ and } \exists e \in \mathcal{U}_{min}(C_i) \text{ with } \Delta(ie) = i\Delta(e) \right\},$$

and

$$\Lambda_{1,c} := \left\{ i \in \Lambda : C_i \text{ has rank } \geq 2 \text{ and } \exists e \in \mathcal{U}_{min}(C_i) \text{ with } \Delta(ie) = -i\Delta(e) \right\}.$$

We deduce from (14.a) and (14.b) that $\Delta(ie) = i\Delta(e)$ for every $i \in \Lambda_{1,l}$, and every $e \in \mathcal{U}_{min}(C_i)$ and $\Delta(ie) = -i\Delta(e)$ for every $i \in \Lambda_{1,c}$, and each $e \in \mathcal{U}_{min}(C_i)$.

For each $i \in \Lambda_{1,0}$ there exists a conjugation j_i on $D_{\sigma(i)}$, a real linear surjective isometry $T_i : C_i \rightarrow D_{\sigma(i)}$ such that $T_i(e) = \Delta(e)$ for all $e \in \mathcal{U}_{min}(C_i) = S_{C_i}$ and $j_i \circ T_i : C_i \rightarrow D_{\sigma(i)}$ is an isometric (complex linear) triple isomorphism (cf. Remark 3.7). For $i \in \Lambda_{1,c}$ we can find a conjugation j_i on $D_{\sigma(i)}$ (the existence is guaranteed by [33, Theorem 4.1]). For $i \in \Lambda_{1,l}$ we set $j_i = Id_{D_{\sigma(i)}}$. Define a real linear mapping $J : N = \bigoplus_{j \in \Lambda_2}^{\ell_\infty} D_j \rightarrow N = \bigoplus_{j \in \Lambda_2}^{\ell_\infty} D_j$, by $J((y_{\sigma(i)})_{i \in \Lambda_1}) := (j_i(y_{\sigma(i)}))_{i \in \Lambda_1}$. The mapping J is a real linear surjective isometry, and by construction, for $i \in \Lambda_{1,0} \cup \Lambda_{1,c}$ and $e, v \in \mathcal{U}_{min}(C_i)$ we have $J\Delta(ie) = j_i\Delta(ie) = ij_i\Delta(e) = iJ\Delta(e)$, and thus

$$TTP(J\Delta(e), J\Delta(v)) = TTP(j_i\Delta(e), j_i\Delta(v)) = TTP(e, v).$$

Clearly, $TTP(J\Delta(e), J\Delta(v)) = TTP(e, v)$, for all $i \in \Lambda_{1,l}$ and $e, v \in \mathcal{U}_{min}(C_i)$, and hence the same conclusion holds for all $e, v \in \mathcal{U}_{min}(M)$ by orthogonality. We have therefore shown that the mapping $J\Delta : \mathcal{U}_{min}(M) \rightarrow \mathcal{U}_{min}(N)$ is a surjective isometry preserving triple transition pseudo-probabilities. Corollary 3.3 asserts that $J\Delta$ extends (uniquely) to a surjective complex-linear (isometric) triple isomorphism Φ from M onto N . Finally, the mapping $J^{-1}\Phi : M \rightarrow N$ is a surjective real linear isometry whose restriction to $\mathcal{U}_{min}(M)$ is Δ . □

Funding Funding for open access publishing: Universidad de Granada/CBUA; Please verify relation to: Universidad de Granada. Author partially supported by Project PID2021-122126NB-C31, financed by: ERDF / Ministry of Science and Innovation - State Research Agency, Junta de Andalucía Grants FQM375 and PY20.00255, and by the IMAG–María de Maeztu Grant CEX2020-001105-M/AEI/10.13039/501100011033. Funding for open access charge: Universidad de Granada / CBUA.

Data availability Statement Data sharing is not applicable to this article as no datasets were generated or analyzed during the preparation of the paper.

Declarations

Conflict of interest The author declares that he has no conflict of interest.

Open Access. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- [1] Barton, T.J., Dang, T., Horn, G.: Normal representations of Banach Jordan triple systems. *Proc. Am. Math. Soc.* **102**(3), 551–555 (1988)
- [2] Barton, T., Timoney, R.M.: Weak*-continuity of Jordan triple products and its applications. *Math. Scand.* **59**, 177–191 (1986)
- [3] Battaglia, M.: Order theoretic type decomposition of JBW^* -triples. *Quart. J. Math. Oxford Ser. (2)* **42**(166), 129–147 (1991)
- [4] Becerra Guerrero, J., López Pérez, G., Peralta, A.M., Rodríguez-Palacios, A.: Relatively weakly open sets in closed balls of Banach spaces, and real JB^* -triples of finite rank. *Math. Ann.* **330**(1), 45–58 (2004)
- [5] Braun, R., Kaup, W., Upmeyer, H.: A holomorphic characterisation of Jordan C^* -algebras. *Math. Z.* **161**, 277–290 (1978)
- [6] Bunce, L.J., Chu, C.-H.: Compact operations, multipliers and Radon-Nikodym property in JB^* -triples. *Pacific J. Math.* **153**, 249–265 (1992)
- [7] Burgos, M., Fernández-Polo, F.J., Garcés, J., Martínez, J., Peralta, A.M.: Orthogonality preservers in C^* -algebras, JB^* -algebras and JB^* -triples. *J. Math. Anal. Appl.* **348**, 220–233 (2008)
- [8] Burgos, M., Garcés, J., Peralta, A.M.: Automatic continuity of biorthogonality preservers between weakly compact JB^* -triples and atomic JBW^* -triples. *Studia Math.* **204**(2), 97–121 (2011)
- [9] Casinelli, G., de Vito, E., Lahti, P., Levrero, A.: Symmetry groups in quantum mechanics and the theorem of Wigner on the symmetry transformations. *Rev. Mat. Phys.* **8**, 921–941 (1997)
- [10] Chevalier, G.: Wigner's theorem and its generalizations. In: *Handbook of Quantum Logic and Quantum Structures*, pp. 429–475, Elsevier Sci. B.V., Amsterdam (2007)

- [11] Dang, T.: Real isometries between JB^* -triples. *Proc. Am. Math. Soc.* **114**(4), 971–980 (1992)
- [12] Dang, T., Friedman, Y.: Classification of JBW^* -triple factors and applications. *Math. Scand.* **61**(2), 292–330 (1987)
- [13] Ding, G.G.: The 1-Lipschitz mapping between the unit spheres of two Hilbert spaces can be extended to a real linear isometry of the whole space. *Sci. China Ser. A* **45**(4), 479–483 (2002)
- [14] Edwards, C.M., Rüttimann, G.T.: On the facial structure of the unit balls in a JBW^* -triple and its predual. *J. Lond. Math. Soc.* **38**, 317–332 (1988)
- [15] Fernández-Polo, F.J., Martínez, J., Peralta, A.M.: Contractive perturbations in JB^* -triples. *J. Lond. Math. Soc.* **2**(85), 349–364 (2012)
- [16] Fernández-Polo, F.J., Peralta, A.M.: Tingley’s problem through the facial structure of an atomic JBW^* -triple. *J. Math. Anal. Appl.* **455**, 750–760 (2017)
- [17] Fernández-Polo, F.J., Peralta, A.M.: Low rank compact operators and Tingley’s problem. *Adv. Math.* **338**, 1–40 (2018)
- [18] Fernández-Polo, F.J., Peralta, A.M.: On the extension of isometries between the unit spheres of a C^* -algebra and $B(H)$. *Trans. Am. Math. Soc.* **5**, 63–80 (2018)
- [19] Friedman, Y.: Physical applications of homogeneous balls. With the assistance of Tzvi Scarr. *Progress in Mathematical Physics*, 40. Birkhäuser Boston, Inc., Boston, MA (2005)
- [20] Friedman, Y., Peralta, A.M.: Representation of symmetry transformations on the sets of tripotents of spin and Cartan factors. *Anal. Math. Phys.* **12**(1), 37–52 (2022)
- [21] Friedman, Y., Russo, B.: Structure of the predual of a JBW^* -triple. *J. Reine u. Angew. Math.* **356**, 67–89 (1985)
- [22] Friedman, Y., Russo, B.: The Gelfand-Naimark theorem for JB^* -triples. *Duke Math. J.* **53**, 139–148 (1986)
- [23] Gehér, G.P.: An elementary proof for the non-bijective version of Wigner’s theorem. *Phys. Lett. A* **378**(30–31), 2054–2057 (2014)
- [24] Hamhalter, J.: Dye’s theorem for tripotents in von Neumann algebras and JBW^* -triples. *Banach J. Math. Anal.* **15**(3), 49–19 (2021)
- [25] Hamhalter, J., Kalenda, O.F.K., Peralta, A.M.: Finite tripotents and finite JBW^* -triples. *J. Math. Anal. Appl.* **490**(1), 124217 (2020)
- [26] Hamhalter, J., Kalenda, O.F.K., Peralta, A.M.: Determinants in Jordan matrix algebras, to appear in *Linear and Multilinear Algebra* (2022). <https://doi.org/10.1080/03081087.2022.2049187>
- [27] Hamhalter, J., Kalenda, O.F.K., Peralta, A.M.: Order type relations on the set of tripotents in a JB^* -triple, preprint (2021). [arXiv:2112.03155](https://arxiv.org/abs/2112.03155)
- [28] Hamhalter, J., Kalenda, O.F.K., Peralta, A.M., Pfitzner, H.: Measures of weak non-compactness in preduals of von Neumann algebras and JBW^* -triples. *J. Funct. Anal.* **278**(1), 108300 (2020)
- [29] Hamhalter, J., Kalenda, O.F.K., Peralta, A.M., Pfitzner, H.: Grothendieck’s inequalities for JB^* -triples: proof of the Barton-Friedman conjecture. *Trans. Am. Math. Soc.* **374**(2), 1327–1350 (2021)

- [30] Harris, L.A.: Bounded symmetric homogeneous domains in infinite dimensional spaces. In: *Proceedings on Infinite Dimensional Holomorphy (Kentucky 1973)* pp. 13–40. *Lecture Notes in Math.* 364. Berlin-Heidelberg-New York: Springer (1974)
- [31] Hervés, F.J., Isidro, J.M.: Isometries and automorphisms of the spaces of spinors. *Rev. Mat. Univ. Complut. Madrid* **5**(2–3), 193–200 (1992)
- [32] Kaup, W.: A Riemann Mapping Theorem for bounded symmetric domains in complex Banach spaces. *Math. Z.* **183**, 503–529 (1983)
- [33] Kaup, W.: On real Cartan factors. *Manuscripta Math.* **92**, 191–222 (1997)
- [34] Kaup, W., Upmeyer, H.: Jordan algebras and symmetric Siegel domains in Banach spaces. *Math. Z.* **157**, 179–200 (1977)
- [35] Loos, O.: Bounded symmetric domains and Jordan pairs. *Lecture Notes. Univ. California at Irvine* (1977)
- [36] Molnár, L.: *Selected Preserver Problems on Algebraic Structures of Linear Operators and on Function Spaces.* *Lecture Notes in Math*, vol. 1895. Springer-Verlag, Berlin (2007)
- [37] Molnár, L.: On certain automorphisms of sets of partial isometries. *Arch. Math. (Basel)* **78**(1), 43–50 (2002)
- [38] Peralta, A.M.: Positive definite hermitian mappings associated with tripotent elements. *Expo. Math.* **33**, 252–258 (2015)
- [39] Peralta, A.M.: Maps preserving triple transition pseudo-probabilities, to appear in *RIMS Kôkyûroku Bessatsu*. [arXiv:2204.03463](https://arxiv.org/abs/2204.03463)
- [40] Peralta, A.M., Tanaka, R.: A solution to Tingley’s problem for isometries between the unit spheres of compact C^* -algebras and JB^* -triples. *Sci. China Math.* **62**(3), 553–568 (2019)
- [41] Wigner, E.P.: *Gruppentheorie und ihre Anwendung auf die Quantenmechanik der Atomspektrum*, Fredrik Vieweg und Sohn (1931)

Antonio M. Peralta
Departamento de Análisis Matemático, Facultad de Ciencias,
Instituto de Matemáticas de la Universidad de Granada (IMAG)
Universidad de Granada
18071 Granada
Spain
e-mail: aperalta@ugr.es

Received: September 10, 2022.

Accepted: December 20, 2022.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.