

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/12290842>

# Hepatitis C virus and Epstein-Barr virus: Dual infection or atypical antibody response

Article in *Microbios* · February 2000

Source: PubMed

---

CITATIONS

2

---

READS

239

2 authors:



José Gutiérrez-Fernández  
University of Granada

472 PUBLICATIONS 3,965 CITATIONS

SEE PROFILE



Manuel Rodríguez-Iglesias  
Universidad de Cádiz

175 PUBLICATIONS 3,981 CITATIONS

SEE PROFILE

Laboratory, Spain) and with confirmatory tests (Immunoblot, RIBA3, Ortho Laboratory, U.S.A.), and the presence of HCV RNA (Amplifcor HCV, Roche Diagnostics Laboratories, Switzerland) was investigated. Briefly, serum samples were tested for HCV RNA by reverse transcription-polymerase chain reaction (PCR), with primers derived from the highly conserved 5'-untranslated region. The specificity of the amplifications was confirmed by hybridization. Negative and positive controls were included in each run (four controls for each set of 20-25 clinical samples). The sensitivity of the assay was 100 RNA copies by 100 µl of serum.

### Results and discussion

In seven patients anti-HCV antibodies (C33 and C22c RIBA bands) were detected but all were PCR negative. It is not known whether these results were due to an HCV infection solely or an atypical response to HCV in the course of infectious mononucleosis. We must emphasize that in hepatitis differential diagnoses can result from serological reactions to HCV and EBV. These may be due to (1) superinfection by EBV in a patient with previous infection by HCV, and (2) a false antibody response to HCV by B lymphocyte non-specific clonal stimulation, in a manner analogous to that which has occurred with other viral infections. In the absence of any patient history of antibody response to HCV, the clinical and serological follow-up alone does not permit these two possible causes of infection to be differentiated.

### References

- FIKAR C.H., McKee C. and Park D. 1994. False positivity of IgM antibody to Epstein-Barr viral capsid antigen during acute hepatitis A infection. *Ped. Infect. Dis. J.* 13 413-4.
- GUTIÉRREZ J. and Maroto C. 1995. False positive antibodies to *Borrelia burgdorferi* in infections by herpes virus. *Biomedical Letters* 52 33-5.
- GUTIÉRREZ J., Piedrola G. and Maroto C. 1997a. Reliability of four methods for the diagnosis of acute infection by Epstein-Barr virus. *J. clin. Lab. Anal.* 11 78-81.
- GUTIÉRREZ J., Rodríguez M., Piedrola G. and Maroto C. 1997b. Behaviour of IgG antibody avidity for the antigen and of IgA in active cytomegalovirus, Epstein-Barr virus, herpes simplex virus and human herpes virus 6 infections. Adaptation of a commercial test. *J. Infect.* 35 25-30.
- GUTIÉRREZ J., Vergara M. J., Piedrola G. and Maroto C. 1999. Clinical reliability of IgG, IgA and IgM antibodies in detecting the Epstein-Barr virus in different stages of infection with a commercial nonrecombinant polyanalytic ELISA. *J. clin. Lab. Anal.* 13 65-8.
- LENNETTE E. T. 1995. Epstein-Barr virus. In *Manual of Clinical Microbiology*. pp 905-10. Edited by P. R. Murray, E. J. Baron, M. A. Tenover and R. H. Tenover. A. S. M. Press.
- WILBER J. C. 1995. Hepatitis C virus. In *Manual of Clinical Microbiology*. pp 1050-5. Edited by P. R. Murray, E. J. Baron, M. A. Tenover and R. H. Tenover. A. S. M. Press.

Accepted 12 January 2000

# Hepatitis C virus and Epstein-Barr virus: dual infection or atypical antibody response

José Gutiérrez<sup>1\*</sup> and Manuel Rodríguez-Iglesias<sup>2</sup>

<sup>1</sup>Department of Microbiology, School of Medicine, University of Granada, St Cecilia Hospital, 12 Av. Madrid, E-18012 Granada, Spain, and <sup>2</sup>Department of Microbiology, University Puerto Real Hospital, University of Cadiz, Cadiz, Spain (\*Reprint address)

Fdo.: Francisco José Fernández Yáñez  
Isla del Negorrido

**Key words:** hepatitis C virus, Epstein-Barr virus, antibodies, RNA

## Abstract

Positive serological reactions against hepatitis C virus (HCV) appeared in the course of Epstein-Barr virus (EBV) infectious mononucleosis. In 429 consecutive patients of Epstein-Barr virus (EBV) with high levels of transaminases, 28 patients were found. The presence of anti-HCV antibodies and HCV RNA was studied in these subjects. In seven patients anti-HCV antibodies (C33 and C22c RIBA bands) were detected, but all were polymerase chain reaction (PCR) negative. These results may have been due solely to a HCV infection or were an atypical response to HCV in the course of infectious mononucleosis.

## Introduction

In the course of acute infection by Epstein-Barr virus (EBV) immunoglobulin M (IgM) to EBV and other infectious agents can be produced (Fikar *et al.*, 1994; Gutiérrez and Maroto, 1995; Gutiérrez *et al.*, 1997a; Gutiérrez *et al.*, 1997b; Gutiérrez *et al.*, 1999). In patients who are immunosuppressed (transplantations, AIDS, haemodialysis) it is frequently the case that dual infection of EBV and hepatitis C virus (HCV), is diagnosed through the detection of specific antibodies (Lennette, 1995; Wilber, 1995). This short communication records the appearance of positive serological reactions against HCV in the course of EBV infectious mononucleosis.

## Materials and methods

In patients (429 *in toto*) with high levels of transaminases 28 patients with Epstein-Barr virus (EBV) primary infection were found. The infection was defined by (1) the presence of IgM against a pool of antigens comprising Epstein-Barr nuclear antigen (EBNA), membrane antigen (MA), early antigen (EA), and viral capsid antigen (VCA) (ELISA; Enzygnost EBV, DadeBehring Laboratory, Germany); (2) no detection of IgM to cytomegalovirus (ELISA; Enzygnost Cytomegalovirus, DadeBehring Laboratory, Germany) and human herpes virus 6 (ELISA; PanBio Laboratory, Australia); and (3) no immunoglobulin G (IgG) to EBNA present (ELISA; Incstar Laboratory, U.S.A.). The patients were studied in the presence of anti-HCV antibodies for screening (ELISA; Bioelisa HCV, Biokit

DEPARTAMENTO DE ESTOMATOLOGIA  
FACULTAD DE ODONTOLOGIA  
UNIVERSIDAD DE GRANADA  
Impulsado y conforme con el documento original.  
Granada, 29 NOV. 2001  
Fdo.: Francisco José Fernández Yáñez  
Isla del Negorrido