

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.e-jds.com

Original Article

Effect of amoxicillin and clindamycin on the gene expression of markers involved in osteoblast physiology

Francisco Javier Manzano-Moreno ^{a,b},
 Anabel González-Acedo ^{b,c}, Elvira de Luna-Bertos ^{b,d*},
 Enrique García-Recio ^{b,c}, Concepción Ruiz ^{b,d,e},
 Candela Reyes-Botella ^{a,b}

^a Biomedical Group (BIO277), Department of Stomatology, School of Dentistry, University of Granada, Granada, Spain

^b Instituto Investigación Biosanitaria, ibs.Granada, Spain

^c Biomedical Group (BIO277), Department of Nursing, Faculty of Health Science, University of Granada, Melilla, Spain

^d Biomedical Group (BIO277), Department of Nursing, Faculty of Health Science, University of Granada, Granada, Spain

^e Institute of Neuroscience, University of Granada, Granada, Spain

Received 27 June 2023; Final revision received 10 July 2023

Available online 18 July 2023

KEYWORDS

Low-speed drilling;
 Bone
 decontamination;
 Amoxicillin;
 Clindamycin;
 Osteoblasts

Abstract *Background/purpose:* Amoxicillin and clindamycin are the most effective decontaminants for intraoral bone grafts before their application in bone regeneration without cytotoxic effects on osteoblasts, but their effects on the gene expression of markers involved in osteoblast growth and differentiation remain unclear. The study objective was to determine the effects of amoxicillin and clindamycin on the gene expression of markers involved in osteoblast growth and differentiation.

Materials and methods: Real-time polymerase chain reaction (RT-PCR) was performed to explore the effect of 150 µg/mL clindamycin or 400 µg/mL amoxicillin on the gene expression by primary human osteoblasts (HOBs) of runt-related transcription factor 2 (Runx-2), osterix (OSX), alkaline phosphatase (ALP), osteocalcin (OSC), osteoprotegerin (OPG), receptor activator for nuclear factor κ B ligand (RANKL), type I collagen (Col-I), bone morphogenetic proteins 2 and 7 (BMP-2 and BMP-7), TGF-β1 and TGF-β receptors (TGF-βR1, TGF-βR2, and TGF-βR3), and vascular endothelial growth factor (VEGF).

Results: Treatment with 150 µg/mL clindamycin significantly increased the gene expression of

* Corresponding author. Department of Nursing, Faculty of Health Sciences, University of Granada, Avenida de la Ilustración s/n, Granada, 18016, Spain.

E-mail address: elviraadlb@ugr.es (E. de Luna-Bertos).

TFG- β 1, TGF- β R1, TGF- β R2, TGF- β R3, RUNX-2, Col-1, OSX, OSC, BMP-2, BMP-7, ALP, VEGF, and RANKL by HOBs. Treatment with 400 μ g/mL amoxicillin significantly increased the gene expression of TGF- β R1, Col-1, OSC, RANKL, and OPG alone.

Conclusion: These findings suggest that 150 μ g/mL clindamycin is the decontaminant of choice to treat intraoral bone grafts before their application in bone regeneration. The osteogenic and antibacterial properties of clindamycin can favor and accelerate the integration of bone grafts in the oral cavity.

© 2023 Association for Dental Sciences of the Republic of China. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Osseointegrated implants are frequently used in oral cavity rehabilitation. Techniques to regenerate lost bone volume are sometimes necessary to reestablish function. Bone substitutes are often used for this purpose, but autologous bone tissue remains the gold standard due to its unique characteristics, including its cellularity and morphogenetic proteins, which increase the potential for growth and integration of the graft in receptor tissue. In most cases, rehabilitation require a relatively small amount of graft material, and particulate bone grafts from intraoral localizations are used to achieve small increases in bone tissue that yield a functional bed for implants. One possible complication of this procedure is contamination by oral cavity microbiota during handling of the graft.^{1,2}

Studies comparing contamination among different methods of gathering intraoral particulate bone³ reported a lower bacterial count with the low-speed biological drilling of medullar bone tissue without irrigation.^{4,5} This technique has also been associated with the presence of viable osteoblasts with proliferative and mineralization capacity in the particulate bone.^{6,7}

Various decontaminants have been used to develop a safe prophylactic antimicrobial protocol for bone grafts.⁸ Our group previously compared the efficacy of two antibiotics (amoxicillin and clindamycin) and an antiseptic (chlorhexidine) on bone samples obtained by biological drilling and observed the greatest decontaminant effect with clindamycin, followed by chlorhexidine and amoxicillin.⁹ The risk of cytotoxicity posed by these decontaminant agents has been tested in fibroblasts, myoblasts, osteoblasts, and epithelial cells, studying their impact on the growth and differentiation of these cells in both fresh and cultured explants.^{10–14} Chlorhexidine was found to induce an abrupt decrease in proliferation and an increase in the percentage of apoptotic cells, whereas low-dose clindamycin increased cell proliferation, alkaline phosphatase activity, and calcification.^{15,16} Other researchers have published similar observations on the impact of these decontaminants on bone cells.^{17,18}

Amoxicillin and clindamycin are currently the two most effective decontaminants without cytotoxic effects on osteoblasts, and the objective of this study was to determine their effects on the gene expression of markers involved in osteoblast growth and differentiation.

Materials and methods

Tissues

To obtain the primary osteoblasts cell line, bone fragments were collected from volunteers during oral surgery. Volunteers were all university students aged 20–23 years from the School of Dentistry (University of Granada) requiring impacted third molar extraction at the School Clinic. The study was approved by the research ethics committee of the University (reference no. 3312/CEIH/2023). All participants signed their informed consent.

Isolation and culture of osteoblasts

The procedure described by Manzano-Moreno et al. (2013)⁶ was used to process the bone fragments. Primary osteoblasts were maintained in Dulbecco's modified Eagle medium (DMEM; Invitrogen Gibco Cell Culture Products, Carlsbad, CA, USA). After adding 100 IU/mL penicillin (Lab Roger SA, Barcelona, Spain), 2.5 mg/mL amphotericin B (Sigma, St. Louis, MO, USA), 50 mg/mL gentamicin (Braun Medical SA, Jaen, Spain), 1% glutamine (Sigma), and 2% HEPES (Sigma) supplemented with 10% fetal bovine serum (FBS; Gibco, Paisley, UK), cultures were kept in a humidified atmosphere at 37 °C with 95% air and 5% CO₂. Next, cells were detached from the flask with 0.05% trypsin (Sigma) and 0.02% ethylenediaminetetraacetic acid solution (EDTA; Sigma) and then rinsed and resuspended in culture medium with 10% FBS (Gibco).¹⁹ Previously published methods were used for the identification and characterization of osteoblasts.²⁰

Gene expression by real-time polymerase chain reaction (RT-PCR)

The methodology reported by Manzano-Moreno et al. (2018)²¹ was used to evaluate the effect of clindamycin and amoxicillin on osteoblast gene expression.

RNA extraction and cDNA synthesis (reverse transcription)

After 24 h of osteoblast culture with 150 μ g/mL clindamycin or 400 μ g/mL amoxicillin in DMEM without penicillin and gentamicin (untreated cells served as controls), mRNA was

Table 1 Primer sequences for the amplification of osteoblasts' cDNA by real-time PCR. Abbreviations: TGF- β 1, TGF- β R1, TGF- β R2, and TGF- β R3 (Transforming growth factor- β 1 and TGF- β receptors); Runx-2 (Runt-related transcription factor 2); VEGF (Vascular endothelial growth factor); OSX (Osterix); BMP-2 and BMP (Bone morphogenetic proteins 2 and 7); ALP (Alkaline phosphatase); Col-I (Type I collagen); OSC (osteocalcin); OPG (Osteoprotegerin); RANKL (Receptor activator for nuclear factor κ B ligand); UBC (Ubiquitin C); PPIA (peptidylprolyl isomerase A); RPS13 (Ribosomal protein S13).

Gene	Sense Primer	Antisense Primer	Amplicon (bp)
TGF- β 1	5'-TGAACCGGCTTTCTGCTTCTCATG-3'	5'-GCGGAAGTCAATGTACAGCTGCCGC-3'	152
TGF- β R1	5'-ACTGGCAGCTGTCATTGCTGGACCAG-3'	5'-CTGAGCCAGAACCTGACGTTGTCATATCA-3'	201
TGF- β R2	5'-GGCTCAACCACCAGGCATCCAGAT-3'	5'-CTCCCCGAGAGCCTGTCCAGATGCT-3'	139
TGF- β R3	5'-ACCGTGATGGGCATTGCGTTTGCA-3'	5'-GTGCTCTGCGTGCTGCCGATGCTGT-3'	173
RUNX-2	5'-TGGTTAATCTCCGAGGTCAC-3'	5'-ACTGTGCTGAAGAGGCTGTTTG-3'	143
VEGF	5'-CCTTGCTGCTCTACCTCCAC-3'	5'-CACACAGGATGGCTTGAAGA-3'	197
OSX	5'-TGCCTAGAAGCCCTGAGAAA-3'	5'-TTTAACTTGGGGCCTTGAGA-3'	205
BMP-2	5'-TCGAAATCCCCGTGACCAG-3'	5'-CCACTTCCACCACGAATCCA-3'	142
BMP-7	5'-CTGGTCTTTGTCTGCAGTGG-3'	5'-GTACCCCTCAACAAGGCTTC-3'	202
ALP	5'-CCAACGTGGCTAAGAATGTCATC-3'	5'-TGGGCATTGGTGTTGTACGTC-3'	175
COL-I	5'-AGAAGTGGTACATCAGCAAG-3'	5'-GAGTTTACAGGAAGCAGACA-3'	471
OSC	5'-CCATGAGAGCCCTCACACTCC-3'	5'-GGTCAGCCAACTCGTCACAGTC-3'	258
OPG	5'-ATGCAACACAGCACAAACATA-3'	5'-GTTGCCGTTTTATCCTCTCT-3'	198
RANKL	5'-ATACCTGATGAAAGGAGGA-3'	5'-GGGGCTCAATCTATATCTCG-3'	202
UBC	5'-TGGGATGCAAATCTTCGTGAAGACCCTGAC-3'	5'-ACCAAGTGAGAGTGGACTCTTTCTGGATG-3'	213
PPIA	5'-CCATGGCAAATGCTGGACCCAAACAAATG-3'	5'-TCCTGAGCTACAGAAGGAATGATCTGGTGG-3'	256
RPS13	5'-GGTGTGACACAAGTACGTTTGTGACAGGC-3'	5'-TCATATTTCCAATTGGGAGGGAGGACTCGC-3'	251

extracted by a silicate gel technique included in the Qia-genRNeasy extraction kit (Qiagen Inc., Hilden, Germany), which involves a DNase digestion step. The amount of extracted mRNA was measured by UV spectrophotometry at

260 nm (Eppendorf AG, Hamburg, Germany), and contamination with proteins was determined according to the 260/280 ratio. An equal amount of RNA (1 μ g of total RNA in 40 μ L of total volume) was reverse-transcribed to cDNA and




Figure 1 Expression of osteoblast genes (TGF- β 1, TGF- β R1, TGF- β R2, and TGF- β R3) treated for 24 h with 150 μ g/mL clindamycin or 400 μ g/mL amoxicillin. Data are expressed as ng of mRNA per average ng of housekeeping mRNAs $\pm$ SD. TGF- β 1, TGF- β R1, TGF- β R2, and TGF- β R3 (Transforming growth factor- β 1 and TGF- β receptors). * P < 0.05, ** P < 0.001.

amplified by PCR using the iScript™ cDNA Synthesis Kit (Bio-Rad laboratories, Hercules, CA, USA) following the manufacturer's instructions.

Real-time polymerase chain reaction (RT-PCR)

The NCBI- nucleotide library and Primer3-design were used to design primers to detect mRNA of the following genes: runt-related transcription factor 2 (Runx-2), osterix (OSX), alkaline phosphatase (ALP), osteocalcin (OSC), osteoprotegerin (OPG), receptor activator for nuclear factor κ B ligand (RANKL), type I collagen (Col-I), bone morphogenetic proteins 2 and 7 (BMP-2 and BMP-7), TGF- β 1 and TGF- β receptors (TGF- β R1, TGF- β R2, and TGF- β R3), and vascular endothelial growth factor (VEGF). According to different authors,^{21–23} it is recommended to use a minimum number of three house-keeping genes. In the case of bone tissue and osteoblast, results were normalized using ubiquitin C (UBC), peptidylprolyl isomerase A (PPIA), and ribosomal protein S13 (RPS13) as housekeeping genes. Primer sequences are given in Table 1.

Quantitative RT-PCR (q-RT-PCR) was performed with the SsoFast™ EvaGreen® Supermix Kit (Bio-Rad laboratories) following the manufacturer's instructions. Samples were amplified in 96-well microplates in an IQ5-Cycler (Bio-Rad laboratories) at a specific annealing temperature for each gene (range, 60 to 65 °C) and an elongation temperature of 72 °C over 40 cycles. PCR reactions were carried out in a final volume of 20 μ L, with 5 μ L of cDNA sample and 2 μ L of each primer. Ct values were plotted against log cDNA dilution to construct standard curves for each target gene. A melting profile was created after each RT-PCR, conducting agarose gel electrophoresis in samples to rule out nonspecific PCR products and primer dimers. The comparative Ct method was applied for the relative quantification of gene expression, and the mRNA concentration of each gene was expressed as ng of mRNA per average ng of housekeeping mRNAs. The cDNA from individual cell experiments was analyzed in triplicate RT-PCR studies.

Statistical analysis

SPSS 26.0 (IBM, Chicago, IL, USA) was used for data analyses. mRNA levels were expressed as means $\pm$ standard deviation (SD). A two-way repeated-measures analysis of variance (ANOVA) was performed, applying the Bonferroni correction when significant interactions were identified. A minimum of three experiments were performed for each assay. $P < 0.05$ was considered significant.

Results

Effect of clindamycin and amoxicillin on gene expression of TGF- β 1 and its receptors (TGF- β R1, TGF- β R2, and TGF- β R3)

Fig. 1 depicts q-RT-PCR results for the gene expression of TGF- β 1 and its receptors (TGF- β R1, TGF- β R2, and TGF- β R3). TGF- β 1 expression was significantly increased *versus*




Figure 2 Expression of osteoblast genes (RANKL and OPG) treated for 24 h with 150 μ g/mL clindamycin or 400 μ g/mL amoxicillin. Data are expressed as ng of mRNA per average ng of housekeeping mRNAs $\pm$ SD. RANKL (Receptor activator for nuclear factor κ B ligand); OPG (Osteoprotegerin). * $P < 0.05$, ** $P < 0.001$.

controls after 24 h of treatment with 150 μ g/mL clindamycin ($P < 0.001$) but not 400 μ g/mL amoxicillin. Treatment with 150 μ g/mL clindamycin significantly increased the expression of TGF- β R1, TGF- β R2, and TGF- β R3 *versus* controls ($P < 0.001$), whereas treatment with 400 μ g/mL amoxicillin significantly increased the expression of TGF- β R1 alone ($P = 0.023$).

Effect of amoxicillin and clindamycin on gene expression of RANKL-OPG complex

Fig. 2 shows q-RT-PCR results for the gene expression of RANKL and OPG. Treatment for 24 h with 150 μ g/mL clindamycin or 400 μ g/mL amoxicillin significantly increased the expression of RANKL *versus* controls ($P < 0.001$ and 0.043, respectively). In addition, the expression of OPG was significantly increased by treatment with 150 μ g/mL clindamycin and 400 μ g/mL amoxicillin ($P = 0.024$ and $P = 0.004$, respectively).

Effect of amoxicillin and clindamycin on the gene expression of Runx2, ALP, OSX, Col-I and OSC

Fig. 3 displays q-RT-PCR results for the expression of osteoblast differentiation markers Runx2, ALP, OSX, Col-I, and OSC. Treatment with 150 µg/mL clindamycin for 24 h significantly increased the expression of Runx2, ALP, OSX, Col-I, and OSC *versus* controls ($P < 0.001$), and treatment with 400 µg/mL amoxicillin significantly increased the expression of Col-I and OSC ($P < 0.001$), whereas neither treatment significantly altered the expression of Runx2, ALP, or OSX.

Effect of amoxicillin and clindamycin on gene expression of BMP-2, BMP-7, and VEGF

Fig. 4 exhibits q-RT-PCR results for the gene expression of BMP-2, BMP-7, and VEGF. Expression of BMP-2, BMP-7, and VEGF was significantly increased *versus* controls after 24 h of treatment with 150 µg/mL clindamycin ($P < 0.001$) but not 400 µg/mL amoxicillin.

Discussion

Amoxicillin and clindamycin, widely used to decontaminate and avoid infectious complications after autologous bone grafting, were found to differ in their effects on the gene expression of osteoblasts. In this way, 150 µg/mL clindamycin significantly increased the expression of genes involved in osteoblast growth (TGF-β1, TGF-βR1, TGF-βR2, TGF-βR3, and VEGF) and differentiation (RUNX-2, Col-1, OSX, OSC, BMP-2, BMP-7, or ALP) *versus* controls, whereas 400 µg/mL amoxicillin only increased the expression of TGF-β R1, Col-1, and OSC. Both treatments modified gene expressions related to osteoblast-osteoclast interactions (RANKL-OPG). We selected these treatment doses/time because in a previous study by our group,¹⁶ we demonstrated that treatment of osteoblasts with 150 µg/mL clindamycin and 400 µg/mL amoxicillin for 24 h did not have a toxic effect on osteoblasts.

The TGF-β superfamily has more than 40 members, including TGF-βs, Nodal, Activin, and BMPs. TGF-β signaling is critical for the regulation of osteoblast differentiation and bone formation, and signaling relays at each stage leads to the final target gene expression. Low to moderately-elevated levels of TGF-β1 were found to stimulate early osteoblast proliferation, and this growth factor is involved in activating the MAPK cascade.^{24,25} The MAPK signaling pathway is known to participate in the regulation of cell growth, development, and division through its influence on nuclear transcription factors^{24,26} and in the activation and signaling of other osteogenic growth factors.^{24,27} The expression of TGF-β1 and its receptors was significantly increased by treatment with 150 µg/mL clindamycin, which may be related to the increased proliferation of human osteoblasts observed by Olvera-Huertas et al. (2022)¹⁶ after treatment with this antibiotic at the same dose. Other authors have reported that a low dose of clindamycin is effective to decontaminate different types of bone grafts or extraction sockets

without impairing the viability, metabolic activity, or proliferation of osteoblasts. However, no evidence has been published on its impact on the gene expression of osteoblast growth and differentiation markers.

There are two main steps in the complex process of osteoblast differentiation: i) osteogenic commitment, characterized by upregulation of the transcription factor Runx2²⁸; and ii) osteogenic differentiation, characterized by expression of the specific osteogenic proteins and transcription factors ALP, OPG, and OSX.^{29,30} Among these, OSX is an essential transcription factor for osteoblast differentiation and bone formation, and its expression indicates the loss of bipotentiality from preosteoblast to osteoblast and chondrocyte. In the present study, treatment with 150 µg/mL clindamycin induced upregulation of the osteoblast expression of Runx-2, ALP, and OSX, which would be related to an increase in osteoblast differentiation, consistent with the by Olvera-Huertas et al. (2022)¹⁶ of higher ALP expression and mineralization after the same treatment.

BMPs have multiple effects on osteoblast proliferation, differentiation, and general physiology. For instance, BMP-2 and BMP-7 have important roles in bone formation/remodeling and development and in osteoblast differentiation, inducing the expression of ALP and other osteoblastic markers and promoting calcium mineralization. In the present study, treatment with 150 µg/mL clindamycin significantly increased the gene expression of BMP-2 and BMP-7 ($P < 0.001$), indicating the activation of osteogenesis.³¹ These findings suggest that stimulation of the BMP-Smad pathway may induce an osteogenic effect that is further sustained by Wnt signaling.^{32,33}

Treatment with clindamycin also increased gene expressions of OSC and COL-I, osteogenic markers of the final stages of osteoblast differentiation that appear at the start of mineralization.

The expression and production of pro-angiogenic factors such as VEGF and angiopoietin (ANG) affect endothelial cell growth, migration, and vessel formation in many tissues and are important for regulating vascular growth in the skeleton. These factors are produced by osteoblasts and osteocytes, among other bone cells, and the present study shows an increased gene expression of VEGF by osteoblasts treated with clindamycin; which may therefore favor angiogenesis in bone tissue.

Treatment with 400 µg/mL amoxicillin was recently found to be an effective decontaminant of intraoral bone grafts and only increased the expression of TGF-β R1, Col-1, and OSC in the present study. Rathbone et al.³² observed a moderate reduction of 26–49% in osteoblastic cell count and alkaline phosphatase activity after treatment with amoxicillin at a dose over 500 µg/mL and a reduction of >75% in cell count at a dose of 2000 g/mL or higher. Selection of the lower dose of 400 µg/mL is designed to maintain the decontaminating efficacy of amoxicillin without impairing the functional capacity of osteoblasts.

RANKL and OPG regulate bone remodeling by respectively stimulating and inhibiting osteoclast differentiation, and they have been found to participate in the complex set of molecular interactions leading to bone loss in peri-implantitis.³⁴ Local deregulation of the RANKL-OPG complex in favor of the pro-osteoclastic component may




Figure 3 Expression of osteoblast genes (RUNX-2, ALP, Col-I, OSX and OSC) treated for 24 h with 150 µg/mL clindamycin or 400 µg/mL amoxicillin. Data are expressed as ng of mRNA per average ng of housekeeping mRNAs $\pm$ SD. Runx-2 (Runt-related transcription factor 2); ALP (Alkaline phosphatase); Col-I (Type I collagen); OSX (Osterix); OSC (osteocalcin). ** $P < 0.001$.

lead to bone resorption at the alveolar site³⁵ and increase the risk and severity of peri-implant disease.³⁶ In the present study, treatment with the two antibiotics (amoxicillin and clindamycin) produced significant changes in the gene expression of RANKL-OPG in osteoblasts, modulating the interaction of osteoblasts and osteoclasts and potentially altering the bone apposition-resorption process.

The antibiotic clindamycin appears to be a good option for the decontamination of intraoral bone grafts, offering a higher decontaminating capacity in comparison to other

agents⁹ as well as increasing the proliferation and differentiation of osteoblasts.¹⁶ These effects are explained by analysis of its impact on the gene expression of markers with a key role in osteoblast physiology, as revealed in the present study.

According to these findings, low-dose clindamycin is the agent of choice to decontaminate intraoral bone grafts before their utilization in bone regeneration. Its osteogenic and antibacterial properties may favor and accelerate the integration of bone grafts in the oral cavity.




Figure 4 Expression of osteoblast genes (BMP-2, BMP-7, and VEGF) treated for 24 h with 150 µg/mL clindamycin or 400 µg/mL amoxicillin. Data are expressed as ng of mRNA per average ng of housekeeping mRNAs $\pm$ SD. BMP-2 and BMP (Bone morphogenetic proteins 2 and 7); VEGF (Vascular endothelial growth factor). ** $P < 0.001$.

Declaration of competing interest

The authors declare that they have no conflict of interest.

Acknowledgements

This study was supported by BIO-277 research group (Junta de Andalucía, Spain).

References

1. Pape HC, Evans A, Kobbe P. Autologous bone graft: properties and techniques. *J Orthop Trauma* 2010;24(Suppl 1):S36–40.
2. Rogers GF, Greene AK. Autogenous bone graft: basic science and clinical implications. *J Craniofac Surg* 2012;23:323–7.
3. Takamoto M, Takechi M, Ohta K, et al. Risk of bacterial contamination of bone harvesting devices used for autogenous bone graft in implant surgery. *Head Face Med* 2013;9:3.
4. Hashemi HM, Beshkar M. Bacterial contamination of autogenous bone collected by rongeur compared with that collected by bone filter during implant surgery. *Br J Oral Maxillofac Surg* 2011;49:474–7.
5. Manzano-Moreno FJ, Herrera-Briones FJ, Linares-Recatala M, Ocaña-Peinado FM, Reyes-Botella C, Vallecillo-Capilla MF. Bacterial contamination levels of autogenous bone particles collected by 3 different techniques for harvesting intraoral bone grafts. *J Oral Maxillofac Surg* 2015;73:424–9.
6. Manzano-Moreno FJ, Rodríguez-Martínez JB, Ramos-Torrecillas J, et al. Proliferation and osteogenic differentiation of osteoblast-like cells obtained from two techniques for harvesting intraoral bone grafts. *Clin Oral Invest* 2013;17:1349–56.
7. Liang C, Lin X, Wang SL, Guo LH, Wang XY, Li J. Osteogenic potential of three different autogenous bone particles harvested during implant surgery. *Oral Dis* 2017;23:1099–108.
8. Tezulas E, Dilek OC, Topcuoglu N, Kulekci G. Decontamination of autogenous bone grafts collected during dental implant site preparation: a pilot study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;107:656–60.
9. Olvera-Huertas AJ, Linares-Recatalá M, Herrera-Briones FJ, Vallecillo-Capilla MF, Manzano-Moreno FJ, Reyes-Botella C. Microbiological analysis of autogenous bone particles obtained by low-speed drilling and treated with different decontamination agents. *Int J Oral Maxillofac Surg* 2021;50:104–8.
10. Almazin SM, Dziak R, Andreana S, Ciano SG. The effect of doxycycline hyclate, chlorhexidine gluconate, and minocycline hydrochloride on osteoblastic proliferation and differentiation in vitro. *J Periodontol* 2009;80:999–1005.
11. Verdugo F, Sáez-Rosón A, Uribarri A, et al. Bone microbial decontamination agents in osseous grafting: an in vitro study with fresh human explants. *J Periodontol* 2011;82:863–71.

12. Faria G, Cardoso CRB, Larson RE, Silva JS, Rossi MA. Chlorhexidine-induced apoptosis or necrosis in L929 fibroblasts: a role for endoplasmic reticulum stress. *Toxicol Appl Pharmacol* 2009;234:256–65.
13. John G, Becker J, Schwarz F. Effects of taurolidine and chlorhexidine on SaOS-2 cells and human gingival fibroblasts grown on implant surfaces. *Int J Oral Maxillofac Implants* 2014;29:728–34.
14. Liu JX, Werner J, Kirsch T, Zuckerman JD, Virk MS. Cytotoxicity evaluation of chlorhexidine gluconate on human fibroblasts, myoblasts, and osteoblasts. *J Bone Jt Infect* 2018;3:165–72.
15. Brunello G, Becker K, Scotti L, Drescher D, Becker J, John G. The effects of three chlorhexidine-based mouthwashes on human osteoblast-like SaOS-2 cells. An in vitro study. *Int J Mol Sci* 2021;22:9986.
16. Olvera-Huertas AJ, Costela-Ruiz VJ, García-Recio E, et al. The effect of chlorhexidine, amoxicillin, and clindamycin on the growth and differentiation of primary human osteoblasts. *Int J Oral Maxillofac Implants* 2022;37:283–8.
17. Cabral CT, Fernandes MH. In vitro comparison of chlorhexidine and povidone-iodine on the long-term proliferation and functional activity of human alveolar bone cells. *Clin Oral Invest* 2007;11:155–64.
18. Naal FD, Salzmann GM, von Knoch F, et al. The effects of clindamycin on human osteoblasts in vitro. *Arch Orthop Trauma Surg* 2008;128:317–23.
19. George J, Klika AK, Higuera CA. Use of chlorhexidine preparations in total joint arthroplasty. *J Bone Jt Infect* 2017;2:15–22.
20. Reyes-Botella C, Montes MJ, Vallecillo-Capilla MF, Olivares EG, Ruiz C. Antigenic phenotype of cultured human osteoblast-like cells. *Cell Physiol Biochem* 2002;12:359–64.
21. Manzano-Moreno FJ, Ramos-Torrecillas J, Melguizo-Rodríguez L, Illescas-Montes R, Ruiz C, García-Martínez O. Bisphosphonate modulation of the gene expression of different markers involved in osteoblast physiology: possible implications in bisphosphonate-related osteonecrosis of the jaw. *Int J Med Sci* 2018;15:359–67.
22. Vandesompele J, De Preter K, Pattyn F, et al. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 2002;3. RESEARCH0034.
23. Ragni E, Viganò M, Rebullà P, Giordano R, Lazzari L. What is beyond a qRT-PCR study on mesenchymal stem cell differentiation properties: how to choose the most reliable house-keeping genes. *J Cell Mol Med* 2013;17:168–80.
24. Xiao G, Jiang D, Gopalakrishnan R, Franceschi RT. Fibroblast growth factor 2 induction of the osteocalcin gene requires MAPK activity and phosphorylation of the osteoblast transcription factor, Cbfa1/Runx2. *J Biol Chem* 2002;277:36181–7.
25. Shen B, Wei A, Whittaker S, et al. The role of BMP-7 in chondrogenic and osteogenic differentiation of human bone marrow multipotent mesenchymal stromal cells in vitro. *J Cell Biochem* 2010;109:406–16.
26. Seger R, Krebs EG. The MAPK signaling cascade. *FASEB J* 1995;9:726–35.
27. Lee KS, Hong SH, Bae SC. Both the Smad and p38 MAPK pathways play a crucial role in Runx2 expression following induction by transforming growth factor-beta and bone morphogenetic protein. *Oncogene* 2002;21:7156–63.
28. Xu J, Li Z, Hou Y, Fang W. Potential mechanisms underlying the Runx2 induced osteogenesis of bone marrow mesenchymal stem cells. *Am J Transl Res* 2015;7:2527–35.
29. Klejna K, Naumnik B, Gasowska K, Myśliwiec M. OPG/RANK/-RANKL signaling system and its significance in nephrology. *Folia Histochem Cytobiol* 2009;47:199–206.
30. Nakashima K, Zhou X, Kunkel G, et al. The novel zinc finger-containing transcription factor osterix is required for osteoblast differentiation and bone formation. *Cell* 2002;108:17–29.
31. Groeneveld EH, Burger EH. Bone morphogenetic proteins in human bone regeneration. *Eur J Endocrinol* 2000;142:9–21.
32. Gregory CA, Gunn WG, Reyes E, et al. How Wnt signaling affects bone repair by mesenchymal stem cells from the bone marrow. *Ann N Y Acad Sci* 2005;1049:97–106.
33. Hamidouche Z, Haÿ E, Vaudin P, et al. FHL2 mediates dexamethasone-induced mesenchymal cell differentiation into osteoblasts by activating Wnt/beta-catenin signaling-dependent Runx2 expression. *FASEB J* 2008;22:3813–22.
34. Duarte PM, de Mendonça AC, Máximo MBB, Santos VR, Bastos MF, Nociti FH. Effect of anti-infective mechanical therapy on clinical parameters and cytokine levels in human peri-implant diseases. *J Periodontol* 2009;80:234–43.
35. Bostanci N, Ilgenli T, Emingil G, et al. Gingival crevicular fluid levels of RANKL and OPG in periodontal diseases: implications of their relative ratio. *J Clin Periodontol* 2007;34:370–6.
36. Duarte PM, de Mendonça AC, Máximo MBB, Santos VR, Bastos MF, Nociti Júnior FH. Differential cytokine expressions affect the severity of peri-implant disease. *Clin Oral Implants Res* 2009;20:514–20.