

Research Article

Gradual Sets: An Approach to Fuzzy Sets

Josefa M. García ¹ and Pascual Jara ²

¹Department of Applied Mathematics, University of Granada, E-18071 Granada, Spain

²Department of Algebra and IMAG (Instituto de Matemáticas), University of Granada, E-18071 Granada, Spain

Correspondence should be addressed to Pascual Jara; pjara@ugr.es

Received 16 November 2021; Accepted 27 September 2022; Published 27 February 2023

Academic Editor: Antonin Dvorák

Copyright © 2023 Josefa M. García and Pascual Jara. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the fuzzy theory of sets and groups, the use of α -levels is a standard to translate problems from the fuzzy to the crisp framework. Using strong α -levels, it is possible to establish a one to one correspondence which makes possible doubly, a gradual and a functorial treatment of the fuzzy theory. The main result of this paper is to identify the class of fuzzy sets, respectively, fuzzy groups, with subcategories of the functorial categories $\text{Set}^{(0,1)}$, resp., $\text{Gr}^{(0,1)}$. In this line, the algebraic potential of this theory will be reached, in forthcoming papers.

1. Introduction

1.1. Gradual Elements. Let X be a set, every subset $S \subseteq X$ is defined by its characteristic function $\chi_S: X \rightarrow \{0, 1\}$, which

is defined by $\chi_S(x) = \begin{cases} 1, & \text{if } x \in S, \\ 0, & \text{if } x \notin S. \end{cases}$ Thus, the concept of

membership is exclusive. However, we can consider, in a wider environment, different degrees of membership: 1 means that the element belongs to the subset; 0 means that it does not belong, and any other real number $0 < \alpha < 1$ would mean a different degree of membership.

The theory, in these terms, is due to Zadeh (see [1]) who introduces a fuzzy subset of a given set X as a map $\mu: X \rightarrow [0, 1]$. From this primitive concept, we can develop a whole theory of sets, relations, maps, numbers, and so on.

In this approach to the fuzzy theory, we begin by relating various mathematical theories; this relationship is evident in the crisp framework, but which in the fuzzy theory presents, so far, some difficulties.

Our approach to the fuzzy concept starts from the definition of a fuzzy element: we adopt the definition given by Dubois and Prade in [2] of a gradual element (if X is set, a gradual element in X is a partial map $e: (0, 1) \rightarrow X$); see also [3] in which gradual elements are applied to gradual numbers or [4] in which they are applied to build gradual

intervals so that a gradual element of a set X is given by a collection of elements of X , each with a degree of membership, ranging in $(0, 1]$: there is always an element of the set X that has a degree of membership 1, and, possibly, other elements with other membership values, but never 0; that is, we do not determine any element of X that has zero degree of membership. This notion of gradual element has been extended to study several problems (see [5]). Let X be a set. A representation level (RL) of a fuzzy concept on X is a partial map $e: [0, 1] \rightarrow \mathcal{P}(X)$ (see also [6, 7]). Nevertheless, we have preferred to maintain the former one as when applying to sets, groups, and other structures, it defines canonically a ground set, group, and so on which is an ambient object suitable for working.

For a greater flexibility in the definition, we may assume that not all possible degrees are reached, so a gradual element is given by a partial mapping from $(0, 1]$ to X : we will call it a partial gradual element. If e is a partial gradual element with definition domain $\text{dom}(e) \subseteq (0, 1]$, for its study, we need to relate partial gradual elements to each other. The problem that arises is when two gradual elements e_1 and e_2 are equal? It is clear that we can only compare e_1 and e_2 where they are defined, that is, in $\text{dom}(e_1) \cap \text{dom}(e_2)$.

This definition of equality is too weak. In fact, we are more interested in knowing if e_1 and e_2 take equal values in a

range $[\alpha, 1]$, for some $\alpha \in (0, 1]$. Taking into account that whenever $\beta \in (0, 1]$ is very small, it is not relevant at all if that $\varepsilon_1(\beta)$ and $\varepsilon_2(\beta)$ are the same or different; we are more interested in knowing whenever ε_1 and ε_2 coincide for values of β close to 1.

Thus, we extend the equality relation to the case, previously indicated, of values in an interval $[\alpha, 1]$. In this way, a relationship is obtained: $\varepsilon_1 R_\alpha \varepsilon_2$ if

$$\varepsilon_{1|[\alpha,1] \cap \text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2)} = \varepsilon_{2|[\alpha,1] \cap \text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2)}. \quad (1)$$

However, this relation is not necessarily an equivalence relation because it depends heavily on $\text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2)$. So, if we want to compare partial gradual elements, we must standardize the definition domain. In other words, we must, for instance, extend $\text{dom}(\varepsilon)$ to the whole $(0, 1]$.

There is a standard method of doing this, consisting of, given $\alpha \in \text{dom}(\varepsilon)$ such that ε is not defined in (β, α) , defining $\varepsilon(\gamma) = \varepsilon(\alpha)$ for all $\gamma \in (\beta, \alpha)$. The condition that has seemed most efficient forces to restrict the partial gradual elements to those whose definition domain verifies that for every $\alpha \in (0, 1]$, there is a minimum ζ of $[\alpha, 1] \cap \text{dom}(\varepsilon)$, to, in this way, extend ε to all $(0, 1]$, defining $\varepsilon(\alpha) = \varepsilon(\zeta)$. We have called inf-compact the subsets of $(0, 1]$ containing 1 and verifying this property. In this way, every partial gradual element ε , with inf-compact definition domain, can be extended, in a unique way, to a gradual element $\bar{\varepsilon}$ with definition domain $(0, 1]$. We call $\bar{\varepsilon}$ the extended gradual element of ε .

We define a total gradual element as a map $\varepsilon: (0, 1] \rightarrow X$, among which we have the extended of the partial gradual elements; and denote by \mathcal{X} the set of all total gradual elements of X . Observe that, when working with total gradual elements, the relation R_α is an equivalence relation.

1.2. Gradual Subsets and Gradual Subgroups. The next step of complexity is to consider a binary operation $*$ in the set X and extend it to gradual elements. The standard method is to define $(\varepsilon_1 * \varepsilon_2)(\alpha) = \varepsilon_1(\alpha) * \varepsilon_2(\alpha)$ for any $\alpha \in (0, 1]$.

We have that if $(X, *)$ has a more complex algebraic structure, for example, if it is a group, a semilattice, the set \mathcal{X} of all gradual elements can have the same property. However, this has not been the line we followed for the study of fuzzy structures in a set \mathcal{X} , the reason is that when considering, for example, a ring structure in \mathcal{X} , although X has a ring structure, this is of little interest, since it has too many zero-divisor elements.

We have chosen therefore to consider a greater degree of abstraction and consider, given a group $(G, *)$, not the set of elements of G , but the set $\mathcal{S}(G)$ of all the subgroups of G . We have an inclusion $\mathcal{S}(G) \subseteq \mathcal{P}(G)$, in the powerset of G , and the elements of $\mathcal{S}(G)$ are the nonempty subsets $\mathcal{P} \in \mathcal{P}(G)$ verifying: $S * S \subseteq S$ and $S^{-1} \subseteq S$. When considering gradual elements $\sigma, \sigma_1, \sigma_2$ of $\mathcal{P}(G) \setminus \{\emptyset\}$, we have new gradual elements: $\sigma_1 * \sigma_2$ and σ^{-1} , and naturally the notions of gradual subgroup and gradual subset appear.

A gradual subset of a set X is a gradual element σ of $\mathcal{P}(X)$, and a gradual subgroup of a group G is a gradual element σ of $\mathcal{P}(G) \setminus \{\emptyset\}$ which is a subgroup, i.e., it will be a gradual element of $\mathcal{S}(G)$. Compare with the notion of RL (representation by levels) introduced in [5].

Observe that in these situations, we have solved the problem of extending partial gradual subsets or subgroups because we can define the image of any element in $(0, 1] \setminus \text{dom}(\sigma)$ equals either \emptyset , for subsets, or $\{e\}$, for subgroups. Therefore, in Sections 3 and 4, we shall consider only total gradual subsets and subgroups.

This study will lead us to relate gradual subgroups with fuzzy subgroups and gradual groups with fuzzy groups, and the same process will allow to relate other structures: rings, modules, and so on.

In our exposition, we will try to establish a framework, for future developments, based on a categorical structure that allows to consider, not only gradual sets but also gradual algebraic and geometrical structures. Before carrying out this work, we have considered necessary to implement an in-depth study that relates gradual and fuzzy sets and subsets.

In the set \mathcal{X} of the gradual subsets of X , we define a closure operator $\sigma \mapsto \sigma^c = \cup \{\sigma(\beta) | \beta \geq \alpha\}$. A gradual subset σ will be a decreasing gradual subset if $\sigma = \sigma^c$. And in the set $\mathcal{F}(X)$ of all decreasing gradual subsets of X , we define an interior operator $\sigma \mapsto \sigma^d = \cup \{\sigma(\beta) | \beta > \alpha\}$; a decreasing gradual subset σ will be a strict decreasing gradual subset whenever $\sigma = \sigma^d$.

Associated to any fuzzy subset μ of X , we have a decreasing gradual subset $\sigma(\mu)$, defined $\sigma(\mu)(\alpha) = \mu_\alpha$, the α -level of μ , for any $\alpha \in (0, 1]$, and a strict decreasing gradual subset $\bar{\sigma}(\mu) = \sigma(\mu)^d$, which is the strong α -level or strong α -cut, of μ . The map $\mu \mapsto \sigma(\mu)$ does not preserve unions of infinite families, and the map $\mu \mapsto \bar{\sigma}(\mu)$ does not preserve infinite intersections; hence, after modifying the intersection, we establish an injective correspondence, preserving union and intersection, from fuzzy subsets to strict decreasing gradual subsets, and find conditions on strict decreasing gradual subsets to be in the image of this map; that condition is property (inf-F). Which is important, in this situation, is that we have an isomorphism, for intersections and unions, between fuzzy subsets and strict decreasing gradual subsets satisfying property (inf-F). As a consequence, properties on fuzzy subsets can be studied via strict decreasing gradual subsets.

In addition, we consider a generalization of the theory of gradual subsets through the use of contravariant functors from the category $(0, 1]$ to the category Set of sets which allow a functorial framework of both theories of gradual subsets and of fuzzy subsets.

This theory, first developed in a context of sets, can be carried out to the more algebraic framework of groups, in which we may establish also a bijection between fuzzy subgroups and a specific class of gradual subgroups and contravariant functors. In particular, this bijection will allow a functorial treatment of fuzzy groups.

This paper is organized in four sections. In the second one, we study and establish the general theory of gradual elements and introduce binary operations in the set of all

gradual elements defined from binary operation in the ground set X . In particular, if we start from a group G , we get a structure of group in the set of gradual elements. Not in all cases, this structure reflects the properties of G and its elements.

For this reason, to make an algebraic development later, in the third section, we study gradual subsets and operators in the set of gradual subsets that will allow to establish a close relationship, an isomorphism, between the set of fuzzy subsets and a set of gradual subsets. This study ends in Theorem 2 in which an isomorphism is established; observe that to obtain the isomorphism, we had to make use of the strict decreasing gradual subsets. To do that, first we consider binary operations in $\mathcal{P}(X)$, the power set of X : the standard ones are the meet (intersection) and the join (union) and translate them to gradual subsets, which are noting more than gradual elements of $\mathcal{P}(X)$. In this section, we also identify a new type of objects through the use of contravariant functors from the category $(0, 1]$ to the category of sets. These contravariant functors, which are identified with directed systems, generalize gradual subsets and fuzzy subsets and allow a functorial framework of these two examples, which will provide a tool capable of dealing with other types of gradual and fuzzy objects such as groups, rings, and so on, and that will allow to work, by using direct limits, with gradual and fuzzy sets, instead of with gradual and fuzzy subsets.

The fourth section is devoted to study the more complex example of gradual groups. After studying the different concepts related to group theory, we establish the most important result, Theorem 2, showing a bijection between equivalence classes of fuzzy subgroups and some specific strict gradual subgroups. This gradual subgroups appear in a natural way after studying two operator on gradual subgroups, one a closure operator and another one an interior operator in the class of all decreasing gradual groups. The formulation of the theory in terms of operators allows to develop a more abstract framework, in this case a functorial one, and hence obtain new properties and relationships between known objects.

2. Gradual Elements

2.1. Definition of Gradual Elements

Definition 1. Let X be a set, a total gradual element of X is a map $\varepsilon: (0, 1] \rightarrow X$, and a partial gradual element of X is a map $\varepsilon: L \subseteq (0, 1] \rightarrow X$, defined on a subset $L \subseteq (0, 1]$ such that $1 \in L$. For simplicity, depending of the context, we use gradual element to refer either to a total gradual element or to a partial gradual element.

For any partial gradual element $\varepsilon: L \subseteq (0, 1] \rightarrow X$, we call L the definition domain of ε and represent it by $\text{dom}(\varepsilon)$. We represent by \mathcal{X} the set of all total gradual elements of X and by $\underline{\mathcal{X}}$ the set of all partial gradual elements.

A gradual element ε' is an extension of the gradual element ε if $\varepsilon'_{|\text{dom}(\varepsilon)} = \varepsilon$.

There is a particularly useful method of extending a partial gradual element ε to a total gradual one; this is the case in which for any $\alpha \in (0, 1]$, there exists $\text{Min}([\alpha, 1] \cap \text{dom}(\varepsilon))$; then, we define a new gradual element $\bar{\varepsilon}$ as follows:

$$\bar{\varepsilon}(\alpha) = \varepsilon(\zeta), \text{ being } \zeta = \text{Min}([\alpha, 1] \cap \text{dom}(\varepsilon)). \quad (2)$$

See Example 2 in which examples of extensions of partial gradual elements appear. Another example is provided whenever we consider the partial gradual element $\varepsilon: \{1\} \rightarrow \{a, b\}$, defined by $\varepsilon(1) = a$. In this case, an extension $\bar{\varepsilon}: (0, 1] \rightarrow \{a, b\}$ is defined by $\bar{\varepsilon}(\alpha) = a$ for any $\alpha \in (0, 1]$; the constant map equals to a .

A subset $L \subseteq (0, 1]$, containing 1, such that there exists $\text{Min}([\alpha, 1] \cap L)$, for any $\alpha \in (0, 1]$, is named an inf-compact subset of $(0, 1]$.

The following are examples of inf-compact subsets of $(0, 1]$:

- (1) Any compact subset $C \subseteq (0, 1]$, containing 1, is inf-compact. In particular, any finite subset and any closed (in \mathbb{R}) subset of $(0, 1]$, containing 1, are inf-compact
- (2) Any ascending sequence in $(0, 1]$, union with $\{1\}$, is inf-compact
- (3) Any interval $[a, b] \subseteq (0, 1]$, union with $\{1\}$, is inf-compact
- (4) Any union of finitely many inf-compact subsets is inf-compact

In the following, the domain of any partial gradual element will be an inf-compact subset of $(0, 1]$, containing 1; hence, any partial gradual element can be extended to a total gradual element.

Lemma 1. *If $\{C_i | i \in I\}$ is a family of inf-compact subsets, containing 1, then $\cap_i C_i$ is inf-compact.*

Proof. For any $\alpha \in (0, 1]$, let $\xi = \text{Inf}([\alpha, 1] \cap (\cap_i C_i))$, and define $\xi_i = \text{Min}([\xi, 1] \cap C_i)$; hence $\xi \leq \xi_i$, for any $i \in I$. On the other hand, since $[\alpha, 1] \cap (\cap_i C_i) = [\xi, 1] \cap (\cap_i C_i) \subseteq [\xi, 1] \cap C_i$, then $\text{Min}([\xi, 1] \cap C_i) = \text{Inf}([\xi, 1] \cap C_i) \leq \text{Inf}([\xi, 1] \cap (\cap_i C_i)) = \xi$. In consequence, $\xi = \xi_i$ for any $i \in I$.

For any element $a \in X$, there exists a partial gradual element, which we represent by ε_a , with $\text{dom}(\varepsilon_a) = \{1\}$, and defined by $\varepsilon_a(1) = a$. We denote also by ε_a the extension $\bar{\varepsilon}_a$. Without lost of generality, we may identify the element $a \in X$ and the gradual element $\varepsilon_a \in \mathcal{X}$ and denote them simply by a .

In this way, a gradual element is nothing more than a collection of elements of X , each one with a degree of membership; thus, if ε is a gradual element, then $\varepsilon(\alpha)$ is an element of X with membership degree α . Since $\varepsilon(1)$ is always defined, we have it as an element of X with the highest membership degree; since $\varepsilon(0)$ is not defined, then there is no any element with zero membership degree. \square

2.2. Relations between Gradual Elements. For any $\alpha \in [0, 1]$, in the set of all partial gradual elements, we define a relation R_α as, for partial gradual elements ε_1 and ε_2 of X , we say $\varepsilon_1 R_\alpha \varepsilon_2$ if

$$\varepsilon_{1|[\alpha,1] \cap \text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2)} = \varepsilon_{2|[\alpha,1] \cap \text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2)}. \quad (3)$$

Observe that if $\alpha \in (0, 1]$ and $\varepsilon_1, \varepsilon_2$ are total gradual elements, then $\varepsilon_1 R_\alpha \varepsilon_2$, whenever $\varepsilon_{1|[\alpha,1]} = \varepsilon_{2|[\alpha,1]}$.

Lemma 2. For any $\alpha, \beta \in [0, 1]$, we have

- (1) If $\alpha, \beta \in [0, 1]$ satisfies $\alpha \leq \beta$, then $R_\alpha \subseteq R_\beta$
- (2) The relation R_α is an equivalence relation in the set of all total gradual elements

The equivalence relation R_α indicates us when two gradual elements are equal at a certain level. For instance,

- (1) If $\alpha = 1$, then we only have an equivalence class for each element of X
- (2) If $\alpha = 0$, then two gradual elements belong to the same equivalence class if, and only if, they coincide in their definition domains

It is necessary to remark that these equivalence relations R_α are not compatible with the extension process. Indeed, if $\varepsilon_1 R_\alpha \varepsilon_2$, not necessarily $\bar{\varepsilon}_1 R_\alpha \bar{\varepsilon}_2$ as the following example shows.

Example 1. Let $X = \{a, b\}$, and define

$$\varepsilon_1(\delta) = \begin{cases} b, & \text{if } \delta = 1, \\ a, & \text{if } \frac{1}{2} \leq \delta < 1, \end{cases} \quad (4)$$

$$\varepsilon_2(\delta) = \begin{cases} b, & \text{if } \delta = 1, \\ a, & \text{if } 0 < \delta \leq \frac{1}{2}. \end{cases}$$

Then, $\varepsilon_1 R_\alpha \varepsilon_2$, for any $\alpha \in (0, 1]$, but $\bar{\varepsilon}_1 R_\alpha \bar{\varepsilon}_2$ if and only if $\alpha = 1$.

Let $f: X \rightarrow Y$ be a map between two sets.

- (1) For any total gradual (resp., partial gradual) element ε of X , we have a total gradual (resp., partial gradual) element of Y defined by the composition $f\varepsilon: \text{dom}(\varepsilon) \rightarrow X \rightarrow Y$; we call $f\varepsilon$ the image of ε by f .
- (2) For any gradual element ε' of Y , a gradual element ε of X is an inverse image of ε' if $\varepsilon' = f\varepsilon$

2.3. Binary Operations and Gradual Elements. There is another method to relate gradual elements of a set X ; this is the

case in which there exists a binary operation in X .

Let G be a set together a binary operation, say $*$, and $\varepsilon_1, \varepsilon_2$ gradual elements of G , we define a new gradual element $\varepsilon_1 * \varepsilon_2$ as

$$(\varepsilon_1 * \varepsilon_2)(\alpha) = \varepsilon_1(\alpha) * \varepsilon_2(\alpha), \quad \text{for any } \alpha \in \text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2). \quad (5)$$

In the case of partial gradual elements $\varepsilon_1, \varepsilon_2$, we have that $\text{dom}(\varepsilon_1 * \varepsilon_2) = \text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2)$.

This operation non necessarily is compatible with the extension construction.

The following example shows that if we start from two partial gradual elements ε_1 and ε_2 , then not necessarily we have the equality: $\overline{\varepsilon_1 * \varepsilon_2} = \bar{\varepsilon}_1 * \bar{\varepsilon}_2$, i.e., then extension map is not necessarily a homomorphism with respect to the binary operation $*$.

Example 2. Let $\varepsilon_1, \varepsilon_2$ be partial gradual elements defined on \mathbb{Z} , defined as:

$$\varepsilon_1(\alpha) = \begin{cases} 2, & \text{if } \frac{1}{2} \leq \alpha \leq 1, \\ 1, & \text{if } \frac{1}{10} \leq \alpha \leq \frac{1}{3}, \end{cases} \quad (6)$$

$$\varepsilon_2(\alpha) = 2, \quad \text{if } \frac{2}{3} \leq \alpha \leq 1.$$

In this case, we have

$$(\varepsilon_1 + \varepsilon_2)(\alpha) = 4, \quad \text{if } \frac{2}{3} \leq \alpha \leq 1. \quad (7)$$

and the extended gradual elements are


$$\bar{\varepsilon}_1(\alpha) = \begin{cases} 2, & \text{if } \frac{1}{3} < \alpha \leq 1, \\ 1, & \text{if } \alpha \leq \frac{1}{3}, \end{cases} \quad (8)$$

$$\bar{\varepsilon}_2(\alpha) = 2, \quad \text{if } \alpha \leq 1,$$

$$\overline{(\varepsilon_1 + \varepsilon_2)}(\alpha) = 4, \quad \text{if } \alpha \leq 1.$$

On the other hand, we have

$$(\bar{\varepsilon}_1 + \bar{\varepsilon}_2)(\alpha) = \begin{cases} 4, & \text{if } \frac{1}{3} < \alpha \leq 1, \\ 3, & \text{if } \alpha \leq \frac{1}{3}. \end{cases} \quad (9)$$



On the other hand, this operation is compatible with the equivalence relations R_α .

Lemma 3. Let G be a set together a binary operation $*$, for any $\alpha \in [0, 1]$ the relations R_α in the set of all total gradual elements (resp., in the set of all partial gradual elements) are compatible with the binary operation, i.e., for gradual elements $\varepsilon, \varepsilon_1, \varepsilon_2$ of G , if $\varepsilon_1 R_\alpha \varepsilon_2$, then $(\varepsilon * \varepsilon_1) R_\alpha (\varepsilon * \varepsilon_2)$ and $(\varepsilon_1 * \varepsilon) R_\alpha (\varepsilon_2 * \varepsilon)$.

Proof. Let $\varepsilon_1, \varepsilon_2, \varepsilon$ be gradual elements such that $\varepsilon_1 R_\alpha \varepsilon_2$, for any $\beta \in [\alpha, 1] \cap \text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2) \cap \text{dom}(\varepsilon)$, we have

$$\begin{aligned} (\varepsilon_1 * \varepsilon)(\beta) &= \varepsilon_1(\beta) * \varepsilon(\beta) \\ &= \varepsilon_2(\beta) * \varepsilon(\beta) \\ &= (\varepsilon_2 * \varepsilon)(\beta). \end{aligned} \tag{10}$$

In some cases, in which G has a richer structure, this structure could be inherited by the sets of gradual elements. Let us show an example. \square

$$\varepsilon_1 R_\alpha \varepsilon_2, \text{ if there is an inf - compact subset } C \subseteq (0, 1] \text{ containing } 1, \text{ such that } \varepsilon_{1|C} = \varepsilon_{2|C}. \tag{12}$$

With the relation R , the problem is that we may have $\text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2) = \{1\}$, and this trivialize this relation.

Hence, to obtain a well-defined structure on partial gradual elements, we may consider only special types of partial gradual elements, for instance, the subset of $\underline{\mathcal{G}}$ constituted by those partial gradual elements who have the same (inf-compact) domain containing 1.

Thus, we can extend Lemma 4 to consider gradual elements defined on an inf-compact subset containing 1.

Proposition 1. Let G be a group, and let $C \subseteq (0, 1]$ be an inf-compact subset containing 1. If $\underline{\mathcal{G}}$ be the set of all partial gradual elements whose domain is C , the following statements hold:

- (1) For any $\alpha \in C$, the relation R_α is an equivalent relation in $\underline{\mathcal{G}}$

Lemma 4. Let G be a group, with binary operation $*$ and neutral element e , the following statements hold:

- (1) The set \mathcal{G} of all total gradual elements is a group with neutral element e , i.e., the total gradual element ε_e
- (2) For any $\alpha \in [0, 1]$, we have that \mathcal{G}/R_α is a group

Proof. For any $\varepsilon, \varepsilon_1, \varepsilon_2 \in \mathcal{G}$, we define, for any $\alpha \in (0, 1]$:

$$(\varepsilon_1 * \varepsilon_2)(\alpha) = \varepsilon_1(\alpha) * \varepsilon_2(\alpha), \quad \text{and} \quad (\varepsilon^{-1})(\alpha) = \varepsilon(\alpha)^{-1}. \tag{11}$$

- (1) In \mathcal{G} , the operation is associative and e is the neutral element. For any $\varepsilon \in \mathcal{G}$, we have ε^{-1} the inverse of ε . Therefore, \mathcal{G} is a group, which is abelian whenever G is.
- (2) It is a direct consequence of being R_α a compatible equivalence relation. \square

Remark 1 (The particular case of partial gradual elements). Let G be a group, in the set $\underline{\mathcal{G}}$ of all partial gradual elements, we have an associative binary operation, $(\varepsilon_1 * \varepsilon_2)(\alpha) = \varepsilon_1(\alpha) * \varepsilon_2(\alpha)$ for any $\alpha \in \text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2)$, but we have “many” possible neutral elements. Thus, to get a useful structure, we must define before an equivalence relation to put together all of them. For instance, given two partial gradual elements $\varepsilon_1, \varepsilon_2$, since $\varepsilon_1 * \varepsilon_2$ is defined on $\text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2)$, there are three possible neutral elements: $e_{|\text{dom}(\varepsilon_1)}, e_{|\text{dom}(\varepsilon_2)}$ and finally $e_{|\text{dom}(\varepsilon_1) \cap \text{dom}(\varepsilon_2)}$, which are different two to two.

We can try to fix this problem in defining an equivalence relation R in $\underline{\mathcal{G}}$ generated by

- (2) In $\underline{\mathcal{G}}$, we have an associative operation
- (3) The extending map from $\underline{\mathcal{G}}$ to \mathcal{G} is a group monomorphism
- (4) For any $\alpha \in C$, the equivalence relation R_α in $\underline{\mathcal{G}}$ is compatible
- (5) The groups $\underline{\mathcal{G}}$ and $\underline{\mathcal{G}}/R_\alpha$ are abelian whenever G is

Proof

- (1) It is reflexive and symmetric, and obviously it is transitive as the domain is the whole set C .
- (2) It is evident as the product is defined componentwise.
- (3) Let $\varepsilon_1, \varepsilon_2 \in \underline{\mathcal{G}}$, and $\xi = \text{Min}([\alpha, 1] \cap C)$, then

$$\begin{aligned}
\overline{(\varepsilon_1 * \varepsilon_2)}(\alpha) &= (\varepsilon * \varepsilon_2)(\xi) \\
&= \varepsilon_1(\xi) * \varepsilon_2(\xi) \\
&= \bar{\varepsilon}_1(\alpha) * \bar{\varepsilon}_2(\alpha).
\end{aligned} \tag{13}$$

- (4) It is similar to the proof on Lemma 3.
(5) It is evident as the product is defined componentwise.

It is clear that it is better to consider total gradual elements instead of partial gradual elements and therefore work in \mathcal{G} .

If the group G has e as neutral element and for any α in $[0, 1]$ we consider the equivalence relation R_α , we may rewrite Lemma 3 obtaining a filtration of subgroups of \mathcal{G} . \square

Proposition 2. *Let G be a group with neutral element e , if for any $\alpha \in [0, 1]$, we define*

$$\mathcal{G}_\alpha = \{\varepsilon \in \mathcal{G} \mid \varepsilon R_\alpha e\}. \tag{14}$$

Then, we have

- (1) For each $\alpha \in (0, 1]$, the subgroup $\mathcal{G}_\alpha \subseteq \mathcal{G}$ is a normal.
- (2) There is a filtration $\{\mathcal{G}_\alpha \mid \alpha \in [0, 1]\}$ where $\mathcal{G}_\alpha \subseteq \mathcal{G}_\beta$ if $\alpha \leq \beta$.
- (3) We have inclusions: $\mathcal{G}_0 \subseteq \mathcal{G}_\alpha \subseteq \mathcal{G}_1 \subseteq \mathcal{G}$, and surjective group homomorphisms:

$$\frac{\mathcal{G}}{\mathcal{G}_0} \longrightarrow \frac{\mathcal{G}}{\mathcal{G}_\alpha} \longrightarrow \frac{\mathcal{G}}{\mathcal{G}_1} \cong G. \tag{15}$$

Observe that in all these examples, it seems that the way to define an operation on gradual elements is to define it componentwise.

If the base set X has a more richer structure, for instance, if it is a ring R , then the corresponding sets \mathcal{R} and $\underline{\mathcal{R}}$ are rings, but there are in these rings many elements which are zero-divisor. So, in this case, the use of gradual elements is not a good option. For that, in this and forthcoming papers, we shall develop a different approach to study algebraic structures. Before doing that, let us study the simplest notion of gradual subset, and after doing this we shall return to consider a set endowed with one or several binary operations, for instance, a group.

3. Gradual Subsets

Once we have established the notion of gradual element of a set X , we shall apply it to define new objects. If we consider a set X and the power set $\mathcal{P}(X)$, we can study gradual elements of $\mathcal{P}(X)$, thereby the concept of gradual subset appears.

3.1. Definition of Gradual Subsets

Definition 2. Let X be a set, and let $\mathcal{P}(X)$ be the power set of X , i.e., $\mathcal{P}(X) = \{S \mid S \text{ is a subset of } X\}$. We define a gradual subset of X as a gradual element of $\mathcal{P}(X)$. We represent by $\sigma: (0, 1] \rightarrow \mathcal{P}(X)$ a gradual subset of X .

Throughout this section, we follow the same assumptions used for gradual elements in the previous section. In this way, we have defined partial gradual subsets and total gradual subsets.

In some sense, gradual subsets are a generalization of gradual elements. Thus, for any gradual element ε and any gradual subset σ , we say ε belongs to σ if for any $\alpha \in \text{dom}(\varepsilon) \cap \text{dom}(\sigma) \subseteq [0, 1]$, we have $\varepsilon(\alpha) \in \sigma(\alpha)$ and write $\varepsilon \in \sigma$. In the same way, given two gradual subsets σ_1, σ_2 , we say that σ_1 is a subset of σ_2 if $\sigma_1(\alpha) \subseteq \sigma_2(\alpha)$ for any $\alpha \in \text{dom}(\sigma_1) \cap \text{dom}(\sigma_2)$, we write $\sigma_1 \subseteq \sigma_2$.

In general, for any gradual subset σ and elements $\alpha, \beta \in \text{dom}(\sigma)$ such that $\alpha \leq \beta$, we have no information about the relationship of $\sigma(\alpha)$ and $\sigma(\beta)$. In some cases, as in the classical one of α -levels in fuzzy set theory, there is an evident relationship, as we shall see later. To work with them, first we introduce the following definitions that reflect the order existing in $(0, 1]$.

Let σ be a gradual subset of X , we say σ is

- (1) Increasing if for any $\alpha, \beta \in \text{dom}(\sigma)$ such that $\alpha \leq \beta$, we have $\sigma(\alpha) \subseteq \sigma(\beta)$. For any increasing gradual subset σ de X , if $\zeta = \text{Min}(\text{dom}(\sigma))$, then $\sigma(\zeta) \subseteq \sigma(\alpha) \subseteq \sigma(1)$ for any $\alpha \in \text{dom}(\sigma)$.
- (2) Decreasing if for any $\alpha, \beta \in \text{dom}(\sigma)$ such that $\alpha \leq \beta$, we have $\sigma(\alpha) \supseteq \sigma(\beta)$. For any decreasing gradual subset σ de X , we have $\sigma(1) \subseteq \sigma(\alpha)$ for any $\alpha \in \text{dom}(\sigma)$.

Let us show some examples of decreasing gradual subsets.

Example 3. Let μ be a fuzzy subset of X , i.e., a map $\mu: X \rightarrow [0, 1]$ that we assume it is not constant equal to 0. For any $\alpha \in (0, 1]$, we define

- (1) α -level of μ as $\mu_\alpha = \{x \in X \mid \mu(x) \geq \alpha\}$. In this case, we have a decreasing gradual subset $\sigma(\mu)$, defined $\sigma(\mu)(\alpha) = \mu_\alpha$ for any $\alpha \in (0, 1]$.
- (2) Strict α -level (or strong α -level) of μ as $\bar{\mu}_\alpha = \{x \in X \mid \mu(x) > \alpha\}$. In this case, we have a decreasing gradual subset $\bar{\sigma}(\mu)$, defined as

$$\bar{\sigma}(\mu)(\alpha) = \begin{cases} \bar{\mu}_\alpha, & \text{for any } \alpha \in (0, 1), \text{ and,} \\ \mu_1, & \text{if } \alpha = 1. \end{cases} \tag{16}$$

- (3) Let $\bar{\mu}$ the fuzzy subset defined by $\bar{\mu}(x) = 1 - \mu(x)$, for any $x \in X$; the α -levels of $\bar{\mu}$ define a decreasing gradual subset $\sigma(\bar{\mu})(\alpha) = \{x \in X \mid \mu(x) \leq 1 - \alpha\}$.

- (4) Inverse α -level of μ as $\mu^\alpha = \{x \in X | \mu(x) \leq \alpha\}$. In this case, we have an increasing gradual subset $\tau(\mu)$ defined $\tau(\mu)(\alpha) = \mu^\alpha$ for any $\alpha \in (0, 1]$.

3.2. *Operators on Gradual Subsets.* The following are examples of constructions that can be carried out for any gradual subset, which will be useful in their study.

Let σ be a gradual subset of X , associated to σ , we define two new gradual subsets:

- (1) The accumulation σ^c of σ is

$$\sigma^c(\alpha) = \cup \{ \sigma(\beta) | \alpha \leq \beta \in \text{dom}(\sigma) \}, \quad \text{for any } \alpha \in \text{dom}(\sigma). \quad (17)$$

It is clear that for any gradual subset σ , the accumulation σ^c is a decreasing gradual subset, and a gradual subset σ is decreasing if and only if, $\sigma = \sigma^c$.

For any gradual subset σ , we have $\sigma \subseteq \sigma^c = \sigma^{cc}$.

- (2) The strict accumulation σ^d of σ is

$$\sigma^d(\alpha) = \begin{cases} \sigma(1), & \text{if } \alpha = 1, \\ \cup \{ \sigma(\beta) | \alpha < \beta \in \text{dom}(\sigma) \}, & \text{if } \alpha \in \text{dom}(\sigma) \setminus \{1\}. \end{cases} \quad (18)$$

It is clear that for any gradual subset σ , the strict accumulation σ^d is a decreasing gradual subset and $\sigma^d \subseteq \sigma^c$. In general, $\sigma \not\subseteq \sigma^d$.

Thus, we have an operator, c , on gradual subsets: $\sigma \mapsto \sigma^c$. The behaviour of c is reflected in the following lemma.

Lemma 5. *Let X be a set, for any gradual subsets $\sigma_1, \sigma_2, \sigma$ of X , the following statements hold:*

- (1) $\sigma \subseteq \sigma^c$
- (2) $\sigma^c = \sigma^{cc}$
- (3) If $\sigma_1 \subseteq \sigma_2$, then $\sigma_1^c \subseteq \sigma_2^c$
- (4) σ^c is the smallest decreasing gradual subset containing σ

Proof. (1), (2), and (3) are easy

- (5) Let τ be a decreasing gradual subset such that $\sigma \subseteq \tau$, then $\sigma^c \subseteq \tau^c = \tau$.

This means that the operator c is a closure operator in the set \mathcal{X} of all gradual subsets of X .

Remember that a closure operator in a poset (partial ordered set) P is a map $c: P \rightarrow P$ satisfying

- (1) $p \leq c(p)$ for any $p \in P$
- (2) For any $p_1, p_2 \in P$ such that $p_1 \leq p_2$ we have $c(p_1) \leq c(p_2)$
- (3) $c(p) = cc(p)$ for any $p \in P$

The elements $p \in P$ such that $c(p) = p$ are named the c -closed elements. Thus, the gradual subsets which are closed for the operator c are the decreasing gradual subsets. Let us denote by $\mathcal{F}(X)$ the set of all decreasing gradual subsets of X .

In the same way, we may consider the operator d , defined as $\sigma \mapsto \sigma^d$; its behaviour is reflected in the following lemma.

Lemma 6. *Let X be a set, for any gradual subsets $\sigma_1, \sigma_2, \sigma$ of X , the following statements hold:*

- (1) $\sigma^d \subseteq \sigma^c$
- (2) If $\sigma_1 \subseteq \sigma_2$, then $\sigma_1^d \subseteq \sigma_2^d$
- (3) $\sigma^d = \sigma^{d^d} = \sigma^{c^d} = \sigma^{dc}$

Proof. (1) and (2) are easy. (3) Indeed, for any $\alpha \in (0, 1]$, we have

$$\begin{aligned} \sigma^{d^d}(\alpha) &= \cup \{ \sigma^d(\beta) | \beta > \alpha \} \\ &= \cup \{ \cup \{ \sigma(\gamma) | \gamma > \beta \} | \beta > \alpha \} \\ &= \cup \{ \sigma(\beta) | \beta > \alpha \} \\ &= \sigma^d(\alpha). \end{aligned} \quad (19)$$

In the same way, for any $\alpha \in (0, 1]$, we have

$$\begin{aligned} \sigma^{c^d}(\alpha) &= \cup \{ \sigma^c(\beta) | \beta > \alpha \} \\ &= \cup \{ \cup \{ \sigma(\gamma) | \gamma \leq \beta \} | \beta > \alpha \}, \\ &= \cup \{ \sigma(\beta) | \beta > \alpha \}, \\ &= \sigma^d(\alpha). \end{aligned} \quad (20)$$

A gradual subset σ is an strict decreasing gradual subset if $\sigma = \sigma^d$. We have \square

Lemma 7. *For any gradual subset σ , the following statements hold:*

- (1) σ^d is the smallest strict decreasing gradual subset contained in σ^c
- (2) σ is a decreasing gradual subset non strict decreasing if, and only if, $\sigma^d \subsetneq \sigma^c$

Proof. Let τ be a strict decreasing gradual subset such that $\tau \subseteq \sigma^c$, then $\tau = \tau^d \subseteq \sigma^{c^d} = \sigma^d$.

This means that the operator d is an interior operator in the set of all decreasing gradual subsets of X .

Remember that an interior operator in a poset P is a map $d: P \rightarrow P$ satisfying

- (1) $d(p) \leq p$ for any $p \in P$
- (2) For any $p_1, p_2 \in P$ such that $p_1 \leq p_2$, we have $d(p_1) \leq d(p_2)$
- (3) $d(p) = d^d(p)$ for any $p \in P$

The elements $p \in P$ such that $d(p) = p$ is named the d -open elements. Thus, the decreasing gradual subsets open for the operator d are the strict decreasing gradual subsets. Let us denote by $\mathcal{F}^d(X)$ the set of all d -open (strict) decreasing gradual subsets. \square

Remark 2. Inspired in these constructions, we consider a new construction of a gradual subset from a partial gradual

subset that allows us to avoid the initial restriction of inf-compact in the domain of definition of partial gradual elements.

Let $\sigma: (0, 1] \rightarrow \mathcal{P}(X)$ be a partial map defined at 1, i.e., $1 \in \text{dom}(\sigma) \subseteq (0, 1]$, and such that $\text{dom}(\sigma)$ is not necessarily inf-compact, we may extend σ to all $(0, 1]$ simply defining $\sigma(\beta) = \emptyset$ if $\beta \notin \text{dom}(\sigma)$. The decreasing gradual subset associated to σ is $\sigma^c: (0, 1] \rightarrow \mathcal{P}(X)$ defined as

$$\sigma^c(\alpha) = \cup \{ \sigma(\beta) \mid \beta \geq \alpha, \beta \in \text{dom}(\sigma) \}, \text{ for any } \alpha \in (0, 1]. \quad (21)$$

The use of decreasing gradual subsets is due to the fact that gradual subsets are wild structures that one can not be managed, in which there is no relationship between its components. On the other hand, when studying subsets of a given set, it seems natural to impose some inclusion relationships and that these inclusions should be parameterized by the order relation in $(0, 1]$.

Remark 3. Observe that we may extend any gradual subset σ on X to $\bar{\sigma}$ on the whole interval $[0, 1]$, in defining $\bar{\sigma}(\alpha) = \begin{cases} \cup \{ \sigma(\beta) \mid \alpha \leq \beta \in (0, 1] \}, & \text{if } \alpha \in (0, 1], \\ \cup \{ \sigma(\beta) \mid \beta \in (0, 1] \}, & \text{if } \alpha = 0, \end{cases}$ for any $\alpha \in [0, 1]$. In consequence, we may consider also decreasing gradual subsets as maps from $[0, 1]$ to X .

3.3. The Algebra of Gradual Subsets. There is a natural relationship between gradual elements and gradual subsets of a given set X . Thus, for any partial gradual element ε , we may define a unitary partial gradual subset $\sigma(\varepsilon)$ as $\sigma(\varepsilon)(\alpha) = \{ \varepsilon(\alpha) \}$, for any $\alpha \in \text{dom}(\varepsilon)$. As we point out before, we have $\varepsilon \in \sigma$ if, and only if, $\sigma(\varepsilon) \subseteq \sigma$, for any gradual subset σ .

In the set $\mathcal{P}(X)$, there are two operations: the intersection and the union; thus, we can translate these two operations to gradual subsets as did in the first section. Following this line, we define, for any gradual subsets, σ_1 and σ_2 :

- (1) The intersection, $\sigma_1 \cap \sigma_2$, as $(\sigma_1 \cap \sigma_2)(\alpha) = \sigma_1(\alpha) \cap \sigma_2(\alpha)$, for any $\alpha \in \text{dom}(\sigma_1) \cap \text{dom}(\sigma_2)$
- (2) The union, $\sigma_1 \cup \sigma_2$, as $(\sigma_1 \cup \sigma_2)(\alpha) = \sigma_1(\alpha) \cup \sigma_2(\alpha)$, for any $\alpha \in \text{dom}(\sigma_1) \cap \text{dom}(\sigma_2)$

In this way, we may consider the algebra of gradual subsets of a given set X with respect to intersection and union.

The definition of intersection and union can also be extended to arbitrary families of gradual subsets. Let $\{ \sigma_i \mid i \in I \}$ be a family of gradual subsets.

- (1) The intersection $\cap_i \sigma_i$ is defined as $(\cap_i \sigma_i)(\alpha) = \cap_i \sigma_i(\alpha)$, for any $\alpha \in (0, 1]$
- (2) The union $\cup_i \sigma_i$ defined as $(\cup_i \sigma_i)(\alpha) = \cup_i \sigma_i(\alpha)$, for any $\alpha \in (0, 1]$

Let $\{ \mu_i \mid i \in I \}$ be a family of fuzzy subsets of a set X , the union, $\vee_i \mu_i$, and the intersection, $\wedge_i \mu_i$, are the fuzzy subsets defined by

$$\begin{aligned} (\vee_i \mu_i)(a) &= \vee_i \mu_i(a) = \text{Sup} \{ \mu_i(a) \mid i \in I \}, \quad \text{for any } a \in X, \\ (\wedge_i \mu_i)(a) &= \wedge_i \mu_i(a) = \text{Inf} \{ \mu_i(a) \mid i \in I \}, \quad \text{for any } a \in X. \end{aligned} \quad (22)$$

Example 4. Let $X = \{a, b\}$ be a set, for any $n \in \mathbb{N} \setminus \{0, 1\}$, we define $\mu_n: \{a, b\} \rightarrow [0, 1]$ by $\mu_n(a) = 1$ and $\mu_n(b) = (1/2) - (1/2^n)$. We have

- (1) $(\vee_n \mu_n)(a) = 1$ and $(\vee_n \mu_n)(b) = (1/2)$
- (2) $\sigma(\vee_n \mu_n)(\delta) = \begin{cases} X, & \text{if } \delta > (1/2), \\ \{a\}, & \text{if } \delta \leq (1/2), \end{cases}$
- (3) $(\cup_n \sigma(\mu_n))(\delta) = \begin{cases} X, & \text{if } \delta \geq (1/2), \\ \{a\}, & \text{if } \delta < (1/2), \end{cases}$

This shows that the inclusion $\sigma(\vee_n \mu_n) \supseteq \cup_n \sigma(\mu_n)$ is proper.

In the same line, we have a similar situation for $\bar{\sigma}$ and the intersection.

Example 5. Let $X = \{a, b\}$ be a set, for any $n \in \mathbb{N} \setminus \{0, 1\}$, we define $\mu_n: \{a, b\} \rightarrow [0, 1]$ by $\mu_n(a) = 1$ and $\mu_n(b) = (1/2) + (1/2^n)$. We have

- (1) $(\wedge_n \mu_n)(a) = 1$ and $(\wedge_n \mu_n)(b) = (1/2)$
- (2) $\bar{\sigma}(\wedge_n \mu_n)(\delta) = \begin{cases} X, & \text{if } \delta \geq (1/2), \\ \{a\}, & \text{if } \delta < (1/2), \end{cases}$
- (3) $(\cap_n \bar{\sigma}(\mu_n))(\delta) = \begin{cases} X, & \text{if } \delta > (1/2), \\ \{a\}, & \text{if } \delta \leq (1/2), \end{cases}$

This shows that the inclusion $\bar{\sigma}(\wedge_n \mu_n) \subseteq \cap_n \bar{\sigma}(\mu_n)$ is proper.

In the set $\mathcal{F}^d(X)$ of all strict decreasing gradual subsets of X , we define two new operations: intersection: $\wedge_i \sigma_i = (\cap_i \sigma_i)^d$, and maintain the old union: $\vee_i \sigma_i = \cap_i \sigma_i$, for every family $\{ \sigma_i \mid i \in I \}$ of strict decreasing gradual subsets of X . With these definitions, we have the following.

Proposition 3. *The union and intersection of strict decreasing gradual subsets are compatible with the union and intersection of fuzzy subsets via the gradual subset $\bar{\sigma}(\mu)$, i.e., for any family of fuzzy subsets $\{ \mu_i \mid i \in I \}$, we have*

- (1) $\vee_i \bar{\sigma}(\mu_i) = \bar{\sigma}(\vee_i \mu_i)$
- (2) $\wedge_i \bar{\sigma}(\mu_i) = \bar{\sigma}(\wedge_i \mu_i)$

Proof. For any $\alpha \in (0, 1]$, we have

$$\begin{aligned} \bar{\sigma}(\vee_i \mu_i)(\alpha) &= \{a \in X \mid (\vee_i \mu_i)(a) > \alpha\} \\ &= \{a \in X \mid \vee_i \mu_i(a) > \alpha\}, \\ &= \cup \{a \in X \mid \text{there exists } i \text{ such that } \mu_i(a) > \alpha\} \\ &= (\cup_i \bar{\sigma}(\mu_i))(\alpha) \\ &= (\vee_i \bar{\sigma}(\mu_i))(\alpha). \end{aligned} \tag{23}$$

In the same way, we can prove the case of $\bar{\sigma}(\wedge_i \mu_i)(\alpha)$.

From this point of view, strict α -levels should be a suitable tool for studying the algebra of fuzzy subsets via decreasing gradual subsets. \square

3.4. Maps. In order to relate two gradual subsets, a standard method consists in defining a map from one to the other. In this context, first we consider a map between the underlying sets containing each gradual subset; the following result show how to associate gradual subsets to gradual subsets via a map.

Lemma 8 (Direct image). *Let $f: X \rightarrow Y$ be a map and denote by f the induced map from $\mathcal{P}(X)$ to $\mathcal{P}(Y)$, the following statements hold:*

- (1) For every gradual element ε of X , we have that $f\varepsilon: \text{dom}(\varepsilon) \rightarrow Y$ is a gradual element of Y .
- (2) Let σ be a gradual subset of X , then $f\sigma: \text{dom}(\sigma) \rightarrow \mathcal{P}(Y)$ is a gradual subset of Y . And we have a map $f_*: \mathcal{X} \rightarrow \mathcal{Y}$ defined $f_*(\sigma) = f\sigma$ for any $\sigma \in \mathcal{X}$.
- (3) Let σ be a gradual subset of X , then $f_*(\sigma^c) = (f_*(\sigma))^c$.

In addition, if $\varepsilon \in \sigma$, then $f\varepsilon \in f\sigma$.

Lemma 9 (Inverse image). *Let $f: X \rightarrow Y$ be a map and denote by $f^{-1}: \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$ the induced map. For any gradual subset τ of Y , we have that $f^*\tau: \text{dom}(\tau) \rightarrow \mathcal{P}(X)$, defined as $f^*\tau(\alpha) = f^{-1}(\tau(\alpha))$, for any $\alpha \in \text{dom}(\tau)$, is a partial gradual subset of X . Thus, we have a map $f^*: \mathcal{Y} \rightarrow \mathcal{X}$, defined $f^*(\tau) = f^*\tau$, for any $\tau \in \mathcal{Y}$.*

In particular, for any gradual subset τ of Y , we have $f^*(\tau^c) = (f^*(\tau))^c$.

Since every element of X and every element of Y are gradual elements and the same for gradual subsets, the notions of injective map and surjective map applied either to gradual elements or gradual subsets are equivalent. In the case of gradual subsets, we have

Lemma 10. *Let $f: X \rightarrow Y$ be a map, then*

- (1) f is injective if, and only if, $f^* \circ f_* = id_{\mathcal{X}}$
- (2) f is surjective if, and only if, $f_* \circ f^* = id_{\mathcal{Y}}$

Our aim will be to establish maps between gradual sets instead of between gradual subsets, i.e., leave out the ground

set X and use only the subsets $\{\sigma(\alpha) \mid \alpha \in (0, 1]\}$. However, we postpone it until the point in which we change the paradigm introducing these gradual sets.

Remark 4. A gradual subset σ of a set X is just a family $\{\sigma(\alpha) \mid \alpha \in (0, 1]\}$ of subsets, indexed in $(0, 1]$. There are particular types of gradual subsets, as decreasing gradual subsets, in which, for any $\alpha, \beta \in (0, 1]$, $\alpha \leq \beta$, there exists a map $j_{\alpha, \beta}: \sigma(\beta) \rightarrow \sigma(\alpha)$: the inclusion, satisfying $j_{\alpha, \beta} j_{\beta, \gamma} = j_{\alpha, \gamma}$ whenever $\alpha \leq \beta \leq \gamma$. In some sense, decreasing gradual subsets are gradual subsets enriched with a family of maps $\{j_{\alpha, \beta} \mid \alpha, \beta \in (0, 1], \alpha \leq \beta\}$ satisfying the above conditions and compatible with the inclusions in X . Thus, we may work with these enriched gradual subsets of X .


An enriched gradual subset of X is a gradual subset σ together with a family of maps $\{f_{\alpha, \beta} \mid \alpha, \beta \in (0, 1], \alpha \leq \beta\}$ satisfying

- (1) $f_{\alpha, \beta}: \sigma(\beta) \rightarrow \sigma(\alpha)$
- (2) $f_{\alpha, \beta} f_{\beta, \gamma} = f_{\alpha, \gamma}$ whenever $\alpha \leq \beta \leq \gamma$
- (3) if $j_\alpha: \sigma(\alpha) \rightarrow X$ is the inclusion, for any $\alpha \in (0, 1]$, then $j_\alpha f_{\alpha, \beta} = j_\beta$, whenever $\alpha \leq \beta$


Observe that, as a consequence of (3), each $f_{\alpha, \beta}$ is an injective map. In particular, enriched gradual subsets are just the decreasing gradual subsets. See also Remark 7.

3.5. Gradual Quotient Sets. The same technique we used to introduce gradual subsets can be applied to define quotient gradual sets of a given set X .

Remember that if X is a set, a subset $S \subseteq X$ is an equivalence class in the class of all injective maps $\{(i, Y) \mid i: Y \rightarrow X \text{ injective}\}$, whenever we consider the equivalence relation: $(i_1, Y_1) \sim (i_2, Y_2)$ if there exists a bijective map $b: Y_1 \rightarrow Y_2$ such that $i_1 = i_2 b$.



Dually, a quotient set of X is an equivalence class in the class of all surjective maps $\{(Z, p) \mid p: X \rightarrow Z \text{ surjective}\}$ when we consider the equivalence relation: $(Z_1, p_1) \sim (Z_2, p_2)$ if there exists a bijective map $b: Z_1 \rightarrow Z_2$ such that $p_2 = b p_1$.



The set of all subsets of X is represented by $\mathcal{P}(X)$, and there exists a bijective correspondence between $\mathcal{P}(X)$ and 2^X . The set of all quotient set of X will be represented by $\mathcal{Q}(X)$, and for any element $Z \in \mathcal{Q}(X)$, we have

- (1) A surjective map $p: X \rightarrow Z$

- (2) An equivalence relation R_p in X defined as $xR_p y$ if $p(x) = p(y)$
- (3) A partition of X into the equivalence classes defined by a relation R .

Each equivalence relation R in X is a subset of $X \times X$ satisfying the properties reflexive, symmetric and transitive. Hence, $\mathcal{Q}(X)$ is in bijection with a subset of $\mathcal{P}(X \times X)$. If we call $\mathcal{Q}(X \times X)$ this subset, it is constituted by all the equivalence relations in X .

A gradual quotient set of X is a gradual element of $\mathcal{Q}(X)$ or equivalently of $\mathcal{Q}(X \times X)$. We represent by ρ a gradual quotient set of X .

3.6. Gradual Subsets and Fuzzy Subsets. As an example of application of the gradual subset theory, let us establish a correspondence between fuzzy subsets and enriched gradual subsets. As we had shown before, see Proposition 3; if we consider the strict decreasing gradual subset $\bar{\sigma}(\mu)$, the correspondence $\mu \mapsto \bar{\sigma}(\mu)$ is a homomorphism with respect to arbitrary union and intersection.

In addition, the gradual subsets $\sigma(\mu)$ and $\bar{\sigma}(\mu)$ are related in a strong way: $\bar{\sigma}(\mu) = \sigma(\mu)^d \subseteq \sigma(\mu)^c = \sigma(\mu)$, using the interior and closure operator. Also, these gradual subsets satisfy the following properties:

- (1) $\mu(x) = \text{Max}\{\alpha | x \in \sigma(\mu)(\alpha)\}$, for any $x \in X$
- (2) $\mu(x) = \text{Inf}\{\alpha | x \notin \bar{\sigma}(\mu)(\alpha)\}$, for any $x \in X$

We say

- (1) A decreasing gradual subset σ satisfies property (F) if there exists $\text{Max}\{\alpha | x \in \sigma(\alpha)\}$ for every $x \in \cup\{\alpha | \alpha \in (0, 1]\}$
- (2) A strict decreasing gradual subset σ satisfies property (inf-F) if $\text{Inf}\{\alpha | x \notin \sigma^d(\alpha)\} = \beta$ satisfies $x \in \sigma(\beta)$, for every $x \in \cup\{\alpha | \alpha \in (0, 1]\}$

As a consequence, we have the following result.

Lemma 11. *Let σ be a decreasing gradual subset, not strict decreasing gradual subset, the following statements are equivalent:*

- (i) σ satisfies property (F)
- (ii) σ^d satisfies property (inf-F)
- (iii) $\cup_{\alpha \in (0,1]} \sigma(\alpha) = \cup_{\alpha \in (0,1]} (\sigma^c(\alpha) \setminus \sigma^d(\alpha))$ (the disjoint union)

Proof. By the hypothesis, we have $\sigma^d \subseteq \sigma^c = \sigma$.

- (i) \Rightarrow (ii) Let $\beta = \text{Max}\{\alpha | x \in \sigma(\alpha)\}$ and $\gamma = \text{Inf}\{\alpha | x \notin \sigma^d(\alpha)\}$, then $x \notin \sigma(\gamma)$ for every $\gamma > \beta$. Since σ is decreasing, $\beta \leq \gamma$. If $\beta < \gamma$, for any ω such that $\beta < \omega < \gamma$, we have $x \notin \sigma(\omega)$, hence $\gamma \neq \text{Inf}\{\alpha | x \notin \sigma^d(\alpha)\}$, which is a contradiction.
- (ii) \Rightarrow (i) Let $\beta = \text{Sup}\{\alpha | x \in \sigma(\alpha)\}$ and $\gamma = \text{Inf}\{\alpha | x \notin \sigma^d(\alpha)\}$. Since σ is decreasing, $\beta \leq \gamma$. If $\beta < \gamma$, for any ω such that $\beta < \omega < \gamma$, we have $x \in \sigma(\omega)$ and $\beta \neq \text{Sup}\{\alpha | x \in \sigma(\alpha)\}$, which is a contradiction.

- (i) \Rightarrow (iii) One inclusion is obvious. Otherwise, if $x \in \cup_{\alpha \in (0,1]} \sigma(\alpha)$, let $\beta = \text{Max}\{\alpha | x \in \sigma(\alpha)\}$, then $x \notin \sigma(\gamma)$ for any $\gamma > \beta$; hence, $x \notin \sigma^d(\beta)$ and $x \in \cup_{\alpha \in (0,1]} (\sigma^c(\alpha) \setminus \sigma^d(\alpha))$.
- (iii) \Rightarrow (i) Let $x \in X$, if $x \notin \cup_{\alpha \in (0,1]} \sigma(\alpha)$, then either $x \in \sigma(1)$, and there exists $\text{Max}\{\alpha | x \in \sigma(\alpha)\} = 1$ or $x \notin \cup_{\alpha \in (0,1]} \sigma(\alpha)$ and $\text{Max}\{\alpha | x \in \sigma(\alpha)\} = 0$. Otherwise, if $x \in \cup_{\alpha \in (0,1]} \sigma(\alpha) = \cup_{\alpha \in (0,1]} (\sigma^c(\alpha) \setminus \sigma^d(\alpha))$, there exists α such that $x \in \sigma(\alpha) \setminus \sigma^d(\alpha)$, hence $\text{Max}\{\alpha | x \in \sigma(\alpha)\} = \alpha$. \square

Remark 5

- (1) As a consequence of this result, for any decreasing nonstrict decreasing gradual subset, satisfying property (F), we have $\text{Max}\{\alpha | x \in \sigma(\alpha)\} = \text{Inf}\{\alpha | x \notin \sigma^d(\alpha)\}$ for any $x \in X$
- (2) In the case of a strict decreasing gradual subset satisfying property (inf-F), we have that it also satisfies property (F); hence the following equalities hold: $\text{Max}\{\alpha | x \in \sigma(\alpha)\} = \text{Inf}\{\alpha | x \notin \sigma^d(\alpha)\}$ for any $x \in X$

Now, we are going to establish correspondences between fuzzy subsets and strict decreasing gradual subsets, which preserves union and intersection. The following result, for finite unions, is well known.

Theorem 1. *Let X be a set, the following statements hold:*

- (1) *The map $\nu: \mu \mapsto \sigma(\mu)$ associates to any fuzzy subset μ of X a decreasing gradual subset satisfying property (F), $\sigma(\mu)$ of X , defined $\sigma(\mu)(\alpha) = \{x | \mu(x) \geq \alpha\}$, and it preserves intersections and finite unions.*
- (2) *The map $\nu: \sigma \mapsto \mu(\sigma)$ associates to any decreasing gradual subset $\sigma = \sigma^c$, satisfying property (F); a fuzzy subset $\mu(\sigma)$ is defined as follows:*

$$\mu(\sigma)(x) = \text{Max}\{\alpha | x \in \sigma(\alpha)\}. \quad (24)$$

In addition, we have $\nu \circ \nu = \text{id}$ and $\nu \circ \nu = \text{id}$, and they preserve finite unions and intersections.

The behaviour with respect to infinite unions can be solved using only strict decreasing gradual subsets instead of decreasing gradual subsets.

Theorem 2. *Let X be a set, the following statements hold:*

- (1) *The map $\nu: \mu \mapsto \bar{\sigma}(\mu)$ associates to any fuzzy subset μ of X a strict decreasing gradual subset $\bar{\sigma}(\mu)$ of X , defined $\bar{\sigma}(\mu)(\alpha) = \{x | \mu(x) > \alpha\}$, and it preserves arbitrary unions and intersections.*
- (2) *The map $\nu: \sigma \mapsto \bar{\mu}(\sigma)$ associates to any strict decreasing gradual subset $\sigma = \sigma^d$, satisfying property (inf-F); a fuzzy subset $\bar{\mu}(\sigma)$ is defined as follows:*

$$\bar{\mu}(\sigma)(x) = \text{Inf}\{\alpha | x \notin \sigma(\alpha)\}. \quad (25)$$

In addition, we have $\nu \nu = \text{id}$ and $\nu \nu = \text{id}$.

Proof. We had already studied the map ν in Proposition 3.

The map ν is well defined due to Theorem 2. Now, we check that the compositions are the identity.

Let μ be a fuzzy subset of X , for any $x \in X$, we have

$$\begin{aligned} \nu \circ \nu(\mu)(x) &= \tilde{\mu}((\tilde{\sigma}(\mu))x) \\ &= \text{Inf}\{\alpha | x \notin \tilde{\sigma}(\mu)(\alpha)\} \\ &= \text{Inf}\{\alpha | \mu(x) \leq \alpha\} \\ &= \mu(x). \end{aligned} \tag{26}$$

On the other hand, let σ be a strict decreasing gradual subset, and $\alpha \in (0, 1]$, we have

$$\begin{aligned} \nu \circ \nu(\sigma)(\alpha) &= \tilde{\sigma}(\tilde{\mu}(\sigma))(\alpha) \\ &= \{x | \tilde{\mu}(\sigma)(x) > \alpha\} \\ &= \{x | \text{Inf}\{\beta | x \notin \sigma(\beta)\} > \alpha\}. \end{aligned} \tag{27}$$

If $\text{Inf}\{\beta | x \notin \sigma(\beta)\} = \delta_x$, then $\delta_x > \alpha$; since σ satisfies property (inf-F), then $x \in \sigma(\delta_x) \subseteq \sigma(\alpha)$. Otherwise, if $x \in \sigma(\alpha) = \sigma^d(\alpha) = \cup\{\sigma(\beta) | \beta > \alpha\}$, there exists $\gamma > \alpha$ such that $x \in \sigma(\gamma)$, hence $\text{Inf}\{\beta | x \notin \sigma(\beta)\} > \gamma > \alpha$, and we have the other inclusion.

As a consequence, we have the final result that establishes an isomorphism between the two lattices. See Proposition 3. \square


Corollary 1. *Let X be a set, there is an isomorphism between the lattice of all fuzzy subset of X and the lattice of all strict decreasing gradual subset of X , satisfying property (F), and they preserve arbitrary unions and intersections.*

3.7. A Functorial Interpretation. Let us consider $(0, 1]$ as a category whose objects are the elements of $(0, 1]$ and homomorphisms: only one, $f_{\alpha,\beta}$, from α to β whenever $\alpha \leq \beta$ and the obvious composition.

Let $F: (0, 1] \rightarrow \text{Set}$ a contravariant functor from the category $(0, 1]$ to the category of sets. Indeed, $\{F(\alpha) | \alpha \in (0, 1]\}$ is a directed system with maps $F(f_{\alpha,\beta}): F(\beta) \rightarrow F(\alpha)$, if $\alpha \leq \beta$. Let $D = \varinjlim F$ be the direct limit of this system.

Let us remember the definition of the direct limit, $D = \varinjlim F$. First, we consider the disjoint union, $\dot{\cup} F(\alpha)$, of the family of sets $\{F(\alpha) | \alpha \in (0, 1]\}$ and in it; the equivalence relation R generated by $a \in F(\alpha)$ is related with $b \in F(\beta)$ if either $\alpha \leq \beta$ and $F(f_{\alpha,\beta})(b) = a$ or $\beta \leq \alpha$ and $F(f_{\beta,\alpha})(a) = b$.


Let $p: \dot{\cup} F(\alpha) \rightarrow (\dot{\cup} F(\alpha))/R$ be the canonical projection, $i_\beta: F(\beta) \rightarrow \dot{\cup} F(\alpha)$ be the inclusion, for any $\beta \in (0, 1]$, and $q_\beta = pi_\beta$ the composition.



The pair $(D, \{p_\alpha | \alpha \in (0, 1]\})$ satisfies the corresponding universal property of the direct limit.

In the particular case in which every map $F(f_{\alpha,\beta})$ is injective, then every q_α is also injective; this means that we can consider every $F(\alpha)$ as a subset of D . Therefore, we have that F defines a decreasing gradual subset of $D = \varinjlim F$, or more generally, a decreasing gradual subset of any overset of D and represent it by $(F, \varinjlim F)$.

This allows to give an interpretation of decreasing gradual subsets in terms of contravariant functors. If we start from a decreasing gradual subset of a set X , then $\{\sigma(\alpha) | \alpha \in (0, 1]\}$, together with the family of inclusions, is a directed system, and if $j_\alpha: \sigma(\alpha) \rightarrow X$, for every $\alpha \in (0, 1]$, is the inclusion, then we have a commutative diagram



and an inclusion $h: D \rightarrow X$, from the direct limit D to X , being $h(D) = \text{Im}(h)$ the union of the family of the subsets $\{\sigma(\alpha) | \alpha \in (0, 1]\}$.

Taking into account this construction, we may show that a decreasing gradual subset σ of a set X is nothing more than a contravariant functor $F: (0, 1] \rightarrow \text{Set}$ of this shape, together with an injective map $D = \varinjlim F \rightarrow X$. As a consequence, we may consider contravariant functors as the central element in the study of decreasing gradual subsets.

Hence, we may define a gradual set as a contravariant functor $F: (0, 1] \rightarrow \text{Set}$ and a decreasing gradual set as a

gradual set such that each map $F(f_{\alpha,\beta})$, whenever $\alpha \leq \beta$, is injective.

Given two gradual sets F_1 and F_2 , a map from F_1 to F_2 is just a natural transformation $\theta: F_1 \rightarrow F_2$, i.e., a set of maps $\{\theta_\alpha | \alpha \in (0, 1]\}$ such that each diagram commutes, whenever $\alpha \leq \beta$.

$$\begin{array}{ccc} F_1(\beta) & \xrightarrow{F_1(f_{\alpha,\beta})} & F_1(\alpha) \\ \theta_\beta \downarrow & & \downarrow \theta_\alpha \\ F_2(\beta) & \xrightarrow{F_2(f_{\alpha,\beta})} & F_2(\alpha) \end{array}$$

The contravariant functors from $(0, 1]$ to Set , i.e., the gradual sets, constitute a category that we shall denote by $\text{Set}^{(0,1]}$. The class of all decreasing gradual sets defines a full subcategory of $\text{Set}^{(0,1]}$, and it is closed under (finite and infinite) unions and intersections. Let us call \mathcal{F} this subcategory.

In the subcategory \mathcal{F} , we define an interior operator, $d: F \mapsto F^d$, as follows:

$$\begin{aligned} F^d(\alpha) &= \cup \{F(\gamma) | \gamma > \alpha\} \\ &= \lim_{\rightarrow (\alpha,1]} F(\gamma), \end{aligned} \tag{28}$$

$$\begin{array}{ccccc} F_1(\gamma) & \longrightarrow & \lim_{\rightarrow (\beta,1]} F_1(\gamma) = F_1^d(\beta) & \longrightarrow & \lim_{\rightarrow (0,1]} F_1(\gamma) \\ \theta_\beta \downarrow & & \downarrow \theta^d(\beta) & & \downarrow \\ F_2(\gamma) & \longrightarrow & \lim_{\rightarrow (\beta,1]} F_2(\gamma) = F_2^d(\beta) & \longrightarrow & \lim_{\rightarrow (0,1]} F_2(\gamma) \end{array}$$

A decreasing gradual set F is a strict decreasing gradual set whenever $F = F^d$ and satisfies property (F) if $D = \lim_{\rightarrow} F = \bullet \cup \{F(\alpha) \setminus F^d(\alpha)\}$, where the union is taken in D .

By the relationship between (inf-F) and (F) properties, we may define a fuzzy set as a strict decreasing gradual set satisfying property (inf-F). In particular, strict decreasing gradual sets satisfying property (inf-F) constitute a full subcategory of \mathcal{F} .

As a consequence, we have the following result.

Theorem 3. *Let F be a gradual set.*

- (1) The following statements are equivalent:
 - (a) F is a strict decreasing gradual set
 - (b) The pair $(F, \lim_{\rightarrow} F)$ is a strict decreasing gradual subset of $D = \lim_{\rightarrow} F$
- (2) The following statements are equivalent:
 - (a) F is a strict decreasing gradual set satisfying property (inf-F), i.e., F is a fuzzy set
 - (b) The pair $(F, \lim_{\rightarrow} F)$ is a strict decreasing gradual subset of $D = \lim_{\rightarrow} F$ satisfying property (inf-F)

and if $\alpha \leq \beta$, then there is an inclusion functor $(\beta, 1] \rightarrow (\alpha, 1]$ and a natural map from $F^d(\beta) = \lim_{\rightarrow (\beta,1]} F(\gamma)$ to $F^d(\alpha) = \lim_{\rightarrow (\alpha,1]} F(\gamma)$.

Proposition 4. *Let F be a decreasing gradual set and $\theta: F_1 \rightarrow F_2$ be a decreasing gradual set map. The following statements hold:*

- (1) F^d is a contravariant functor from $(0, 1]$ to Set ; hence, it is a gradual set
- (2) If $\alpha \leq \beta$, the natural map $F^d(\beta) \rightarrow F^d(\alpha)$ is injective; hence, F^d is a decreasing gradual set
- (3) There exists a natural map $\theta^d: F_1^d \rightarrow F_2^d$, defined, for every $\beta \in (0, 1]$, as the only map making commutative the following diagram.
- (4) d defines an endofunctor of \mathcal{F} , which is an interior operator in \mathcal{F} .

Remark 6. The use of decreasing gradual sets allows to avoid the use of decreasing gradual subsets. Indeed, a decreasing gradual subset is, in some sense, more natural: we can build the category of decreasing gradual sets as a subcategory of the functor category $\text{Set}^{(0,1]}$. Otherwise, decreasing gradual subsets are referenced to a set, the same does not happen with decreasing gradual sets; although, as we have the direct limit, the direct system itself acts as a real set. With fuzzy subsets, we have the same situation. Observe that in the fuzzy situation, when we consider the directed systems and the direct limit, we are considering only those elements with a positive, nonzero, membership degree, i.e., we do not consider those with zero membership degree. See also Remark 4.

Remark 7. In looking for an abstract model for gradual subsets of a set X , our first candidate was the functor category $\text{Set}^{(0,1]}$. However, unfortunately, with this category, we do not obtain faithful representation of all gradual subsets. One may consider the gradual subset σ of a nonempty set X defined by $\sigma(\delta) = \begin{cases} X, & \text{if } \delta = (1/2), \\ \emptyset, & \text{if } \delta \neq (1/2). \end{cases}$ Obviously, we can not obtain σ using contravariant functors from the category $(0, 1]$: the reason is that there are no maps from X to \emptyset (it is

not an enriched gradual subset). This have been overcome when we consider enriched gradual sets. This model works perfectly and meets all our expectations whenever we consider decreasing gradual subsets.

Remark 8. Observe that in our construction, we have fixed the categories Set and $(0, 1]$ and considered contravariant functor. If we change contravariant for covariant, we get increasing gradual sets. On the other hand, the category Set has some peculiarities: one is that there \emptyset which is an initial and not a final object; the other is that there are objects A and B such that $\text{Hom}_{\text{Set}}(A, B) = \emptyset$; these force the use of increasing or decreasing gradual sets to assure writing the theory in a functor language using the usual order relation in $(0, 1]$. Some of these restrictions will be removed once we change the category Set for another category as Gr (the category of groups) or $\text{Mod} - A$ (the category of right A -modules).

4. Gradual Subgroups

In Section 2.3, we have studied gradual subsets of $\mathcal{P}(X)$ for any set X and considered in $\mathcal{P}(X)$ the operations:

$$\begin{aligned} (\sigma_1 * \sigma_2)(\alpha) &= \sigma_1(\alpha) * \sigma_2(\alpha), \quad \text{for every } \sigma_1, \sigma_2 \text{ and } \alpha \in (0, 1], \\ \sigma^{-1}(\alpha) &= \sigma(\alpha)^{-1}, \quad \text{for every } \sigma \text{ and } \alpha \in (0, 1]. \end{aligned} \tag{30}$$

Definition 3. Let G be a group (we eliminate the symbol $*$ and represent the product just as juxtaposition); a gradual subgroup of G is a gradual subset σ of G , satisfying

- (1) $\sigma * \sigma \subseteq \sigma$
- (2) $\sigma^{-1} \subseteq \sigma$

Proposition 5. Let e be the neutral element of a group G and σ a gradual subgroup, the following statements hold:

- (1) $e \in \sigma(\alpha)$ for any $\alpha \in (0, 1]$
- (2) $\sigma(\alpha)$ is a subgroup of G for any $\alpha \in (0, 1]$

Therefore, if $S(G)$ is the set of all subgroups of G , a gradual subgroup of G is just a gradual element of $S(G)$.

Proof

- (1) Let $a \in \sigma(\alpha)$, then $a^{-1} \in \sigma(\alpha)$; hence, $e = a * a^{-1} \in \sigma(\alpha)$
- (2) Let $a, b \in \sigma(\alpha)$, then $b^{-1} \in \sigma(\alpha)$; hence, $a * b^{-1} \in \sigma(\alpha)$

If ε is a gradual element of a group G , for any $\alpha \in (0, 1]$, we define $\langle \varepsilon \rangle(\alpha) = \langle \varepsilon(\alpha) \rangle$, the gradual subgroup of G generated by ε . A gradual subgroup σ of G is cyclic if there exists a gradual element ε such that $\sigma = \langle \varepsilon \rangle$.

We may also define finitely generated gradual subgroups: a gradual subgroup σ is finitely generated if there are gradual elements $\varepsilon_1, \dots, \varepsilon_t$ such that for any $\alpha \in L$ we have

intersections and union. We can repeat the same procedure whenever we have a binary operation in X and translate it into $\mathcal{P}(X)$, or a subset of $\mathcal{P}(X)$, in the natural way. Thus, our aim in this section is to study gradual subsets of a given set X , together with an additional algebraic structure in X ; to do that we shall consider the simplest example of groups.

4.1. Gradual Subgroups. Let X be a nonempty set with a binary operation $*$, we define in $\mathcal{P}(X) \setminus \{\emptyset\}$ new binary and unary operations by

$$\begin{aligned} S_1 * S_2 &= \{s_1 * s_2 \mid s_i \in S_i\}, \text{ for every } S_1, S_2 \in \mathcal{P}(X) \setminus \{\emptyset\} \text{ and,} \\ S^{-1} &= \{s^{-1} \mid s \in S\}, \text{ for every } S, \in \mathcal{P}(X) \setminus \{\emptyset\}. \end{aligned} \tag{29}$$

Thus, we may define an operation on gradual subsets of X (for simplicity, in this section, for a set X , a gradual subset of X is a gradual element of $\mathcal{P}(X) \setminus \{\emptyset\}$) by

$\sigma(\alpha) = \langle \varepsilon_1(\alpha), \dots, \varepsilon_t(\alpha) \rangle$. We represent this σ simply as $\langle \varepsilon_1, \dots, \varepsilon_t \rangle$. \square

Proposition 6. Let σ be a gradual subgroup of a group G . The following statements are equivalent:

- (i) σ is finitely generated
- (ii) There exist gradual elements $\varepsilon_1, \dots, \varepsilon_t$ such that $\sigma = \langle \varepsilon_1, \dots, \varepsilon_t \rangle$
- (iii) There exists a positive integer t such that each subgroup $\sigma(\alpha)$ can be generated by t elements

Observe that due to Proposition 5, gradual subgroups of G can be identified with subgroups of the direct product, indexed in $(0, 1]$, of copies of G .

4.2. Normal Gradual Subgroups. By the aforementioned identification, the study of gradual subgroups is very simple. Thus, a gradual subgroup of a group G is normal if for any $\alpha \in (0, 1]$, we have that $\sigma(\alpha) \subseteq G$ is a normal subgroup of G . If σ is a normal gradual subgroup of G , for every $\alpha \in (0, 1]$, we have a quotient group $G/\sigma(\alpha)$.

Let σ be a normal gradual subgroup of G , for every $\alpha \in (0, 1]$, we have a gradual quotient group $G/\sigma(\alpha)$; hence, a gradual quotient set ρ of G is defined as $\rho(\alpha) = G/\sigma(\alpha)$, for every $\alpha \in (0, 1]$. We may represent it also by G/σ . For every $\alpha \in (0, 1]$, there is a group homomorphism $G \rightarrow G/\sigma(\alpha) = (G/\sigma)(\alpha)$.

We define a gradual quotient group of G a gradual quotient set η of G such that for every $\alpha \in (0, 1]$, the projection $p(\alpha): G \rightarrow \eta(\alpha)$ is a group homomorphism.

Proposition 7. Let G be a group, then

- (1) For every normal gradual subgroup σ of G , we have G/σ a gradual quotient group of G
- (2) For every gradual quotient group η of G , there is a normal gradual subgroup κ of G , defined $\kappa(\alpha) = \text{Ker}(G \rightarrow \eta(\alpha))$, for any $\alpha \in (0, 1]$

Lemma 12. Let $f: G \rightarrow G'$ be a group homomorphism.

- (1) For any gradual subgroup σ of G , we have that $f_*\sigma: (0, 1] \rightarrow G'$, defined $f_*(\sigma)(\alpha) = f(\sigma(\alpha))$ as a gradual subgroup of G'
- (2) For any gradual subgroup τ of G' , we have that $f^*\tau: (0, 1] \rightarrow G$, defined $f^*\tau(\alpha) = f^{-1}(\tau(\alpha))$ as a gradual subgroup of G
- (3) If τ is normal, then $f^*\tau$ is normal

Let σ_1, σ_2 be gradual subgroups of G , we define $\sigma_1 \subseteq \sigma_2$ and say σ_1 is a subgroup of σ_2 , if $\sigma_1(\alpha) \subseteq \sigma_2(\alpha)$ for any $\alpha \in (0, 1]$.

Lemma 13. Let σ_1, σ_2 be normal gradual subgroups of G , the following statements are equivalent:

- (i) $\sigma_1 \subseteq \sigma_2$
- (ii) For any $\alpha \in (0, 1]$, there exist group homomorphisms h_α such that the following diagrams commute

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \sigma_1(\alpha) & \longrightarrow & G & \longrightarrow & G/\sigma_1(\alpha) \longrightarrow 0 \\
 & & \downarrow & & \parallel & & \downarrow h_\alpha \\
 1 & \longrightarrow & \sigma_2(\alpha) & \longrightarrow & G & \longrightarrow & G/\sigma_2(\alpha) \longrightarrow 0
 \end{array}$$

Remark 9. In consequence, to include G/σ_2 inside this theory, we could introduce the notion of gradual quotient group of a gradual quotient group G/σ_1 ; hence, study gradual objects which are not related to an ambient group. The same can be done in considering gradual subgroups. Thus, we could introduce the notion of gradual group or enriched gradual group, in a similar way as we did for gradual subsets and sets.

If σ_1 and σ_2 are gradual subgroups of G , we have a gradual subset $\sigma_1\sigma_2$ of G and not necessarily a gradual subgroup; we get a gradual subgroup whenever one of them is normal and, in this case, we have the following.

Lemma 14. Let σ_1, σ_2 be gradual subgroups of G such that σ_1 is normal, then

- (1) $\sigma_1\sigma_2$ is a gradual subgroup of G
- (2) $\sigma_1\sigma_2/\sigma_1$ defined $(\sigma_1\sigma_2/\sigma_1)(\alpha) = \sigma_1(\alpha)\sigma_2(\alpha)/\sigma_1(\alpha)$ is a gradual subgroup of G/σ_1

This theory can be enriched whenever we consider maps between the different $\sigma(\alpha)$'s, i.e., enriched gradual subgroups. For instance, when there is an inclusion, $\sigma(\beta) \subseteq \sigma(\alpha)$ whenever $\alpha \leq \beta$.

4.3. Decreasing Gradual Subgroups. A gradual subgroup σ of G is decreasing if for any $\alpha, \beta \in (0, 1]$ such that $\alpha \leq \beta$, we have $\sigma(\beta) \subseteq \sigma(\alpha)$. For any decreasing gradual subgroup σ of G , for every $\alpha \in (0, 1]$, we have that $\sigma(1) \subseteq \sigma(\alpha)$.

Let σ be a gradual subgroup, we define the accumulation σ^c of σ as

$$\sigma^c(\alpha) = \langle \cup \{ \sigma(\beta) | \beta \geq \alpha \} \rangle, \text{ for any } \alpha \in (0, 1]. \quad (31)$$

It is clear that σ^c is a decreasing gradual subgroup, and a gradual subgroup σ is decreasing if, and only if, $\sigma = \sigma^c$. In particular, we have the following properties of the operator $\sigma \mapsto \sigma^c$.

Lemma 15. Let G be a group, for every gradual subgroups $\sigma, \sigma_1, \sigma_2$, the following statements hold:

- (1) $\sigma \subseteq \sigma^c$
- (2) $\sigma^c = \sigma^{cc}$
- (3) If $\sigma_1 \subseteq \sigma_2$, then $\sigma_1^c \subseteq \sigma_2^c$
- (4) σ^c is the smallest decreasing gradual subgroup containing σ
- (5) If σ is a normal gradual subgroup, then σ^c is a normal subgroup and $\sigma^c(\alpha)$ is the set of all products of elements in $\cup \{ \sigma(\beta) | \beta \geq \alpha \}$ for any $\alpha \in (0, 1]$

Proof. Each element of $\langle \cup \{ \sigma(\beta) | \beta \geq \alpha \} \rangle$ is a product $a_1 * \dots * a_t$, for some $a_i \in \sigma(\beta_i)$, and $\beta_i \geq \alpha$. For any $b \in G$, we have $g * a_1 * \dots * a_t * g^{-1} = (g * a_1 * g^{-1}) * \dots * (g * a_t * g^{-1}) \in \langle \cup \{ \sigma(\beta) | \beta \geq \alpha \} \rangle$.

This means that the map $\sigma \mapsto \sigma^c$ is a closure operator in the set \mathcal{G} of all gradual subgroups of G which is compatible with the product in G . The set of all c -closed gradual subgroups of G is denoted by $\mathcal{F}(G)$. \square

Proposition 8. Let σ_1, σ_2 be gradual subgroups of G , then $\langle \sigma_1\sigma_2 \rangle^c = \langle \sigma_1^c\sigma_2^c \rangle$. In addition, if either σ_1 or σ_2 is normal, then $(\sigma_1\sigma_2)^c = \sigma_1^c\sigma_2^c$.

Proof. In fact, we have that both $\langle \sigma_1\sigma_2 \rangle^c(\alpha)$ and $\langle \sigma_1^c\sigma_2^c \rangle(\alpha)$ are the subgroup generated by the subset $\cup \{ \sigma_1(\beta) \cup \sigma_2(\beta) | \beta \geq \alpha \}$.

Since σ_1 is normal, each element of $(\sigma_1 * \sigma_2)^c(\alpha)$ is a product $a * b$; for $a \in \sigma_1^c(\beta)$ and $b \in \sigma_2^c(\gamma)$, some $\beta, \gamma \geq \alpha$ is formed, then $a * b \in \sigma_1^c(\alpha) * \sigma_2^c(\alpha)$. The converse is similar.

In the same way, we may define the strict accumulation σ^d of σ as

$$\sigma^d(\alpha) = \begin{cases} \sigma(1), & \text{if } \alpha = 1, \\ \langle \cup \{ \sigma(\beta) | \beta > \alpha \} \rangle, & \text{for any } \alpha \in (0, 1). \end{cases} \quad (32)$$

We have that σ^d is a decreasing gradual subgroup and σ^d is normal whenever σ is. Some properties of the operator $\sigma \mapsto \sigma^d$ are the following, whose proof is similar to the proof of Lemma 15 and Proposition 8. \square

Lemma 16. Let G be a group, for any gradual subgroups $\sigma, \sigma_1, \sigma_2$, the following statements hold:

- (1) $\sigma^d \subseteq \sigma^c$.
- (2) If $\sigma_1 \subseteq \sigma_2$, then $\sigma_1^d \subseteq \sigma_2^d$.
- (3) $\sigma^d = \sigma^{d d} = \sigma^{c d} = (\sigma^c)^d$.
- (4) If σ is a normal gradual subgroup, then σ^d is a normal subgroup, and $\sigma^c(\alpha)$ is the set of all products of elements in $\cup \{\sigma(\beta) | \beta > \alpha\}$ for any $\alpha \in (0, 1]$.
- (5) $(\sigma_1 \sigma_2)^d = \langle \sigma_1^d \sigma_2^d \rangle$. In addition, if either σ_1 or σ_2 is normal, then $(\sigma_1 \sigma)^d = \sigma_1^d \sigma^d$.

A gradual subgroup σ is an strict decreasing gradual subgroup whenever $\sigma = \sigma^d$, and we have the following.

Lemma 17. For any gradual subgroup σ , we have that σ^d is the biggest strict decreasing gradual subgroup contained in σ^c .

These results mean that the map $\sigma \mapsto \sigma^d$ is an interior operator in the set $\mathcal{F}^d(X)$ of all decreasing gradual subgroups of G .

4.4. Gradual Subgroups and Fuzzy Subgroups. We shall show that there exists a strong relationship between gradual subgroups of a group G and fuzzy subgroups of G . Remember that if G is a group, a fuzzy subgroup μ of G is a nonconstant, equal to 0, map $\mu: G \rightarrow [0, 1]$ satisfying $\mu(xy^{-1}) \geq \mu(x) \wedge \mu(y)$, for any $x, y \in G$. In particular, if e is the neutral element of G , then $\mu(e) \geq \mu(x)$ and $\mu(x) = \mu(x^{-1})$ for any $x \in G$. Our aim is to identify fuzzy subgroups with some particular decreasing gradual subgroups.

First, we need to realize some modifications to have well-defined gradual groups starting from a fuzzy group.

In the set of all fuzzy subgroups μ of G , we define an equivalence relation: $\mu_1 \sim \mu_2$ if $\mu_1(x) = \mu_2(x)$ for any $x \neq e$. In order to choose a canonical element in each equivalence class, following an idea in [8], for any fuzzy subgroup μ , we define μ^1 as follows: $\mu^1(x) = \begin{cases} \mu(x), & \text{if } x \neq e, \\ 1, & \text{if } x = e. \end{cases}$ Observe that each equivalence class $[\mu]$ has a unique element of the shape μ^1 , whenever μ^1 is a fuzzy subgroup.

Lemma 18. Let μ be a fuzzy subgroup of a group G , then μ^1 is a fuzzy subgroup and μ^1 is normal whenever μ is.

Proof. Let $x, y \in G$, then $\mu^1(xy^{-1}) = \mu(xy^{-1}) \geq \mu(x) \wedge \mu(y) = \mu^1(x) \wedge \mu^1(y)$ whenever $x, y, xy^{-1} \neq e$. On the other hand, if $xy^{-1} = e$, then $\mu^1(xy^{-1}) = 1 \geq \mu^1(x) \wedge \mu^1(y)$; if $x \neq e, y = e$, then $\mu^1(y) = 1 \geq \mu^1(x)$, and we have $\mu^1(xy^{-1}) = \mu^1(x) = \mu^1(x) \wedge \mu^1(y)$.

A decreasing gradual subgroup σ satisfies property (F) if there exists $\text{Max}\{\alpha | x \in \sigma(\alpha)\}$ for every $x \in \cup_\alpha \sigma(\alpha)$. An strict decreasing gradual subgroup σ satisfies property (inf-F) if $\gamma = \text{Inf}\{\beta | x \notin \sigma^d(\beta)\}$ satisfies $x \in \sigma(\gamma)$ for any $x \in G$.

Let σ be a decreasing gradual subgroup; let us denote $\sigma^* (\alpha) = \sigma^c(\alpha) \setminus \sigma^d(\alpha)$, the difference set, for every $\alpha \in (0, 1]$. \square

Lemma 19. Let σ be a decreasing gradual subgroup such that $\sigma^c \neq \sigma^d$; the following statements are equivalent:

- (a) σ^c satisfies property (F)
- (b) σ^d satisfies property (inf-F)
- (c) $\dot{\cup}_{\alpha \in (0,1]} \sigma^*(\alpha) = \cup \{\sigma(\alpha) | \alpha \in (0, 1]\}$

Let $[\mu]$ be the equivalence class of the fuzzy subgroup μ ; we define a gradual subset $\sigma(\mu)$ of G by

$$\sigma(\mu)(\alpha) = \{x \in G | \mu^1(x) \geq \alpha\}. \tag{33}$$

Lemma 20. The map $\nu: [\mu] \mapsto \sigma(\mu)$ is well defined, and $\sigma(\mu)$ is a decreasing gradual subgroup satisfying property (F).

Proof. Observe that for any class $[\mu]$, there exists only one fuzzy subgroup μ such that $\mu = \mu^1$.

Let σ be a decreasing gradual subgroup satisfying property (F); we define a fuzzy subset $\mu(\sigma)$ by

$$\mu(\sigma)(x) = \begin{cases} \text{Max}\{\gamma | x \in \sigma(\gamma)\}, & \text{if } x \neq e, \\ 1, & \text{if } x = e. \end{cases} \tag{34}$$

\square

Lemma 21. With the above notation, $\mu(\sigma)$ is a fuzzy subgroup, and we have a map $\nu: \sigma \mapsto [\mu(\sigma)]$ from the set of all decreasing gradual subsets satisfying property (F) to the set of all classes of fuzzy subgroups.

Now, we have the announced relationship of gradual subgroups and fuzzy subgroups.

Theorem 4. Let G be a group, the maps $\nu: [\mu] \mapsto \sigma(\mu)$ and $v: \sigma \mapsto [\mu(\sigma)]$ define a bijective correspondence between the following:

- (1) Equivalence classes of fuzzy subgroups $[\mu]$ of G
- (2) Descending gradual subgroups σ of G satisfying property (F)

Proof. Let $\mu = \mu^1$ be a fuzzy group, for any $x \in G$, we have

$$\begin{aligned} \mu(\sigma(\mu))(x) &= \text{Max}\{\alpha | x \in \sigma(\mu)(\alpha)\}, \\ &= \text{Max}\{\alpha | \alpha \leq \mu(x)\}, \\ &= \mu(x). \end{aligned} \tag{35}$$

On the other hand, let σ be a decreasing gradual subgroup satisfying property (F), for any $\alpha \in (0, 1]$, we have

$$\begin{aligned} \sigma(\mu(\sigma))(\alpha) &= \{x | \mu(\sigma)(x) \geq \alpha\}, \\ &= \{x | \text{Max}\{\beta | x \in \sigma(\beta)\} \geq \alpha\}, \\ &= \sigma(\alpha). \end{aligned} \tag{36}$$

\square

Lemma 22. Let μ_1, μ_2 fuzzy subgroups; let us define $[\mu_1][\mu_2]$ as $[\mu_1\mu_2]$.

Proof. The product $[\mu_1][\mu_2]$ is well defined. Let $[\mu] = [\mu']$, for any μ_2 , we have

$$\begin{aligned} \mu\mu_2(x) &= \text{Sup}\{\mu(y) \wedge \mu_2(z) | yz = x\}, \\ &= \text{Sup}\{\mu'(y) \wedge \mu_2(z) | yz = x\}, \\ &= \mu'\mu_2(x). \end{aligned} \tag{37}$$

□

Remark 10. Unfortunately, in Theorem 4, the map ν is not a homomorphism with respect to the product of classes of fuzzy subgroups. Indeed, for any μ_1, μ_2 , we have

$$\begin{aligned} (\sigma(\mu_1)\sigma(\mu_2))(\alpha) &= \sigma(\mu_1)(x)\sigma(\mu_2)(\alpha), \\ &= \{x | \mu_1(x) \geq \alpha\} \{x | \mu_2(x) \geq \alpha\}, \\ &\subseteq \{x | \text{Sup}\{\mu_1(y) \wedge \mu_2(z) | yz = x\} \geq \alpha\} \\ &= \{x | (\mu_1\mu_2)(x) \geq \alpha\}, \\ &= \sigma(\mu_1\mu_2)(x). \end{aligned} \tag{38}$$

This inclusion could be strict as the following example shows.

Example 6. We define fuzzy subgroups μ_1 and μ_2 of \mathbb{Z} as follows:

$$\mu_1(x) = \begin{cases} 0, & \text{if } x \in \mathbb{Z} \setminus 2\mathbb{Z}, \\ 1 - \frac{2^t}{3^t}, & \text{if } x \in 2^t\mathbb{Z} \setminus 2^{t+1}\mathbb{Z}, \\ 1, & \text{if } x = 0. \end{cases} \tag{39}$$

$$\mu_2(x) = \begin{cases} 0, & \text{if } x \in \mathbb{Z} \setminus 3\mathbb{Z}, \\ \frac{1}{2} - \frac{1}{3^t}, & \text{if } x \in 3^t\mathbb{Z} \setminus 3^{t+1}\mathbb{Z}, \\ 1, & \text{if } x = 0. \end{cases}$$

We claim $(\mu_1 + \mu_2)(2) = \text{Sup}\{\mu_1(y) \wedge \mu_2(2 - y) | y \in \mathbb{Z}\} \leq (1/2)$. Indeed, we have two possibilities:

- (1) $\mu_1(y) > 1/2$, then $y \in 4\mathbb{Z}$, i.e., there exists $k \in \mathbb{Z}$ such that $y = 4k$. Hence, $\mu_2(2 - y) = \mu_2(2 - 4k) = \mu_2(2(1 - 2k)) < 1/2$ as $2 - y \neq 0$.
- (2) $\mu_1(y) < 1/2$.

In both cases, we have $\mu_1(y) \wedge \mu_2(2 - y) < 1/2$. In addition, we can choose y such that $\mu_1(y) \wedge \mu_2(2 - y)$ is as closed to $1/2$ as we desire. For any $2 \leq t, s \in \mathbb{N}$, there exist $k, h \in \mathbb{Z}$ such that $2^{t-1}k - 3^s h = 1$; hence, $2 - 2^t k = 2(1 - 2^{t-1}k) = 3^s h$; now, if we take $y = 2^t k$, then $\mu_1(y) \geq 1 - (1/3^t)$ and $\mu_2(2 - y) \geq (1/2) - (1/3^t)$. In consequence, $(1/2) > \mu_1(y) \wedge \mu_2(2 - y) \geq (1/2) - (1/3^t)$, which implies that $(\mu_1 + \mu_2)(2) = (1/2)$ and $2 \in (\mu_1 + \mu_2)_{(1/2)}$. On the

other hand, we have $(\mu_1)_{(1/2)} + (\mu_2)_{(1/2)} = 4\mathbb{Z}$ and $2 \notin (\mu_1)_{(1/2)} + (\mu_2)_{(1/2)}$.

We shall change the assignation defined by ν to consider $\tilde{\nu}: [\mu] \mapsto \tilde{\sigma}(\mu)$, in which $\tilde{\sigma}(\mu)$ is a strict decreasing gradual subgroup satisfying property (inf-F) and is defined by

$$\tilde{\sigma}(\mu)(\alpha) = \begin{cases} \{x \in G | \mu^1(x) > \alpha\}, & \text{if } \alpha \neq 1, \\ \{x \in G | \mu^1(x) = 1\}, & \text{if } \alpha = 1, \end{cases} \tag{40}$$

whose inverse is $\tilde{\nu}: \sigma \mapsto [\tilde{\mu}(\sigma)]$, defined as

$$\tilde{\mu}(\sigma)(x) = \begin{cases} \text{Inf}\{\gamma | x \notin \sigma(\gamma)\}, & \text{if } x \neq e, \\ 1, & \text{if } x = e. \end{cases} \tag{41}$$

Thus, we have the following theorem.

Theorem 5. With the above notation, we have

- (1) $\tilde{\sigma}(\mu)$ is a strict decreasing gradual subgroup satisfying property (inf-F), and $\tilde{\nu}$ is well defined.
- (2) $\tilde{\mu}(\sigma)$ is a fuzzy subgroup.
- (3) The maps $\tilde{\nu}$ and $\tilde{\nu}$ define a bijective correspondence between equivalence classes of fuzzy subgroups $[\mu]$ of G and strict descending gradual subgroups σ of G satisfying property (inf-F).
- (4) $\tilde{\nu}$ is a homomorphism with respect to the product of classes of fuzzy subgroups.

Proof

- (1) First, we observe that $\tilde{\sigma}(\mu) = \sigma(\mu)^d$; hence, it is a strict gradual subgroup, and by Lemma 19, it satisfies property (inf-F). It is well defined as $\sigma(\mu)$ is uniquely defined; hence, it is $\tilde{\sigma}(\mu)$.
- (2) It is a direct consequence of Lemma 16.
- (3) We can mimic the proof of Theorem 4.
- (4) For any μ_1, μ_2 , we have

$$\begin{aligned} (\sigma(\mu_1)\sigma(\mu_2))(\alpha) &= \sigma(\mu_1)(x)\sigma(\mu_2)(\alpha), \\ &= \{x | \mu_1(x) > \alpha\} \{x | \mu_2(x) > \alpha\}, \\ &= \{x | \text{Sup}\{\mu_1(y) \wedge \mu_2(z) | yz = x\} > \alpha\}, \\ &= \{x | (\mu_1\mu_2)(x) > \alpha\}, \\ &= \sigma(\mu_1\mu_2)(x). \end{aligned} \tag{42}$$

□

4.5. Normal Fuzzy Subgroups. A fuzzy subgroup μ of a group G is normal if $\mu\eta = \eta\mu$ for any fuzzy subset η or equivalently if $\mu(xy) = \mu(yx)$ for any $x, y \in G$ (see [9]). We are interested in relating normal fuzzy subgroups and normal gradual subgroups. We have defined a gradual subgroup σ to be normal if $\sigma(\alpha) \subseteq G$ is a normal subgroup for any $\alpha \in (0, 1]$.

Lemma 23. Let μ_1, μ_2 be a fuzzy subgroup such that $\mu_1 \sim \mu_2$ and μ_1 is normal, then μ_2 is normal.

Proof. By hypothesis $\mu_1(xy) = \mu_1(yx)$ for every $x, y \in G$, if $xy, yx \neq e$, then $\mu_2(xy) = \mu_2(yx)$. If $xy = e$, then $x = y^{-1}$, hence $yx = e$, and we have $\mu_2(xy) = \mu_2(yx)$.

As a consequence, if μ is a normal fuzzy subgroup, then every fuzzy subgroup in $[\mu]$ is normal; in particular, μ^1 is normal. \square

Theorem 6. *Let μ be a fuzzy subgroup, the following statements are equivalent:*

- (i) μ is normal
- (ii) $\bar{\sigma}(\mu)$ is normal

Proof. We may assume, without loss of generality, that $\mu = \mu^1$. Let $g \in G$ and let us consider the fuzzy subset $\eta(g)$ defined as the characteristic function of $\{g\}$, then

$$\begin{aligned} (\eta(g)\mu)(x) &= \text{Sup}\{\eta(g)(x_1) \wedge \mu(x_2) \mid x = x_1x_2\} \\ &= \eta(g)(g) \wedge \mu(g^{-1}x) \\ &= \mu(g^{-1}x), \end{aligned} \tag{43}$$

and in the same way $(\mu\eta(g))(x) = \mu(xg^{-1})$. Then,

$$\begin{aligned} \bar{\sigma}(\mu) &= \bar{\sigma}(\eta(g^{-1})\mu\eta(g)) \\ &= \bar{\sigma}(\eta(g^{-1}))\bar{\sigma}(\mu)\bar{\sigma}(\eta(g)) \\ &= g^{-1}\bar{\sigma}(\mu)g. \end{aligned} \tag{44}$$

Therefore, $\bar{\sigma}(\mu)$ is normal. Conversely, if $\bar{\sigma}(\mu)$ is normal, for any element $g \in G$, we have $g^{-1}\bar{\sigma}(\mu)g = \bar{\sigma}(\mu)$; hence, $\eta(g^{-1})\mu\eta(g) = \mu$, and μ is a normal fuzzy subgroup. \square

4.6. Gradual Groups. From any decreasing gradual subgroup σ of a group G , we have two families: one $\{\sigma(\alpha) \mid \alpha \in (0, 1]\}$ is a family of groups and the other $\{i_{\alpha,\beta}: \sigma(\beta) \longrightarrow \sigma(\alpha) \mid \alpha \leq \beta\}$ is the family of the inclusions. To include these objects inside a more general theory, we shall consider contravariant functors from $(0, 1]$ to Gr , the category of groups.

For any contravariant functor $F: (0, 1] \rightarrow Gr$ and every $\alpha \leq \beta$, we have now a group homomorphism from $F(\beta)$ to $F(\alpha)$, and the pair $(\{F(\alpha) \mid \alpha \in (0, 1]\}, \{F(f_{\alpha,\beta}) \mid \alpha \leq \beta\})$ is a direct systems of groups and group homomorphisms; hence, there exists its direct limit, say $\lim F$.

We define a gradual group $\overrightarrow{\sigma}$ as a contravariant functor $F: (0, 1] \rightarrow Gr$, and a gradual group homomorphisms from F_1 to F_2 is just a natural transformation from F_1 to F_2 . Therefore, we can consider the category of gradual groups and gradual group homomorphisms, which we denote by \mathcal{G} .

An example of such a gradual group is provided by any decreasing gradual subgroup σ of a group G . In this case, the direct limit $\lim \sigma$ is isomorphic to a subgroup of G ; indeed, it is the union $\overrightarrow{\sigma} \cup \{\sigma(\alpha) \mid \alpha \in (0, 1]\}$.

Following this example, for any arbitrary gradual group F , we say F is a decreasing gradual group whenever each $F(f_{\alpha,\beta})$, for $\alpha \leq \beta$. The class of all decreasing gradual groups is denoted by \mathcal{F} . To well understand the structure of decreasing gradual groups, we build an operator (an endofunctor) d in \mathcal{F} , defined on objects as follows: for any $F \in \mathcal{F}$, we define

$F^d(\alpha) = \lim_{\longrightarrow (\alpha,1]} F(\gamma)$, for every $\alpha \in (0, 1]$. We collect these results in the following proposition, whose proof, after the theory developed in Section 3, is straightforward.

Proposition 9. *Let F be a decreasing gradual group and $\theta: F_1 \longrightarrow F_2$ be a decreasing gradual map. The following statements hold:*

- (1) F^d is a decreasing gradual group
- (2) d is an endofunctor of the full subcategory \mathcal{F} of \mathcal{G}
- (3) d is an interior operator in \mathcal{F} .

A strict decreasing gradual group is a decreasing gradual group F such that $F = F^d$.

At this point, it is convenient to remark that we have gradual groups and gradual subgroups. Contrary to decreasing gradual subgroups, that need of an ambient or a ground group, decreasing gradual groups have it included: it is the direct limit of the direct system that the gradual group defines. This situation allows us to formulate a more attractive category theory of gradual objects which includes the usual constructions of the category of groups. In this context, decreasing gradual groups, strict decreasing gradual groups and fuzzy groups can be identified with adequate subcategories; see the forthcoming paper [10], in which we study gradual and fuzzy modules over a ring.

5. Conclusion

Our goal in this article has been to introduce more general notions than the fuzzy subset in order to find a framework in which to develop a simpler theory that allows testing new techniques and establishing new results in fuzzy theory. In this sense, we start from the concept of gradual element with the goal of introducing gradual subsets. At this point, we establish a bijective correspondence between fuzzy subsets and a particular kind of gradual subsets (strictly decreasing gradual subsets), that satisfies property (inf-F). The more interesting property of this correspondence is that it preserves arbitrary unions and intersections of fuzzy subsets.

In a second degree of abstraction, we consider a gradual subset as a contravariant functor from the category $(0, 1]$ to the category of sets, which allows us to define the notion of fuzzy and gradual sets without the use of an ambient set. Thus, we have three degrees of abstraction, the first one corresponds to fuzzy subset; the second one to gradual subsets, identifying fuzzy subsets as some particular gradual subsets; and the third one to contravariant functors from $(0, 1]$ to the category of sets, or directed systems of sets, identifying decreasing gradual subsets as those systems with injective maps. Observe that in each abstraction level, we have the objects studied in the previous one. We also establish the corresponding theory for groups in two different but compatible ways (1) defining contravariant functors to the category of groups; $Gr^{(0,1]}$, and (2) defining groups in the functor category $\text{Set}^{(0,1]}$.

One of the goals of this paper was to find a framework to study together the two crisp sets associated with each fuzzy set, and we have proven that groups and gradual groups

allow it to do so. On the other hand, the use of category theory tools will allow to extend this working method to other structures, of which the sets and groups studied are only an example.

Data Availability

In this article, we include all the data that support this research. The references included at the end of the article can be used to complement partial aspects of these data.

Disclosure

An older version has been submitted as arxiv in Cornell University according to the following link: <https://export.arxiv.org/abs/1812.07521>. This is an updated version of that manuscript.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

This work was partially supported by the Research Group “FQM-266-Anillos y Módulos.” This work was supported by grant A-FQM-394-UGR20 from Programa Operativo FEDER 2014–2020 and Consejería de Economía, Conocimiento, Empresas y Universidad de la Junta de Andalucía (Spain) and by the “María de Maeztu” Excellence Unit IMAG, reference CEX2020-001105-M, funded by MCIN/AEI/10.13039/501100011033/.

References

- [1] L. A. Zadeh, “Fuzzy sets,” *Information and Control*, vol. 8, no. 3, pp. 338–353, 1965.
- [2] D. Dubois and H. Prade, “Gradual elements in a fuzzy set,” *Soft Computing*, vol. 12, pp. 165–175, 2008.
- [3] J. Fortin, D. Dubois, and H. Fargier, “Gradual numbers and their application to fuzzy interval analysis,” *IEEE Transactions on Fuzzy Systems*, vol. 16, no. 2, pp. 388–402, 2008.
- [4] R. Boukezzoula, S. Galichet, L. Foulloy, and M. Elmasry, “Extended gradual interval arithmetic and its application to gradual weighted averages,” *Fuzzy Sets and Systems*, vol. 257, pp. 67–84, 2014.
- [5] D. Sánchez, M. Delgado, M. A. Vila, and J. Chamorro-Martínez, “On a non-nested level-based representation of fuzziness,” *Fuzzy Sets and Systems*, vol. 192, pp. 159–175, 2012.
- [6] T. Martin, “The $X-\mu$ representation of fuzzy sets,” *Soft Computing*, vol. 19, pp. 1497–1509, 2015.
- [7] T. Martin and B. Azvine, “The $X-\mu$ approach: Fuzzy quantities, fuzzy arithmetic and fuzzy association rules,” in *Proceedings of the 2013 IEEE Symposium on Foundations of Computational Intelligence (FOCI)*, pp. 24–29, Singapore, April 2013.
- [8] I. Jahan, “The lattice of L -ideals of a ring is modular,” *Fuzzy Sets and Systems*, vol. 199, pp. 121–129, 2012.
- [9] J. Mordeson and D. Malik, *Fuzzy Commutative Algebra*, World Scientific, Singapore, 1998.
- [10] J. M. Garcia and P. Jara, “Gradual and fuzzy modules. Functor categories,” *Functor categories*, p. 26, University of Granada, Granada, Spain, 2021.
- [11] S. Nădăbana, “Some fundamental properties of fuzzy linear relations between vector spaces,” *Filomat*, vol. 30, no. 1, pp. 41–53, 2016, <http://www.jstor.org/stable/24898409>.
- [12] A. Piegat and M. Plucinski, “Fuzzy number addition with the application of horizontal membership functions,” *The Scientific World Journal*, vol. 2015, Article ID 367214, 16 pages, 2015.
- [13] G. Wang and J. Li, “Approximations of fuzzy numbers by step type fuzzy numbers,” *Fuzzy Sets and Systems*, vol. 310, pp. 47–59, 2017.
- [14] D. Lewis and T. Martin, “ $X-\mu$ Fuzzy Association Rule Method,” in *Proceedings of the 2013 13th UK Workshop on Computational Intelligence (UKCI)*, pp. 144–150, Guildford, UK, September 2013.
- [15] F. Mora-Camino and C. A. Nunes Cosenza, “Fuzzy Dual Number. Theory And Applications,” *Studies in Fuzziness and Soft Computing*, Springer, Berlin, Heidelberg, Germany, 2018.